



HAL
open science

Recovering bone and tooth proteins in arid environments using palaeoproteomics -A case study from the Late Stone Age site of Toteng, Botswana

Louise Le Meillour, Séverine Zirah, Joséphine Lesur, Matthieu Lebon, David Pleurdeau, Antoine Zazzo

► To cite this version:

Louise Le Meillour, Séverine Zirah, Joséphine Lesur, Matthieu Lebon, David Pleurdeau, et al.. Recovering bone and tooth proteins in arid environments using palaeoproteomics -A case study from the Late Stone Age site of Toteng, Botswana. Scientific Symposium Frontiers in Heritage Science, Feb 2019, Paris, France. 10, pp.181 - 186, 2019, 10.13140/RG.2.2.21813.12008 . hal-02551368

HAL Id: hal-02551368

<https://hal.science/hal-02551368>

Submitted on 22 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - ShareAlike 4.0 International License

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/331122690>

Recovering bone and tooth proteins in arid environments using palaeoproteomics – A case study from the Late Stone Age site of Toteng, Botswana

Poster · February 2019

DOI: 10.13140/RG.2.2.21813.12008

CITATIONS

0

READS

148

6 authors, including:



Louise Le Meillour

Muséum National d'Histoire Naturelle

6 PUBLICATIONS 4 CITATIONS

[SEE PROFILE](#)



Séverine Zirah

Muséum National d'Histoire Naturelle

106 PUBLICATIONS 2,499 CITATIONS

[SEE PROFILE](#)



Joséphine Lesur

Muséum National d'Histoire Naturelle

88 PUBLICATIONS 603 CITATIONS

[SEE PROFILE](#)



Matthieu Lebon

Muséum National d'Histoire Naturelle

81 PUBLICATIONS 940 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



ANR PaleoCet [View project](#)



Origin and diffusion of domestic animals in Africa [View project](#)

Recovering bone and tooth proteins in arid environments using palaeoproteomics

A case study in the Late Stone Age site of Toteng, Botswana



Louise LE MEILLOUR^{a,b,*}, Séverine ZIRAH^b, Joséphine LESUR^a, Matthieu LEBON^c, David PLEURDEAU^c & Antoine ZAZZO^a

^a Unité Archéozoologie, Archéobotanique : Sociétés, Pratiques et Environnements (AASPE), MNHN, CNRS, Paris, France.

^b Unité Molécules de Communication et Adaptations des Microorganismes (MCAM), MNHN, CNRS, Paris, France

^c Unité Histoire naturelle de l'Homme Préhistorique (HNHP), MNHN, CNRS, Paris, France

* llemeillour@mnhn.fr



The advent of herding practices remains a key event in human societies in terms of social, cultural, economical changes and regarding their interactions with animals.

In Africa, no remain of wild ancestors species of sheep *Ovis aries* nor goat *Capra hircus* has never been reported. This suggests that those domestic species have been introduced from the epicentre of their domestication : the Near East^[1].

The first occurrences of those animals remains in the archaeological record are attested since the 7th millenium BCE in Egypt and Lybia^[2].

But their appearance in the Southern part of the continent is only dated to the 2nd century BCE [3].

The modalities of their diffusion across the continent in such large amount of time is still debated : cultural diffusion (percolation) or population migrations; Eastern or Western route?

Regarding the strong evidences of extant exchanges between Eastern and Southern Africa, we propose to document the hypothesis of the Eastern route using biomolecular approaches through a larger project (Fig. 1). Here, we will focus on methodological developments of palaeoproteomics in the context of African remains.



Fig. 1: Archaeological sites included to the PhD project

Because of the strong morphological similarities between the caprines species and with other small wild bovids, emphasized by the African burying conditions, the distinction of archaeological remains is not always possible.

We aimed to use a novel biomolecular method for species determination of archaeological material: palaeoproteomics (Fig. 2).

Based on the amino acid variations on preserved proteins, this method allow species identification of remains where DNA might not be conserved, due to increased diagenetic processes amplified by arid environmental conditions.




Fig. 2: General principle of palaeoproteomics analyses on archaeological material

and examined by one of the authors (JL). Most of them were too fragmented and altered to propose morphological identifications beyond family^[8].

Estimation of remains organic phase preservation

In order to estimate the organic content of Toteng remains, 0.5 mg of powdered sample were applied on the diamond of a Attenuated Total Reflectance Fourier transform infra-red spectrometer. Anvil pressure was adjusted to normalise $\nu_3\text{PO}_4$ band to an absorbance of 0.5 (Fig. 3).

The ratio between amide I and PO_4 bands area is then used to estimate nitrogen and collagen weight percent content (%wt) according to a method described by Lebon and collaborators^[9].




Fig. 3: ATR FT-IR spectrum of a modern bone reference (*B. taurus*). Areas used for organic content preservation estimation appear in grey.

METHODS

Paleoproteomics

Because no chemical preparation protocol was never adapted to remains from arid environments, we aimed at identifying one that allow the best protein sequence coverage and permits species identifications (Fig. 4). Once proteins were properly extracted and enzymatic hydrolysis succeeded, we performed high performance liquid chromatography coupled to tandem mass spectrometry analyses for characterization^[9].




Fig. 4: Flow chart of palaeoproteomics protocol conducted in this study

Context of the PhD

Aim

Material & Methods

Results

Abstract

Conclusion

A collagen content threshold of ~ 3 wt% under which it would be risky to perform palaeoproteomics analyses is proposed.

Tris-EDTA buffer is identified as the best demineralizing agent when dealing with remains from arid environment.

The presence of domestic sheep *Ovis aries* at the Late Stone Age site of Toteng, Botswana, is confirmed based on preserved proteins from one of the degraded splinters.

In Africa, the absence of wild ancestors' remains of domestic caprines indicates an exogenous origin of domestic sheep and goat. Archaeological evidences suggest that they were introduced in the Northern part of the continent by the 7th millenium BCE, but only arrived in the Southern part by the 2nd century BCE. Moreover, the morphological similarities existing between the two species among them, and with other small wild bovids like antelopes combined with the large amount of highly fragmented remains due to arid environmental conditions have substantially limited the use of classical comparative anatomy analyses.

We report here the use of a promising method, palaeoproteomics, on degraded bone and tooth splinters from the Late Stone Age site of Toteng, Botswana. This study is among the first to focus on African archaeological remains, strongly degraded by arid burying conditions and diagenesis. The purpose of our work is to both document the presence of domestic caprines at this site, but also to develop a new methodology adapted to archaeological remains from arid environments.

BEST SUITED PROTOCOL FOR REMAINS FROM ARID ENVIRONMENTS

All 18 Toteng remains presented contrasted organic phase preservation. Nine out of the 18 were selected for palaeoproteomics analyses regarding a range of preservation distributed between 1.91% and 5.38%^[8] of collagen content.




Fig. 5: Number of assigned peptides using available databases per sample and per decalcification protocol used. Preserved collagen content (coll. wt%) are presented above each sample.

Within the nine tested remains for palaeoproteomics analyses, four yielded preserved peptides that were assigned to proteins using available databases (Fig. 5). Identified remaining proteins were $\alpha 1$ and $\alpha 2$ chains of collagen type I. Tris-EDTA buffer allowed the best protein sequence coverage.

SPECIES IDENTIFICATION BASED ON PROTEIN CHARACTERISATION





Fig. 6: MS/MS spectra of family and species markers of alpha 1 chain of collagen type I preserved in Toteng samples and identified in this study.

$\alpha 1$ chain of collagen type I allowed the refining of species identification. One remain presented the suidae marker (Fig. 6, top left), two other the equidae marker (top right) and one remain was attributed to the domestic caprine species, *Ovis aries* (bottom center)^[8].

All the $\alpha 1$ chain of collagen type I markers presented here were never reported in the literature and will allow species identification based on the least changing chain of this type of collagen.

REFERENCES

- [1] VIGNE, J.-D., ET AL., 2012, *Ethnozootechnie* 91, 11-19.
 [2] LESUR, J., 2017, *Presses universitaires du Midi*, 205 p.
 [3] PLEURDEAU D., ET AL., 2012, *PLoS ONE*, 7, e40340.
 [4] CAMPBELL A. C., 1992, 671-677.
 [5] ROBBINS L., ET AL., 1998, *South Afr. Arch. Bull.*, 53, 125-132.
 [6] SADR K., 1997, *Current Anthropology*, 38, 104-112.

This work was supported by a grant from the ANR under the LabEx ANR-10-LABX-0003-BCDiv, in the program "Investissements d'avenir" ANR-11-IDEX-0004-02 and an ATM grant from the Muséum National d'Histoire Naturelle under the name "ProtéArch". The authors would like to thank the National Museum of Botswana for access to the Toteng remains and expressly M. Mvimi for obtaining the exporting permit, without whom this study will not have been possible. The authors also want to thank the Plateau Technique de Spectrométrie de Masse Bio-Organique and the Plateau de Spectrométrie Infrarouge of the MNHN for accessing the instruments and X. Gallet for his help for FTIR analyses. Finally, the author would like to thank the organizers of the 8th Bone Diagenesis Meeting for provoking the opportunity of publishing in a Special Issue of the *Paleo3* journal.

ACKNOWLEDGEMENTS