



HAL
open science

Emergence of regional cultural traditions during the Lower Palaeolithic: the case of Frosinone-Ceprano basin (Central Italy) at the MIS 11–10 transition

Marie-Hélène Moncel, Italo Biddittu, Giorgio Manzi, Barbara Saracino, Alison Pereira, Sébastien Nomade, Christine Hertler, Pierre Voinchet, Jean-Jacques Bahain

► To cite this version:

Marie-Hélène Moncel, Italo Biddittu, Giorgio Manzi, Barbara Saracino, Alison Pereira, et al.. Emergence of regional cultural traditions during the Lower Palaeolithic: the case of Frosinone-Ceprano basin (Central Italy) at the MIS 11–10 transition. *Archaeological and Anthropological Sciences*, 2020, 12 (8), 10.1007/s12520-020-01150-x . hal-02950771

HAL Id: hal-02950771

<https://hal.science/hal-02950771>

Submitted on 6 Nov 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Emergence of regional cultural traditions during the Lower Paleolithic: the**
2 **case of Frosinone-Ceprano basin (Central Italy) at the MIS 11-10 transition**

3 Marie-Hélène Moncel (1), Italo Biddittu (2), Giorgio Manzi (2, 3), Barbara Saracino (2), Alison Pereira
4 (4), Sébastien Nomade (5), Christine Hertler (6), Pierre Voinchet (1), Jean-Jacques Bahain (1)

5 (1) UMR 7194 HNHP, CNRS-MNHN-UPVD, Department « Homme et environnement », National
6 Museum of Natural History, Institut de Paléontologie Humaine, 1 rue René Panhard, Paris
7 (France)

8 (2) Istituto Italiano di Paleontologia Umana, Via Ulisse Aldrovandi 18, Roma (Italy)

9 (3) Dipartimento di Biologia Ambientale, Sapienza Università di Roma, Roma (Italy)

10 (4) École française de Rome, Piazza Farnese, Rome (Italy)

11 (5) UMR 8212 CEA Saclay, LSCE, Gif-sur-Yvette (France)

12 (6) Senckenberg Forschungsinstitut, Frankfurt/Main (Germany)

13

14 **Abstract**

15 Decades of fieldwork in the Frosinone-Ceprano basin (Latin Valley, Latium, central Italy) have shed
16 light on numerous open-air Lower Palaeolithic localities, delivering a human fossil calvarium,
17 thousands of scattered faunal remains and a large collection of lithic industries, including core-and-
18 flake type lithic series (Mode 1) and Acheulean assemblages (Mode 2).

19 The continuously growing number of available geochronological data (obtained by $^{40}\text{Ar}/^{39}\text{Ar}$ on
20 volcanic minerals, ESR/U-series on large mammal teeth and ESR on bleached fluvial quartz) allow
21 today the construction of a reliable and precise chronological framework for the Lower Palaeolithic
22 sites of this area of the Latin valley. The archaeological horizons with bifaces all appear to belong to a
23 relatively short Middle Pleistocene time range, between about 410 and 350 ka, coeval to the end of
24 the interglacial MIS 11 and to the beginning of the following glacial MIS 10.

25 The Acheulean tools are often associated with cores and flakes. Bifaces are mainly made on
26 limestone, secondary flint and quartz. The archaeological corpus also yielded tools on fragments of
27 large herbivore bones. Comparisons between technological strategies and paleo-anthropological
28 data at the global scale are now meaningful and enable us to decipher hominin behaviour at a more
29 regional scale. Such careful work is becoming essential in the frame of the recent discoveries showing
30 that the MIS 11-10 period was a pivotal period characterized by the appearance of several new
31 archaeological features later associated with the Neanderthal lineage in Western Europe. We present
32 here the first in-depth technological study of the Acheulean lithic corpus from the major
33 archaeological sites from the Frosinone-Ceprano basin including the Campogrande localities (CG9
34 and CG10, intermediate and upper levels), Colle Avarone, Selvotta, Isoletta (level 4), Lademagne
35 (upper and lower levels) and Masseria Castellone.

1 For this work, we focus on biface shaping strategies and demonstrate technological features
2 suggesting the existence of networks connecting the different human occupation sites. Technological
3 data are compared to other penecontemporaneous Italian sites to discuss the hypotheses and
4 characteristics of such early evidence of regionalization in Europe in this specific area of Central Italy.
5 It seems indicate that glacial conditions characterized by millennial rapid climatic oscillations could
6 have been favourable to the development of specific vegetation propitious to human settlement in
7 South-western Europe. European vegetation, as it drives the biomass availability for large herbivores,
8 seems hence have played crucial role in the mobility and settlement of human groups.

9

10 **Key-words**

11 Italy; Ceprano basin; Acheulean; Marine Isotopic Stages 11-10; bifaces; regionalization

12

13 **Introduction**

14 From the hominin evolution perspective, the period ranging between 450 and 350 ka, corresponding
15 to the interglacial/glacial cycle constituted by Marine Isotopic Stages (MIS) 11 and 10, stands out in
16 Western Europe as a crucial period. This time threshold, associated in Europe to the Lower
17 Palaeolithic (and the Acheulean culture) and to the transition towards the Middle Palaeolithic, is still
18 under-investigated and only few studies discussing the reasons and modes of the appearance and
19 transmission of Middle Palaeolithic innovations in Western Europe have been published despite the
20 importance of this period to understand the human settlement of the continent (Moncel et al.,
21 2020a).

22 The Acheulean culture is commonly linked to the ability to produce bifaces. Other behaviours are
23 punctually evidenced such as the management of the raw material territory, the production of large
24 flakes for the LCTs and some innovations such as the independency of the stone shape for the
25 debitage (*i.e* Moncel et al., 2015, 2020b). Western European archaeological assemblages show that
26 the MIS 11 interglacial was a threshold period associated with early evidence of regionalization of
27 cultural traditions after the intense MIS 12 glacial period, considered as a major climatic crisis for
28 hominins and fauna in this area (Moncel et al., 2017, Pereira et al., 2018). After this glacial phase, we
29 observe an increase in the number of well-preserved occupation levels and the appearance of new
30 subsistence behaviours (hunting organization, specialization on one or some species of large
31 mammals, seasonal occupations depending on the location of the site and land use patterns) and
32 technical innovations (appearance of core technologies, increase in light-duty tools, fire
33 domestication) in Western Mediterranean and more especially in Italy (Villa et al., 2016, Pereira et
34 al., 2017, 2018). These behavioural changes and innovations suggest, by their nature and diversity,
35 an increase in cognitive capacities of the hominins illustrated by new skills and social interactions

1 with site networks. Furthermore, the beginnings of the control of fire could have contributed to the
2 development of behavioural abilities and to the extension and organization of territories. However,
3 the reasons underlying these behavioural changes, pushing back the roots of the Neanderthal world,
4 have yet to be identified. (1) Was this behavioural evolution rooted in previous traditions? (2) Was it
5 due to the arrival(s) of new populations or to the on-site evolution of Neanderthals from *Homo*
6 *heidelbergensis* or another Middle Pleistocene hominin ancestor? (3) Was there a population
7 increase conducive to early regional networks of sites and the diffusion of innovations? (4) Was it
8 due to local environmental adaptation to climatic changes? (Moncel et al., 2018).

9 The last hypothesis is supported by the available climatic and environmental reconstructions. Indeed,
10 MIS 12 is known as one of the most arid glacial periods of the Quaternary (Shackleton, 1987, Tzedakis
11 et al., 2003, Masson-Delmotte et al., 2010, Rohling et al., 2014). In contrast, the subsequent MIS 11
12 interglacial was exceptionally long (more than 20 ka, whereas typical interglacial duration is ~13 ka,
13 Tzedakis et al., 2012). Such a long-lasting interglacial period after a harsh glacial stage could have
14 been favourable to the development of vegetation propitious to animal and human settlement in
15 Western Europe. European vegetation, as it drives the biomass availability for large herbivores,
16 affects the mobility of animal and human groups and it is therefore crucial for this point of view.

17 In order to better understand the impact of the MIS 11-10 period on human settlements in Central
18 Italy, we have studied lithic assemblages recovered from sites located in the Frosinone-Ceprano
19 (Pofi) basin (Latin Valley, Latium, Central Italy). This area is renowned for the discovery in 1994 of a
20 fragmented human calvarium, firstly erroneously attributed to *Homo erectus* and now relied to
21 *Homo heidelbergensis* (Ascenzi et al. 1996, 2000; Manzi et al., 2001; 2010; Manzi, 2012, 2016; Di
22 Vincenzo et al., 2017). Initially considered as dated around 900-800 ka, this human fossil was recently
23 re-dated by a multidisciplinary approach, including magnetostratigraphic and $^{40}\text{Ar}/^{39}\text{Ar}$ analyses (e.g.,
24 Nomade et al., 2011; Biddittu et al., 2019). The refined chronology indicates that humans probably
25 lived in Central Italy during the cold period of marine isotope stage (MIS) 12 and the subsequent
26 warm period of MIS 11. Despite its archaic morphological features and lack of Neanderthal traits, the
27 Ceprano calvarium is therefore contemporaneous with derived hominins such as the ancestors of
28 Neanderthals (Manzi et al. 2010; Manzi, 2016).

29 Decades of surveys in the Frosinone-Ceprano basin have yielded lithic industries in numerous
30 localities dispersed in a relatively small area. The series are limited in number but sufficient to
31 compare the technological processes between localities and to discuss possible connections between
32 sites. The geochronological and palaeo-environmental framework available for Central Italy permits
33 then to discuss the hypothesis of networks of sites and investigate a possible early cyclic process of
34 regionalization during prehistoric times, much earlier than those highlighted for late Neanderthals
35 during MIS 4-3 (*i.e.*, Burke et al, 2008; Deujard and Moncel, 2010; Faivre et al 2017).

- 1 Several issues motivate the analysis of the Frosinone-Ceprano lithic series:
- 2 -What are the modalities of biface production and core technologies during the MIS 11/10 timescale
- 3 in the basin?
- 4 -Are there common technological features and differences in biface production evidenced in all the
- 5 studied localities?
- 6 -Do these technological features provide evidences of traditions and/or regionalization in the Italian
- 7 peninsula and more generally in Southern Europe during the MIS 11/10 timescale?
- 8 -If common technological strategies exist, can we assume that the localities are part of a network of
- 9 nearby sites in a small area?
- 10 -How can we explain the differences in technological strategies and toolkits (presence or not of Large
- 11 Cutting Tools-LCTs)? Could they be due to the location or functions of the sites, or eventually to the
- 12 diversity of traditions inside the basin (palimpsest of networks)?

13

14 **The archaeological data**

15 ***General archaeological and geological context***

16 The archaeological sites investigated in our study are located in the Latin Valley (Latium, Central

17 Apennines), on the Tyrrhenian side of the Apennines chain (**Fig. 1**). This valley is geographically

18 enclosed between mountains to the northeast (Simbruini-Ernici mounts) and southwest (Lepini-

19 Ausoni-Aurunci carbonatic mounts), and volcanic complexes to the west (Colli Albani) and southeast

20 (Roccamonfina) (Marra et al., 2009; Rouchon et al., 2008). The tectonic depression of the Latin Valley

21 consists of several extensional basins formed because of the evolution of the Late Pliocene-Middle

22 Pleistocene Apennines normal fault system (Carrara et al., 1995; Boari et al., 2009). The Frosinone-

23 Ceprano basin, in which all the archaeological sites investigated in this study (Campogrande, Colle

24 Avarone, Selvotta, Isoletta, Lademagne and Masseria Castellone) are located, represent along with

25 the Anagni basin the main sedimentary basins of the valley. Due to the intense volcanic activity

26 occurred during the Middle Pleistocene along the Tyrrhenian marge, volcanic effluents (tephra layers

27 or free volcanic minerals) were regularly deposited in the sedimentary sequences recorded in these

28 basins, including archaeological ones. The main volcanic sources of these effluents are mainly located

29 along the Tyrrhenian marge (Colli Albani, Roccamonfina, Sabatini and Vico volcanic districts), but

30 include also the local Middle Latin valley volcanic field complex (MLVVF), sometimes called

31 Ernici/Volschi cones, located in the heart of the Latin Valley (Civetta et al., 1979; Karner and Renne,

32 1998; Giannetti, 2001; Rouchon et al., 2008; Marra et al., 2009, Pereira et al., submitted). This

33 complex consists of about twenty monogenetic cones, the most active of which being the Pofi and

34 Ceccano edifices (Basilone et Civetta, 1975; Civetta et al., 1979, see **Fig. 1**). All these volcanic edifices

1 were very active during MIS 11 and MIS 10, causing direct impacts on the various environments
2 bearing the Frosinone-Ceprano basin human occupations. All the archaeological sites presented in
3 this contribution are coeval with this local volcanic activity from a narrow time range of about 50 ka
4 (c.a. 410-350 ka), spanning from the MIS 11 optimum to the MIS 10 glacial onset (see **Fig. 2**, Manzi,
5 2016; Pereira et al., 2018).

6 From a climatic point of view, the MIS 11- MIS 10 period was, in Central Italy and more largely in the
7 Mediterranean basin, suitable to the development and subsistence of hominin populations. Indeed,
8 the MIS 11 was one of the warmest and longest (20-30 kyrs) Quaternary interglacials (Karner and
9 Marra, 2003) and the MIS 10 Glacial stage was characterized in the Mediterranean basin by
10 mild/temperate conditions followed at the end by a brief period more drought. Several
11 palaeoclimatic and palaeoenvironmental records of the Mediterranean area (e.g. sea surface
12 temperature records, Girone et al., 2013; Lake Ohrid and Tenaghi Philippon pollen diagrams, Tzedakis
13 et al., 2006, Sadori et al., 2016, Wagner et al., 2019) suggest indeed rather favourable climatic
14 conditions during MIS 10 with no significant reduction of the arboreal taxa. In the neighbour Boiano
15 basin (Molise), pollen's assemblage studies led by Orain et al., (2013) have demonstrated the
16 persistence of high edaphic humidity during the MIS 11-10 period. In this study the authors
17 suggested that these specific conditions may have privileged the existence of refuge area for arboreal
18 taxa and thus could have provided subsistence resources and had a significant effect on the
19 demography of human populations (Orain et al., 2013, Moncel and Schreve in 2016). It is also worth
20 noticing that palynological analyses of the sedimentary unit stratigraphically located just below the
21 layer where was found the Ceprano calvarium in the Campogrande archaeological area have been
22 recently provided (Margari et al., 2018). These two sedimentary layers are separated by an erosional
23 surface attributed by the authors to the glacial stage MIS 12. The pollen assemblage of this layer is
24 dominated by the arboreal species *Corylus*, *Carpinus betulus*, deciduous *Quercus* and *Abies*. These
25 analyses have allowed the authors to identify two forested phases. Based on biostratigraphical and
26 sedimentological data this sequence has been attributed to the MIS 13 interglacial and by analogy
27 the authors encouraged a MIS 11 age for the archaeological levels of Campogrande. All these climatic
28 specific conditions of the Mediterranean basin may also have contributed to the good preservation
29 of archaeological sites in Central Italy during this period. This time range is also coeval to a significant
30 renewal of the faunal groups takes. The post- MIS 10 period corresponds to the transition between
31 the Galerian and Aurelian Large Mammal's periods (Milli and Palombo, 2005, Marra et al., 2018). It
32 should be noticed that the most part of the paleontological assemblages of the sites investigated in
33 this study are all attributed to the final Galerian.

34

35 ***Campogrande, Colle Avarone and Selvotta localities***

1 The stratigraphic sequence of the Frosinone-Ceprano basin comprises an extensive lacustrine
2 formation, called the Lirino lake sequence, mainly recorded in the South-eastern part of the basin
3 (Devoto, 1965; Carrara et al., 1995; Centamore et al., 2007). This lacustrine formation is oriented
4 NW-SE (from Ceprano to Cassino-Pignataro Interamna) and reached its maximum extension (35 km)
5 during the Middle Pleistocene.

6 The deposits are characterized by a total maximum thickness of about 100 m and were divided into
7 two main geological units, LR1 at the base and LR2 at the top (Devoto, 1965; Carrara et al., 1995;
8 Centamore et al., 2007, Centamore and Dramis, 2010, Manzi et al., 2010). Most of the Lirino
9 formation is made up of fluvio-lacustrine and volcanic derived deposits. The final phase of the Lirino
10 basin infilling (top of LR2), associated to lacustrine facies, was dated by Nomade et al. (2011) to
11 about 353 ± 4 ka (2σ analytical uncertainties). This final phase of the Lirino sequence (LR2) was
12 identified in the sequences of the investigated sites.

13 Similar stratigraphic contexts have been evidenced by I. Biddittu in 1974 on the Campogrande, Colle
14 Avarone and Selvotta outcrops that are mainly constituted by gravels, sands and sandy-clays.
15 Quaternary deposits present a thickness of about twenty meters in these three localities (between
16 125 and 102 m a.s.l.) and lay in unconformity above Messinian fluvio-lacustrine deposits (Biddittu,
17 1974). Regarding the function of the sites where fauna and artefacts are found are generally at a
18 comparable altitude and in levels of sand and small gravels, which suggest the presence of vast
19 territories close to fluvial and peri-lacustrine formations. These areas seem to have been explored by
20 hominins periodically, particularly along rivers. These hominines would thus have carried out
21 different types of activities related to the exploitation of plant and animal resources.

22

23

24 *Campogrande localities _ CG 9 (CG9A = 41°31'48.6" 13°28' 54.1" and CG9B = 41°31'49.9" 13°28'*
25 *54.0") and CG 10 (CG10A = 41°31'32.2" 13°28' 54.5", CG10B = 41°31'31.1" 13°28' 53.3" and CG10C =*
26 *41°31'29.7" 13°28' 57.1")*

27 The Campogrande archaeological area (120-115 m a.s.l.) is located 3 km SE of the city of Ceprano (see
28 **Fig. 1**). Palaeontological and archaeological remains were found in twelve localities of this area,
29 named from CG 1 to CG 12 (CG for Campogrande), some of which containing more than one
30 archaeological level. CG 1 corresponds to the pioneering site, where the Ceprano calvarium was
31 found (Ascenzi et al., 1996, 2000; Manzi et al., 2001, 2010; Manzi, 2012, 2016; Di Vincenzo et al.,
32 2017).

33 Two excavations were carried out at CG9 between 2001 and 2006 (Biddittu et al., 2019). They started
34 from a surface located about 3 m deep with respect to the ancient excavating level associated to a

1 paleo-channel constituted by gravels and sands. In this excavating area were found fauna fossils and
2 Acheulean pieces. All the materials studied in this contribution were yielded from both these
3 excavation campaigns and previous surveys. This level attests to the presence of waterways that
4 have drained paleo-surfaces that contained both the remains of faunas and the lithic artefact.
5 According to these new excavations data and previous ones, it appears that the all archaeological
6 levels identified at Campogrande are restricted into the end of the Lirino paleo-lacustrine sequence,
7 in the northern part of this paleolake (Manzi, 2016).

8 The geological layers at Campogrande are rich in volcanic material originating from neighbouring
9 volcanoes and in 2011, Nomade et al. obtained an $^{40}\text{Ar}/^{39}\text{Ar}$ age of 353 ± 4 ka (2σ analytical
10 uncertainties) for a clayey geological layer rich in K-feldspars, stratigraphically located above the
11 palaeo-anthropological horizon. This chronological constraint thus providing a *terminus ante quem*
12 age to the human fossil (see **Fig. 1, 2**). Among the twelve Campogrande archaeological localities, two
13 have delivered an abundant archaeological assemblage: CG 9 and CG 10 (Manzi et al., 2016). Most of
14 the archaeological material recovered at CG 9 and CG 10 is on display at the Pietro Fedele Museum
15 at Pofi (Frosinone province, Latium) (Biddittu et al., 2019). The simplified stratigraphic log of
16 Campogrande sequence, with the positions of the CG 9 and CG 10 archaeological levels, is shown in
17 **Fig. 2** (modified from Pereira et al., 2018). At CG 9, only one significant archaeological level was
18 found in the upper part of the section (CG 9 upper), whereas at CG 10, three different human
19 occupations levels were identified (CG 10 lower, CG 10 intermediate and CG 10 upper). Between
20 2001 and 2006, fieldwork investigations were conducted at CG 9, resulting in the enrichment of the
21 upper CG 9 archaeological corpus (Margari et al., 2018). Acheulean industries are recovered in level
22 CG 9 upper, CG 10 intermediate and CG 10 upper. Few lithic tools were found in intermediate CG 10
23 and lower CG 10 is only associated with core-and-flake industries. These archaeological horizons
24 (excepted lower CG 10) are stratigraphically located above the layer dated in Nomade et al., 2011,
25 suggesting a maximum age of 353 ± 4 ka (2σ analytical uncertainties) for the archaeo-surfaces.

26 *Colle Avarone (41° 32' 10.44"N 3° 28' 38.22"E)*

27 Concerning both Colle Avarone and Selvotta it is worth noticing that findings made over the years
28 came for most of them, from archaeological excavations, while others were collected in a small
29 agricultural plot where the archaeological surface constituted by sands and gravels layer was clearly
30 evidenced. Colle Avarone archaeological site (about 120 m a.s.l.), very close to the Campogrande
31 localities, is located in the lower valley of the Sacco River, a few kilometres upstream of the
32 confluence with the Liri River (Biddittu, 1974, see **Fig. 1**). The locality was discovered in 1973 by Italo
33 Biddittu in a quarry ("pozzolana cava") which no longer exists.

34 A small excavating area conducted in a sedimentary level constituted by sands and gravels have been
35 opened in 1975 by I. Biddittu. This level was identified on a small gravel pit exploited at that time at

1 the base of a section of about 4 m high in front of an older quarry (Colle Avarone in the original
2 definition was perhaps Colle Cavarone for the presence of quarries). From this small essay came out
3 a bone "dagger" made from an elephant rib whose length was determined by an intentional fracture
4 facilitated by a notch performed transversely in the bone. This essay is about 15 m from the point
5 where the recent sampling was done on the levels alongside the road.

6 One main archaeological level, containing Acheulean and bone industries, was identified (Biddittu,
7 1974). From a sedimentological and geological point of view, the archaeological remains were
8 discovered into fluvio-lacustrine deposits rich in gravels and K-rich volcanic minerals (leucites and
9 potassium feldspars) corresponding to the final infilling of the Lirino paleolake basin. A simplified
10 stratigraphic log of Colle Avarone is presented in **Fig. 2** and further stratigraphic information can be
11 found in Biddittu, 1974 and Biddittu and Segre, 1984. The archaeological level found below a
12 paleosol, is mainly constituted by gravels and altered leucites. Biface tools, usually symmetric, are
13 dominant in the lithic assemblage (Biddittu and Segre, 1984). The palaeontological assemblage
14 mainly comprises Middle Pleistocene large mammal species (or taxa) (*Palaeoloxodon antiquus*, *Sus*
15 *scrofa*, *Equus caballus*, *Bos primigenius*, *Cervus elaphus*), avifauna (*Stercorarius longicaudatus*, *Anser*
16 *brachyrhynchus*) and turtles (*Testudo* sp.). The birds and the occurrence of *Cuon* sp. remains suggest,
17 according to Biddittu and Segre (1984), a cold environment during the human occupation level.

18

19 *Selvotta* ((41° 32' 10.44"N 3° 28' 38.22"E)

20 Selvotta locality is directly adjoined to the section of Colle Avarone (about 120 m a.s.l., Biddittu,
21 1974, see **Fig. 1**). So far, no scientific publication has been devoted to this archaeological locality.
22 Only one Acheulean archaeological level has been brought to light and the most part of the lithic
23 tools were surface collected. Both lithic and bone tools were recovered at this site. Italo Biddittu
24 proposed a geological correlation between the archaeological level of Selvotta and the
25 archaeological level of Colle Avarone (CA and the archaeological levels CG 9 and CG 10 upper from
26 the Campogrande localities (see Fig. 2). All these levels are characteristic fluvio-lacustrine deposits,
27 with gravels and volcanic minerals, of the Lirino paleo-lake that occurred during the MIS 11/MIS 10
28 transition period (Nomade et al., 2011; Biddittu et al., 2019).

29

30 ⁴⁰Ar/³⁹Ar dating on single K-feldspar crystals extracted from Selvotta and Colle Avarone sandstones
31 are currently in progress to confirm or infirm the stratigraphic attributions and correlations.

32

33 ***Isoletta, Lademagne and Masseria Castellone localities***

34 *Isoletta* (41°31'44.86"N, 13°34'0.16"E)

1 The Isoletta archaeological site (~105 m a.s.l., Biddittu, 1974, 2004; Pereira et al., 2018), discovered
2 in the 1970s, is located at the confluence between the Sacco and Liri rivers only a few kilometres SE
3 of Ceprano city. A large excavating area was conducted in 1998 and 1999 and was focused on the
4 Acheulean level (Isoletta L4) and the inferior sedimentary levels only. The stratigraphic description of
5 the site was established during this excavation following the high-speed train railway construction,
6 which that helped exposed a 30-m-thick section (Biddittu, 2004). The sequence, which is no longer
7 accessible, is mainly composed of alluvial deposits including sands, gravels and volcanic material
8 related to the final lacustrine phase of the Lirino basin infilling (see **Fig. 2**). Two archaeological levels
9 were evidenced in the Isoletta sequence but only the uppermost level presents Acheulean industries
10 (Level 4, labelled GA6Z in Pereira et al., 2018 (Biddittu, 2004)). Palaeontological remains found in
11 level 4 are associated with the Galerian large mammal assemblage (Biddittu, 1974; Milli and
12 Palombo, 2005).

13 Rare mode 1 artefacts and remains of mammals, birds, amphibians and fish were found in the lowest
14 archaeological level (Pereira et al., 2018). A tibia of elephant was found in this level, it was
15 intentionally fractured by the hominids. The lower part of the sequence is more lacustrine, very rich
16 in silt, and has delivered abundant molluscs and vegetation debris.

17

18 *Lademagne (41°31'16.68"N, 13°35'0.72"E)*

19 The Lademagne site, discovered in 1965 (105 m a.s.l.), was found in a very active tectonic area in the
20 North-West part of the Lirino lake sedimentary basin. It is only located 5 km west of Isoletta near the
21 city of San Giovanni Incarico (Biddittu et al., 2012, see **Fig. 1**). Most of the Lademagne sequence is
22 made up of clays, sands, silts, gravels and reworked volcanic deposits corresponding to an alluvial
23 terrace. Gastropods found in level 3 (*Planorbis* sp., *Valvata cristata*, *Pisidium* sp.) indicate a lacustrine
24 habitat (Comerci et al., 2015). At Lademagne, two archaeological layers separated by 5 m of
25 thickness of sediments were identified with very similar archaeological and palaeontological
26 assemblages (Biddittu et al., 2012). They are comprised in gravels and sands sedimentary levels
27 (Biddittu et al., 2012). Unlike at Isoletta, Acheulean and bone tools were found in both levels. The
28 Lademagne palaeontological assemblage is associated with the Galerian Large Mammal Age (Milli
29 and Palombo, 2005; Biddittu et al., 2012) and the upper level assemblage includes *Palaeoloxodon*
30 *antiquus*, *Hippopotamus* sp., *Dicerorhinus* sp., *Dama clactoniana*, *Equus ferus*, *Cervus elaphus*,
31 *Castor* sp. and some remains of *Bos* sp., *Equus* sp. and *Ursus* sp. (Biddittu et al., 2012).

32

33 *Masseria Castellone locality (41°31'12.69"N, 13°35'28.87"E)*

34 Masseria Castellone is located at 400 m from the Lademagne archaeological site (see **Fig. 1**) and only
35 a river ditch separates the two sites. The lithic tools (only a few bifaces in limestone) were discovered

1 in a geological horizon similar to the “Lademagne upper” archaeological layer, rich in sands and
2 gravels (**Fig. 2**) and found at the same altitude (i.e., 105 m a.s.l.). It could suggest that they belong to
3 the same sedimentary unit and are therefore of the same age. So far, no data have been published
4 on this archaeological horizon.

5 From a general point of view, the overall areas where fauna and artefacts were found in the studied
6 zone are generally at a comparable altitude and in levels of sand and small gravels, which suggest the
7 presence of vast territories close to fluvial and peri-lacustrine formations during a period of intense
8 volcanic activity between 410 and 340 ka according to all available $^{40}\text{Ar}/^{39}\text{Ar}$ constraints. These areas
9 seem to have been explored by hominins periodically, particularly along rivers. These hominines
10 would thus have carried out different types of activities related to the exploitation of plant and
11 animal resources. Traces of human activity on faunal remains are mainly represented by the
12 presence of frequent bone artefacts that are for some of them not straightforward. Bone tools were
13 recognized both in the upper level (Selvotta, Colle Avarone, Campogrande 9A, Isoletta L4,
14 Lademagne Upper level) and in the lower level (Isoletta ESR1, Lademagne Lower Level). The faunal
15 remains are characterized by alteration surfaces due to exposure and in rare cases fluctuation
16 especially in the smallest fragments (common aspect also observed in in situ levels with slow water
17 circulation). The presence of artefacts on bone is a characteristic of all the Acheulean sites of
18 southern Lazio from Fontana Ranuccio to Pontecorvo- Aquino

19 **Material and method**

20

21 **Material**

22 The aforementioned geochronological multi-method study ($^{40}\text{Ar}/^{39}\text{Ar}$ on single grains, ESR on
23 bleached quartz and ESR/U-series) at Isoletta and Lademagne (Pereira et al., 2018, Voinchet et al.,
24 2019) and the stratigraphic correlations proposed considering the geological similarities between the
25 deposits (facies, altitude of the levels, proximity) to link these two localities with the other studied
26 sites in the Frosinone-Ceprano basin constrained the age of the Acheulean archaeological levels of
27 these localities between 410 and 370 ka, *i.e.* as contemporaneous of the period ranging from MIS 11
28 interglacial optimum to the MIS 10 glacial onset.

29 Two groups of localities with artefacts can be distinguished: (1) the localities on which were
30 evidenced flakes, cores and pebble-tools (for instance Campogrande CG 10 lower, or Isoletta lower)
31 (**Fig. 3**), and (2) the localities demonstrating a core technology and some Large Cutting Tools (LCTs)
32 including bifaces (Campogrande 10 intermediate, Campogrande 10 upper, Campogrande 9
33 intermediate and upper, Colle Avarone, Selvotta, Isoletta L4, Lademagne lower and upper and
34 Masseria Castellone) (**Table 1**).

1 Regarding the series for which no bifaces were found, the heavy-duty component is composed of
2 pebble-tools or chopper-cores, convergent or not, on thick limestone, quartz and quartzite
3 pebbles/cobbles. Flakes may come from biface production or flaking. Cores are on thick and large
4 quadrangular limestone pebbles, but also on small flint pebbles/nodules. The debitage is not
5 intensive. These cores bear a few short, deep and hinged removal scars, on orthogonal or multiple
6 surfaces adapted to the shape of the blank. There are also some unifacial or bifacial cores with
7 asymmetrical flaking surfaces.

8 Eleven lithic series containing LCTs were also studied. Part of them corresponds to incomplete
9 archaeological series recovered from excavations and from surface collecting and therefore
10 submitted to taphonomic processes (as is usual for open-air Palaeolithic sites), but these series were
11 nonetheless located quite precisely in the same stratigraphic sequence and considered relevant for a
12 comparative study of reduction processes (Biddittu et al., 2019). The cutting edges of the artefacts do
13 not show any signs of damage due to high-level disturbances. When the pieces are partially covered
14 by breccia or when the scars are smoothed by chemical processes, the technological study provided
15 was only partial (**Table 1**). The artefacts in flint are better preserved than the ones in limestone. The
16 cutting edges are usually fresh suggesting that the lithic tools were not much transported. The
17 alteration on limestone tools seems post-depositional and *in situ*. Small tools, for which flint was
18 preferably used, are rare because the drainage of running water tends to disperse small objects. The
19 larger bifaces remain more concentrated.

20

21 **Method**

22 In this work, we studied (according to Kleindienst, 1961; Leakey, 1971; Boëda et al., 1990; Geneste,
23 1991; Texier and Roche, 1995; Roche, 2005; Goren-Inbar and Sharon, 2006; McPherron, 2006; Key et
24 al., 2016; Lycett et al., 2016; McNabb et al., 2018; Torre de la and Mora, 2018):

25 (1) The mode of debitage on cores by analysing the organization and management of the blank and
26 the technological characteristics of the end-products (including types of retouch) ;

27 (2) The different modalities of biface production at the final shaping phase for the LCTs: we focused
28 our work on studying the geometry and identification of the blank and determining when possible,
29 the number, size and organization of the removals as well as the number of series of the removals,
30 extension and organization of shaping on both faces, secondary retouch and location, shape of
31 cutting edges, type of tip management.

32

33 We applied classical statistical methods including box-plots to compare the tool sizes between the
34 investigated localities. To estimate the internal variability between lithic assemblages, we used the
35 Multiple Correspondence Analysis with the SYSTAT software based on a list of quantitative and

1 qualitative parameters. A list of relevant parameters was selected (length, width and thickness in
2 mm, raw materials, extension of removals on each face, number of series of removals, retouch on
3 the edges, tip shaping and bilateral and bifacial symmetry). All the tables and graphs represent 80%
4 of the variability of the data with two components.

5

6 **Results: technological analyses of the lithic component for each locality**

7

8 ***Campogrande (CG) 9 intermediate (or B)***

9

10 In this series three bifaces were identified (see **Table 1**). These tools in flint and limestone may
11 possibly be on flakes (with the platform residue as a proximal back) (**Fig. 4**). Shaping covers a large
12 part of one surface and the periphery, with more removals on the extremity.

13

14 ***Campogrande (CG) 9 upper (or A)***

15 This series consists of one bifacial and thick core and 11 limestone bifaces characterised by varied
16 lengths (80 to 180 mm long) (see **Table. 1**):

17 Two bifaces are short completely or partially worked, with a symmetrical cross-section, made by
18 series of large and short removals. The tip of one of them is curved. The blank may be a flake. Three
19 tools are plano-convex bifaces shaped on half of the piece, sometimes with a back (on a flake?), with
20 a pointed or round tip and retouch on the two cutting edges.

21 Then six tools are plano-convex bifacial tools. The flatter surface is shaped by large removals which
22 thin the whole face, while the opposite convex face is managed by series of removals limited to the
23 upper half and the periphery of the tool.

24

25 ***Campogrande (CG) 10 intermediate (or B).***

26 Apart from some cores on small flint nodules (less than 50 mm long) for the production of a few
27 small flakes, this series is composed of a heavy-duty and a light-duty component.

28 The light-duty component consists of a micro-biface on a limestone pebble, a micro convergent
29 pebble-tool and a point on a micro-flint nodule (30 to 50 mm long). They are minimally shaped by a
30 few unifacial and bifacial removals, which do not really modify the form of the blank.

31 The heavy-duty tools are composed of a “rabot”, a thick scraper on a limestone split pebble, and
32 three limestone bifaces. One of these bifaces is minimally shaped with a thick base. Shaping covers
33 the whole volume with numerous and abrupt removals. The cutting edges are sinuous with no final
34 retouch. Another one is an asymmetrical elongated biface in limestone, with shaping covering most

1 of the surfaces and with final retouch on the edges. Finally, the last one that is in limestone is
2 symmetric and made on a half-pebble. Shaping is not very invasive.

3

4 ***Campogrande (CG) 10 upper (or A)***

5 A dozen pieces are cores and/or chopper-cores (**Fig. 5**). They are made on small oval and
6 quadrangular pebbles (in flint and limestone) of poor-quality (50 mm long) (50-100 mm). The
7 debitage systematically follows the natural morphology of the blank. It is orthogonal or unifacial
8 unipolar (on the largest face or on the core edge). The number of removals is limited and the
9 products are irregular and frequently hinged. We did not evidence any preliminary preparation of the
10 core. The platforms are usually cortical or use previous removals.

11 The series totals 22 heavy-duty tools on limestone. Flint nodules are too small to make heavy-duty
12 tools. Three categories of heavy-duty tools are described:

13 -The first one corresponds to a thick pointed scraper (rabot-type) made by deep and large removals
14 on a small part of the periphery of a broken limestone pebble.

15 -The second category is composed of five elongated plano-convex bifaces on badly preserved
16 limestone (**Fig. 6**). The mean length of these bifaces is around 200 mm, making them among the
17 longest of the series. Some cortical residues suggest that the blank could be a flat and elongated
18 cobble. The bifacial volume is globally managed by one series of invasive removals on both faces.
19 Some removals are hinged. The shaping is alternate or done face by face when visible. The sinuous
20 edges are not rectified by retouch. The base of the tool is shaped by convergent removals (forming a
21 thin cutting edge), or left unworked (back or natural round cortical base). The tip is always thin, and
22 the shaping of the tip by small removals is part of general shaping. The maximum thickness of the
23 tool is on the proximal or the middle part of the biface.

24 -The third category is constituted by sixteen partial bifacial tools in limestone. This category is more
25 diversified than the previous ones, and includes short and elongated pieces (from 110 to 210 mm).
26 The blank is a flake for most of the tools (**Fig. 7, 8**) and the shaping mainly concerns the upper part of
27 the tool, covering the entire surface or only the periphery. The flatter face is either covered by large
28 removals or short removals. The opposite surface indicates series of semi-abrupt frequently hinged
29 removals, rarely rectified by final series of short removals and retouch. In some cases, only one edge
30 is worked, and the tip and cutting edge are opposite a back or a cortical side. The cutting edges are
31 consequently sinuous. The pointed or round tip is made by small or large removals related to the
32 shaping process of the lateral cutting edge(s). The base is always cortical, with or without a back. The
33 cross-section is plano-convex. One piece shows a longitudinal "coup de tranchet" while another piece
34 shows some transversal removals on the transversal tip (due to use or resharpening?).

35

1 Most of the tools are asymmetrical bifacial tools on limestone for which the blank appears to play a
2 major role, with minimal shaping on flakes and more invasive shaping on elongated pebbles. All the
3 tools are made by a single series of removals with no final retouch, producing sinuous edges. The
4 main feature is the presence of a back on some pieces, for elongated bifacial pointed tools and
5 backed bifacial tools.

6

7 ***Colle Avarone***

8 This locality yielded 11 cores in flint and limestone (see Table. 1) made on small nodules or pebbles.
9 The cores are bifacial centripetal, orthogonal, unipolar or unifacial or bifacial chopper-cores with
10 abrupt removals. There is no preparation of the core. One core is attributed to the Levallois core
11 technology by the organization of the removals (see Moncel et al., 2020).

12 Regarding the LCTs, Colle Avarone yielded the largest series (n = 87). These tools in limestone,
13 excepting one on flint, were more or less well-preserved.

14 One LCT was partial and irregular on a flake, made by some removals on the edges and the apical
15 extremity. Its tip is pointed.

16 Three short bifaces were evidenced. Their bases are thick and large and their cross-section are
17 symmetrical. Shaping covers the total volume with series of large and short frequently hinged
18 removals. Final retouch is punctual on cutting edges more or less sinuous. . The tip is round or
19 quadrangular and the base is shaped by abrupt removals.

20 Nine elongated asymmetrical (plano-convex) and totally shaped biface (n = 9) were found. Shaping
21 manages the general volume with large removals on both faces, shorter removals on the base and
22 the round tip. Cortical patches are preserved on the thick base. On the tip, removals are frequently
23 transversal. For one piece, the tip is nose-shaped. Final retouch often finishes working the periphery
24 of the tool and does not modify the sinusoidal edges. The thickest part of the tool is in the middle of
25 the elongated bifaces.

26 Seven elongated symmetrical totally shaped bifaces (**Fig. 9, 10**) are made by large invasive removals
27 that shape the global volume, without final retouch, except for one tool on a flake. The edges are
28 sinusoidal. The round or pointed tip is made by transverse removals except for one tool with a “coup
29 de tranchet”.

30 Five short bifaces with a thick and cortical base are triangular tools made by total shaping on a flake
31 by invasive and short removals on one face, shorter on the opposite flat face with final retouch near
32 the tip. The cross-section is plano-convex or symmetrical.

33 Seven partial bifacial tools on flat and elongated pebbles conducive to minimal management, a
34 plano-convex cross-section and sinuous edges with hinged removals. Two types of management
35 were observed: (1) on the upper part and covering the surfaces, (2) peripheral on part of both sides.

1 The thin tip is mainly pointed with series of short removals. Cortical patches are invasive, on the base
2 and sometimes with a cortical back (pebble side). The maximum thickness is on the middle or
3 proximal part.

4 Sixteen partial bifacial tools on a flake including 14 in limestone and 2 in flint. In the same way as for
5 pebbles, shaping is peripheral or covers the upper part. The cross-section depends on the shape of
6 the flake where the flat surface is more or less covered by flat removals. Removals on the convex
7 surface are hinged. The base is thick, sometimes with a back. The edges are sinuous. The tip is
8 pointed or round and formed by short removals and some retouch. The two flint tools show a similar
9 shaping with many small removals on the tip on the longest tool, one edge is retouched, associating a
10 bifacial point and a lateral scraper.

11 Eighteen partial elongated bifaces on possible pebble-blocks show an independent shaping on both
12 edges and the tip, made by alternate, peripheral or partial deep removals, without final retouch. The
13 thin tip is sometimes shaped by more removals (transversal or lateral). The cross-section is
14 asymmetrical, sometimes twisted, with a back or a trihedral section.

15 Twenty-nine partial elongated and triangular bifaces on a flake on limestone and one flint (**Fig. 8, 9**)
16 show a minimal shaping with short removals located on the upper half of the flake with frequent
17 shaping of the thin tip, retouched or not, and both edges. Invasive removals cover one face more
18 than the other one. The cross-section is asymmetrical with a proximal thick and cortical back or a
19 shaped thick base. The edges are rectilinear or sinusoidal. In some cases, final retouch covers one
20 whole edge, creating what looks like a lateral scraper. The work is thinner on the tip of the single flint
21 biface.

22 Ten elongated bifaces with total shaping in limestone, asymmetrical ($n = 7$) or symmetrical in cross-
23 section ($n = 3$) are shaped by series of short and invasive removals. The edges are sinuous or
24 rectilinear. The cross-section is asymmetrical and some of the bifaces are backed. The tip is round or
25 pointed, sometimes curved. The middle part of the tool is the thickest.

26 Eight bifacial pieces with a transversal cutting edge are on limestone except for one on flint. They are
27 on flakes shaped by peripheral and short removals, which are more invasive on one face. The
28 transversal apical edge is made by transversal and lateral removals without final retouch. One tool
29 shows a "coup de tranchet". The base is thick, sometimes with a back (the preserved cortical butt of
30 the flake). The cross-section is asymmetrical.

31

32 If we summarize the series (**Fig. 10**), we can observe three main categories of tools. One is composed
33 of elongated tools with a cortical back or created by removals, often on flakes, a plano-convex cross-
34 section, made by large and flat removals on one face and semi-abrupt removals on the opposite face.
35 The second category is composed of triangular, elongated or short, partial tools with invasive

1 removals and retouch despite sinusoidal edges. Finally, the last category is made of crudely shaped
2 tools.

3
4

5 ***Selvotta***

6 The series is composed of 38 limestone and quartzite tools, and two flint tools. The heavy-duty tools
7 are composed of one “rabot” and a thick scraper on a half limestone pebble. A backed biface on a
8 flake looks like a bifacial scraper. LCTs can be divided into three main groups. Two trihedral pieces
9 are on limestone and totally shaped by deep and invasive alternate removals with a thick base and a
10 thin tip. The cutting edges are sinuous with no final retouch. The cross-section is symmetrical or
11 asymmetrical. Fifteen plano-convex with total management of the tool including two in flint (**Fig. 11**)
12 are elongated bifaces (150-200 mm long), mostly worked by large then shorter removals, then
13 retouched on the edges. The tip is pointed or round. The cross-section is plano-convex with a thicker
14 base. Sixteen plano-convex bifaces in limestone and quartzite show a management of the two cutting
15 edges and the tip and peripheral and partial shaping of large flakes (cutting edges and tip) (**Fig. 11,**
16 **12**).

17 To summarize, most of the assemblage is composed of plano-convex bifaces with partial
18 management of the two cutting edges and the tip, or total management of the volume. For the tools
19 with a whole management, they are shaped by series of invasive and short face by face removals and
20 the cutting edges are sometimes rectified by final and partial retouch on the convex face. They are
21 elongated, triangular or oval. The tip is mainly pointed and thin and is prepared by small removals or
22 retouch. The thickest part of the tool is the middle part. The two flint tools can be categorized as
23 whole bifacial tools. They are oval or pointed, elongated, plano-convex, and completely shaped with
24 general management of the bifacial volume. For the partial tools, when the blank is still visible, half
25 pebbles, flat pebbles or large flakes are shaped by peripheral removals. The tool base is often left
26 unworked. The tip is pointed worked by small removals or retouch and its management is an
27 extension of cutting edge shaping.

28

29 ***Isoletta L4 (GA6Z)***

30 The 13 bifaces are in limestone. They are elongated, although partially or totally shaped in different
31 ways, by deep and large removals and rarely by final retouch (**Fig. 13**). The maximum thickness of the
32 tool is on the proximal half or in the tool centre.

33 Five technological groups can be distinguished based on asymmetrical or symmetrical cross-sections.
34 Four tools are made by large removals covering a part of the piece with no final retouch. The tip is
35 pointed and narrowed by deep removals. The cutting edges are sinusoidal. The maximum thickness is

1 at the base of the tool. Five tools are made by large and short removals on the whole volume with
2 rare final retouches. The tip is narrowed by removals and the base is also worked and thinned. The
3 maximum thickness is at the centre of the tool. One asymmetrical tool is made by large removals on
4 the upper part With a tip thinned and round. Two tool made on a flat pebble are partial with large
5 removals on the two cutting edges and the tip. Finally, a large flake is partially shaped and some final
6 retouch.

7

8 ***Lademagne lower***

9 The series is composed of a dozen cores (polyhedral cores on large blocks with no preparation and
10 unifacial cores), some chopper-cores and choppers on limestone and flint. The presence of a flake
11 that can be described as a Levallois flake is noteworthy (**Fig. 14**). There are also 14 bifacial pieces, 2
12 with a transversal cutting edge, 5 with partial management and eight with general bifacial
13 management. The tools are on poor-quality limestone, except for one on flint (**Fig. 15, 16**). The two
14 tools with a transversal cutting edge are on limestone and are completely bifacial and symmetrical
15 with invasive removals The partial bifacial tools are mainly on flakes and plano-convex with sinuous
16 edges. Removals cover either the upper part, or the periphery of the flake. The tip is thin and curved
17 while the base is thick and generally left unworked or only slightly worked. Bifaces with general
18 bifacial management are the predominant category. All these tools are in limestone, except for one
19 in flint. They are elongated or short, made by several series of removals covering the whole surfaces.
20 The cross-section is plano-convex for the thin tip and sometimes symmetrical for the thick base. Final
21 retouch is sometimes visible on the whole periphery while the edges remain sinuous. The shorter
22 bifaces show a round or transversal cutting edge while the longer ones have a carefully worked
23 pointed tip. The shaping mode seems to be alternate despite a plano-convex cross-section. The flint
24 biface is the most worked tool of the series and final retouch rectifies the edges.

25

26 ***Lademagne upper***

27 Debitage activity is on limestone cores on pebbles. Core technology is unifacial, orthogonal or on
28 multiple surfaces. Large flakes are the main blanks used for bifacial technology when determinable.
29 We can mention one *déjeté* flake 15 cm long with limited unifacial shaping of the two convergent
30 edges.

31 Apart from one flint tool, the bifaces are in limestone of varying quality (accidents occurred during
32 the production for three pieces). These hinged scars are due to the bad quality of some limestone
33 but also to technical reasons (too flat face of the tool and bad angles of the cutting edges).

34 Two groups can be distinguished. There thirteen partial bifacial tools , mainly on flakes or broken or
35 flat pebbles, with preserved cortical backs (**Fig. 17**), which could account for the curved aspect of the

1 longitudinal section of the tool. The tip is managed by some bifacial removals. One or both edges of
2 the tools are shaped by short removals producing a plano-convex cross-section without final retouch.
3 Edge angles vary between 75 and 85°. A lateral or proximal back is preserved (butt of the flake?), and
4 the tools look like backed bifacial scrapers. Twelve elongated or short bifaces show a general
5 management of the volume and a final retouch. The cross-section is symmetrical (alternate shaping)
6 or plano-convex (face by face shaping) (**Fig. 17**). The angles of the cutting edges are inferior to those
7 of the previous group, between 51 and 72°. The shaping mode could be linked to the type of blank
8 (flake or pebble). The base is sometimes unworked, with a back (part of a flake?) or is part of the
9 overall bifacial management. The thickest part of the tools is the central area. The tip is pointed or
10 round, made by general tool management removals, by one large transverse removal or small
11 retouch.

12

13 ***Masseria Castellone***

14 The series is composed of six tools, few more or less elongated in limestone, and of varied
15 dimensions. They are worked by large and deep and/or short alternate removals limited to the upper
16 part of the blank and almost all show a symmetrical cross-section. Although only part of the tool is
17 shaped, we observe general management of the bifacial volume. These tools are one asymmetrical
18 bifacial “dejeté” scraper, one elongated and symmetrical tool (limande-type), one symmetrical tool
19 on a flake with a proximal back, one curved and symmetrical tool shaped on the whole periphery by
20 short removals with a proximal back (**Fig. 18**), one asymmetrical partial tool on a flat pebble, and
21 finally one symmetrical triangular and partial tool.

22

23 **Discussion**

24 Before discussing the tools themselves, it seems essential to give some general information
25 concerning what we know so far about the origin and nature of raw material used. A constant
26 characteristic of the various sites is that the stone tools recovered are made on medium-sized and
27 small limestone gravels. Pebbles are mainly smaller than the mean size of the bifaces. Raw materials
28 used to elaborate the bifaces were probably collected in adjacent sites to those of the frequentation,
29 even if no specific selection is evidenced currently. The presence of some types of flakes
30 demonstrates the probability of the use of the flaking on anvil despite the lack of large pebbles
31 necessary for this technique. Tests carried using pebbles collected in the bed of current active rivers
32 in the area has shown that only a very limited number of limestone pebbles have the necessary
33 compactness and quality require to be used. The field observations also suggest that the bifaces
34 were made in other unknown neighbouring areas. However, all the raw materials used are products

1 of the erosion of the Cretaceous formations derived from the mountains that surround the Sacco and
2 Liri valleys.

3

4 ***Common features and differences between the lithic series in the localities of the Ceprano basin***

5 Acheulean culture in the Frosinone area is more frequent than the Middle Paleolithic that is
6 represented by sporadic objects except for some rare sites. The lithic series of the Ceprano-Frosinone
7 basin are mainly manufactured on fine-grained, generally of poor-quality limestone that are
8 sporadically associated with better quality flint, quartzite and quartz. Some bifaces are shaped on
9 large flakes obtained from cobbles and were rarely abandoned at the localities, excepted in the
10 upper archaeological level of Lademagne. Raw material types differ for core technology and shaping.
11 Limestone is the preferred one for the LCTs and heavy-duty component whereas flint and quartz are
12 mainly used for flaking, although a few LCTs are made on flint and quartzite in the Campogrande and
13 Lademagne localities. This is probably due to the size of the available nodules and pebbles, as flint is
14 generally available as small nodules.

15 Core technologies demonstrate common technological rules between the investigated localities.
16 Small cores on flint nodules are present, excepted at Lademagne where large-sized limestone cores
17 are also found. The cores are never prepared and are always related to the form of the blank. The
18 debitage affects one or two faces (unifacial, bifacial or orthogonal) with unipolar removals. Some
19 round pebbles produced "slices". Cores are never exhausted and the debitage is minimal. The
20 Levallois core technology is absent, except at Lademagne lower and Colle Avarone where one
21 "Levallois-type" flake is evidenced.

22 Heavy-duty tools are the main component of the collected lithic series. Three types are common to
23 the series but in different proportions:

24 (1) Elongated tools, named bifaces, characterized by complete bifacial shaping of the tool volume
25 and thin tips. When the cross-section is symmetrical, the single series of removals is never rectified
26 by final retouch. The edges of these tools are sinuous. When the cross-section is asymmetrical the
27 removals are numerous and a final retouch sometimes rectifies the edges. Series differ between the
28 localities by the presence/absence of final retouch.

29 (2) Partial tools, named bifacial tools. They are made by successive removals on the upper part of the
30 tool, of the cutting edges and of the tip. Cross-sections are mainly asymmetric with a preserved back.

31 (3) Heavy "scrapers", "rabots" on pebbles and pebble-tools.

32 The shaping mode is not linked to the type of raw material. The bifaces on flint share common
33 technological features with the tools on other kinds of stones. However, the mode and intensity of
34 shaping are consistent with the raw material and the shape of the original pebble. The partial bifacial
35 tools are mainly made on (easily identifiable) flakes, considered as pre-forms, which do not require

1 extensive management. These categories are not stages in the shaping process but represent clear
2 distinct tool types with specific purposes (whole volume management, manufacture of the cutting
3 edges and the tip) and sizes, even if all the tools show an asymmetric cross-section. Some partial
4 bifacial tools indicate that resharpening would have been possible since a limited area of the upper
5 part of the tool is shaped by additional series, which can be reworked independently. Conversely,
6 most of the bifaces with total volume manufacture cannot be resharpened without modifying the
7 whole volume and reducing the size of the tool. The whole bifaces are symmetrical in form and
8 elongated while the partial bifacial tools are shorter, not generally elongated and frequently
9 asymmetrical in form and cross-section. Bifaces are the longest pieces (around 200 mm) (**Fig. 19**).
10 Although asymmetrical tools with retouch on one side evoke bifacial scrapers rather than bifaces, the
11 technological study does not indicate strong overall tool reduction. It is important to mention that
12 each tool category frequently displays the same size and size variation for each locality. This suggests
13 a mental model or template for making these tools and the flexibility to adapt the model to the blank
14 (pebble or flake).

15
16 **Table 2** demonstrates that some technological features of the LCTs may partially distinguish the
17 series. Internal variability affects above all the shaping mode of the tip and retouch on the edges, and
18 episodically the presence of a “coup de tranchet”, twisted bifaces and backed tools. Elongated tools
19 are the most common type of tool, as opposed to the short tools and large flakes that are often used
20 for shaping. However, the number of tools differs between localities and the quantity can affect the
21 technological record and picture. We performed a Multiple Correspondence Analysis with a set of
22 variables characterizing technological features (extension removals along faces A and B, series of
23 removals, final retouches, tip shaping, presence or absence of cross-sectional and/or lateral
24 symmetry). Our analysis shows two groups of sites. One group forms a major cluster and one group
25 isolates the sites CG9 upper, Isoletta 4 and Masseria Castellone, which are visible in **table 2 and fig.**
26 **20**. These three original series differ by less careful shaping of the tip and edges and more
27 asymmetrical tools. Number and extension of removals on both faces distinguishes the two groups
28 along dimension 2 (**Fig. 20**), which calls into question differences in reduction sequence phases by
29 locality (mobility of tools between sites or longer reduction sequences in some sites as a result of
30 longer occupations or specific activities). The size of the gravel pebbles is almost always less than the
31 biface size, suggesting that the bifaces were partially made in other areas than those frequented.

32
33

34 ***The Ceprano basin: an original case in Italy?***

1 On a broader scale, it is essential to compare the networks of sites studied here with other Italian
2 Acheulean sites from the same period. Hereafter, the term “Acheulean groups” corresponds to
3 hominin groups able to perform biface production. However, the fact that some localities do not
4 yield bifaces and/or LCTs does not mean that hominins occupying these sites were not capable to
5 produce these kinds of tools. It may mainly depend on the “function” of the site. The quantity of
6 artefacts collected at Campogrande, Isoletta and Lademagne is higher than in the other studied
7 localities such as Colle Avarone or Selvotta (~ 600 m from Ceprano) (Biddittu, 1974). Despite the
8 variable number of artefacts found in these sites, we demonstrate that the technological and
9 morphological features highlighted share common strategies and specific features that may suggest
10 regional trends of traditions persisting for about 50 ka (from MIS 11 interglacial toward the MIS 10
11 glacial inception). For instance, the bifacial tools and bifaces discovered in the Ceprano basin are
12 generally longer than in other Italian sites studied so far. It could be explained by the dimension and
13 quality of the available raw materials but also by the types of activities or regional routines of the
14 knappers of the area.

15 If we integrate our records at a broader scale, the Acheulean layer “m” of the Torre in Pietra (MIS 10)
16 archaeological site, located 26 km northwest of Rome, is characterized by bifaces similar to the
17 Ceprano basin ones, characterized with a high proportion of asymmetric tools (Villa et al., 2016). At
18 Guado San Nicola, located in Molise (Peretto et al., 2016) and also dated to the MIS 10 glacial onset
19 (Pereira et al., 2016), bifaces represent a total of 3-4% of the series and are mainly made in flint and
20 rarely in silicified limestone, on slabs and some flakes. The raw material does not seem to influence
21 the shaping mode and produces asymmetric pieces. We furthermore observe careful shaping of the
22 tips, including final retouch and limited technical investment. Finally, at Castel di Guido (Boschian and
23 Saccà, 2015), close to Torre in Pietra, bifacial tools on elephant bones and on pebbles (various
24 stones) indicate common shaping strategies at this site, regardless of raw materials. Moreover, most
25 of the LCTs are partial bifacial tools characterized mainly by management of the bifacial tip and a
26 lateral bifacial scraper.

27 As mentioned above, the core technology of our series is similar to what is observed at Torre in
28 Pietra (layer m), dated to the MIS 10/MIS 9 transition (Villa et al., 2016; Soriano and Villa, 2017), but
29 also to the younger butchery site of La Polledrara di Cecanibbio (near Rome) (MIS 9, 325 ± 6 ka,
30 Pereira et al., 2017). These assemblages without bifaces and on small flint nodules do not show clear
31 technological changes for more than 100 ka in Latium (i.e. between MIS 11 and MIS 9). The Levallois-
32 like flake found in the lower archaeological of Lademagne (older than 405 ka, Pereira et al., 2018)
33 and Colle Avarone could indicate the occasional use of this technology, as also highlighted at the site
34 of Guado San Nicola (Molise) during the MIS 11/10 transition (Peretto et al., 2016; Pereira et al.,
35 2016; Moncel et al., 2016, 2020), and in the Acheulean site of Rosaneto in Calabria (Biddittu et al.,

1 1984). The presence of this kind of technology supports the hypothesis of a long transition phase
2 between the Lower and Middle Palaeolithic and the coexistence of these two entities in Europe, but
3 also in Asia and Africa during this period for long period (Akhilesh et al., 2018; Deino et al., 2018).

4 5 ***Evidence of regional networks of sites from MIS 11 onwards in Europe?***

6 The small number of known sites in Western Europe anterior to the MIS 12 glacial period suggests
7 either the bad preservation of the sites or a low population density perhaps reflecting the isolation of
8 small and mobile hominin groups before 450 ka. The first hypothesis raises the question of the good
9 preservation of the sites after this arid glacial stage in Central Italy and more generally in Western
10 Europe. Some studies suggest a good preservation of the sites as well as an increase of the
11 population densities in Mediterranean due to the suitable climatic conditions of the MIS 11
12 interglacial, that is one of the warmest and longest (20-30 kyrs) Quaternary interglacials (Karner and
13 Marra, 2003, Moncel et al., 2018a).

14 We here propose to explore some hypotheses on social behaviours. The hominins groups probably
15 adapted their behaviours to local constraints, which may explain both the diversity and common
16 features of toolkit morphology and modes of shaping for this period. Consequently, they were able
17 to maintain social networks and to permanently occupy some areas. This hypothesis might also
18 explain the lack of clear regional traditions, either among a metapopulation or disconnected groups
19 (White and Ashton, 2003; Premo and Hublin, 2009; Premo and Kuhn, 2010; White et al., 2011;
20 Schreve et al., 2015; Moncel et al., 2015; Moncel and Schreve, 2016; Moncel and Ashton, 2018).
21 Hominins only occupied marginal areas under favourable environments, especially in the northwest
22 of the continent. Latitudinal climatic gradients are consistent with the persistence of occupations in
23 the south during periods of climatic deterioration (Moncel et al., 2016). These climatic oscillations
24 regularly compelled hominins to retreat and expand or led to their disappearance despite evidence
25 of occupations at the MIS 16 in the North-West of Europe under cold conditions or during a cooler
26 period into the glacial event (see *e.g.* Antoine et al., 2019). Even in the Italian peninsula, periodic
27 contractions and expansions of hominin groups during rapid climatic variations were clearly shown
28 during MIS 13 and MIS 12 at Valle Giumentina in Abruzzo, for example (Degeai et al., 2017).

29 A large-scale expansion of Acheulean assemblages over Western Europe is discernible after the
30 glacial period known as MIS 12, and various regional traditions appeared and evolved in both the
31 north and south of Europe (Ashton et al., 2008; Moncel et al., 2015; Ashton and Scott, 2016). The
32 long interglacial MIS 11 followed by a short and relatively mild glacial period MIS 10, especially
33 weakly marked in Southern Europe, seems to have been a crucial period. This period already records
34 some evidence of behavioural changes towards the early Middle Palaeolithic, such as for instance,
35 indications of sporadic, more complex and standardized core technologies, organized hunting or

1 changes in land use patterns (Conard et al., 2015; Moigne et al., 2016; Moncel et al., 2016). Early fire
2 control (Roebroeks and Villa, 2011) could have contributed to new hominin abilities to expand their
3 territories and modify their behaviour accordingly. After 500 ka, lithic assemblages are characterized
4 once again by diverse technological behaviours, with or without bifacial tools, but it is possible to
5 identify regional trends due to the higher quantity of lithic assemblages. We observe the
6 standardization of technological processes in biface production, despite the persistence of the
7 diversity of large toolkits. The diversity of biface thickness and shaping intensity are either due to raw
8 material constraints, site location or traditions. The diversity in the cross-sections of plano-convex or
9 symmetrical bifaces depends on the shaping mode and the intensity and location of retouch
10 (unifacial retouch for a scraper on one edge), and also points to varied toolkit purposes and
11 functions, in the same way as for the early Acheulean (Iovita et al., 2017).

12 The morphological results for LCTs differ between geographical areas suggesting regional trends in or
13 between regions. For instance, in the UK, ovate bifaces dominate in some sites (Hoxne), while
14 pointed bifaces are more numerous in other sites (Swanscombe) (Ashton and Lewis, 2012; Moncel et
15 al., 2015). This suggests the arrival of new population(s) or the diffusion of new traditions, during
16 periods of low sea levels and/or local evolutions. Pointed bifaces also exist in France at the same
17 time (*i.e.* Saint-Pierre-les-Elbeuf) (Cliquet et al., 2009; Lamotte and Tuffreau, 1997; Limondin-Lozouet
18 et al., 2010, 2015).

19 After the MIS 12, core technologies become more complex in Western Europe, with intense
20 development of debitage and production control, including the first traces of Levallois technology in
21 Italy, for example (Peretto et al., 2016). Land use management indicates the fragmentation of
22 reduction processes, an increase in the circulation of long-distance lithic raw materials, despite the
23 persistent use of predominantly local raw materials (Moncel et al., 2012; Herisson et al., 2016;
24 Ravon, 2017).

25 Some sites in the Mediterranean region from this time period (*i.e.* MIS 11-10) yielded assemblages
26 suggesting the widespread diffusion of behaviours over a short period of time beginning during MIS
27 11, and above all, specificities in each area related to raw materials (Gran Dolina TD 10, Spain
28 (Atapuerca, García-Medrano et al., 2015). For example, the sites of Terra Amata (South of France, de
29 Lumley et al., 2008) and Menez Dregan (West of France, Ravon, 2017) demonstrate a high quantity of
30 pebbles on beach sites, with micro-regional traditions strongly impacted by the local environment
31 (marine formations) (Ravon, 2017). Equally, large flakes are used more in Southern Europe than in
32 the northwest, possibly due to an adaptation to raw material availability (flint nodules in the north *v.*
33 coarse-grained pebbles and blocks in the south).

34 Of course, changes in climate cycles (after MIS 12 and during the warm and long MIS 11 interglacial),
35 in demography, and in the density of occupations could have influenced these behavioural changes.

1 They could have encouraged the emergence of regional traditions, a wider system of mobility in the
2 landscape, demographic expansion and genetic exchanges (growth and regular adaptation without
3 demographic discontinuities), as well as new skills amplified by more or less rapid processes of
4 population dynamics and cultural change (Guilfoyle, 2005; Winton, 2005; Bocquet-Appel and
5 Tuffreau, 2009; Chazan, 2009; Dennell et al., 2011; Jagher, 2011; Richter, 2011; Manzi et al., 2011;
6 Kleinen et al., 2014; Grove, 2016; Moncel and Schreve, 2016; Moncel et al., 2016; Henry et al., 2017;
7 Vaesen and Houkes, 2017; Carrión and Walker, 2019). The extension of reliable areas for hominin
8 occupations and better adaptation to diverse environments could explain a phenomenon of “product
9 copying” rather than “process copying”, as described by d’Errico et al. (2017) for the Middle Stone
10 Age in South Africa. This would explain the origin of distinct regional traditions due to the flexible
11 behaviour of hominin groups with common toolkits but varied manufacturing processes depending
12 on raw materials.

13

14 In this general frame, the Frosinone-Ceprano basin localities offer the opportunity to observe
15 technological trends and variability from a regional perspective and to describe lithic series within a
16 narrow time frame over a small territory. Of course, we cannot certify that the occupations were
17 contemporaneous and took place at the same time. However, we are working on the “long” time and
18 the time windows of the ages of the sites is relatively short enough to question and compare the
19 occupations in a delimited territory. Despite limited variations, the common behaviours identified in
20 biface production processes and in the morphological results between the localities raise a new
21 question. How can we explain the high morphological homogeneity (with some types of frequent
22 homogeneous sizes) of assemblages from this basin and their low technological variability? The
23 expertise used for making the LCTs suggests the existence of a common mental template and the
24 flexibility of expert knappers, regardless of the number of pieces and the raw materials (some flint is
25 shaped in the same way as limestone). This could indicate a sort of “network” of sites with strong
26 relationships over time between mobile groups occupying a small region in a short window of time,
27 sharing a micro-regional tradition, possibly with sub-divisions of a metapopulation covering large
28 areas. Studies of late Neanderthal populations suggest low effective population size and
29 demographic isolation and some evidence tends to indicate a similar structure for the earlier period
30 (Ríos et al., 2019). Socio-demographic processes could also affect cognition by the diffusion of new
31 behaviours in a limited area which could maintain stable behaviours, where efficient, leading to the
32 isolation of regional populations (Hamilton et al., 2007; Shipton et al., 2009; Hill et al., 2011;
33 Malinsky-Buller and Hovers, 2019). Slight differences in technology and morphology could be due to
34 variable parameters such as activities, the duration of occupation or site location, but also to
35 individual preferences or the adaptation of expert knappers to the available blanks. We can mention

1 for instance some relevant features of bifaces that, unlike core technologies, distinguish localities:
2 occasional presence of tools with a transverse edge (cleaver-like), intensity of tip-shaping on bifaces,
3 presence of final retouch on bifaces, ratio of short LCTs that are not due to reduction during the
4 shaping process, use of a “coup de tranchet”, presence of twisted bifaces and backed bifaces.
5 LCTs in the Ceprano basin localities are often made on poorly preserved local limestone and are
6 often covered by patches of breccia, rendering the identification of detailed land-use patterns over
7 territories difficult (residential camps or occasional habitats). Moreover, most of the limestone and
8 flint pebbles can be found *in situ*. The topography of the basin could provide an idea of a potential
9 territory with the efficient management of space in an area rich in available blocks, pebbles, flint
10 nodules and large game. Moreover, in some localities (Isoletta, Campogrande), retouched fragments
11 of large mammal bones show the exploitation of elephants and other large herbivore carcasses and
12 bone fragments and this trend is not an isolated case in Latium (Fontana Ranuccio, Torre in Pietra, La
13 Polledrara di Cecanibbio and Castel di Guido) (Piperno and Biddutti, 1978; Biddittu et al., 1979;
14 Ascenzi and Segre, 1997; Radmilli and Boschain, 1996; Segre et al., 2009; Muttoni et al., 2009;
15 Anzidei et al., 2012; Saccà, 2012; Boschian and Saccà, 2015; Santucci et al., 2016; Villa et al., 2016;
16 Pereira et al., 2017). Habitat quality and resources (availability of large fragments of bones) could
17 explain the density of occupations and networks of sites with similar activities, with hominins
18 circulating between localities depending on seasonal resources.
19 The type of mobility and occupations based on current models of mobility (foraging radius and
20 logistical radius, hierarchical sites) cannot be confidently identified in our corpus (*e.g.*, Binford, 1980,
21 1982; Kühn, 1992). When data are available, most of the sites of this period are butchery sites and at
22 the same time record domestic activities, along rivers, lakeshores, and stone outcrops, with or
23 without a hierarchical structure of occupations (aggregation-dispersal mechanism) and are
24 dependent on local ecological conditions and seasonal variations (as demonstrated at the Acheulean
25 site GBY in Israel, by Herlinger and Goren-Inbar, 2019). Each series of our corpus with LCTs could
26 portray several occupations at favourable spots or one main occupation, and the hypothesis of
27 catchment in some cases must be considered for the smallest series. Our series contain few flakes
28 issued from shaping processes and no evidence of *in situ* biface production. However, that does not
29 mean that bifaces were only mobile tools transported from site to site and introduced to the sites in
30 a finished state. Due to the lack of extensive excavations at our localities, it is not possible to observe
31 activity areas. However, it is important to mention that some LCTs occasionally show crushing marks
32 on their edges, indicating use, and a more or less large quantity of these tools was abandoned.
33 Perhaps these tools were easily reproducible owing to the widespread availability of large formations
34 of pebbles. Binford (1980) suggested that environments offering homogeneously distributed resources
35 would have been exploited by frequent residential mobility, while those supplying heterogeneous

1 resources would be best exploited by low residential mobility and by logistic displacement for
2 specific resources. Mithen (1994) and Bocquet-Appel and Tuffreau (2009) suggested a possible
3 higher mobility of small groups during interglacial periods. Grove (2009) demonstrates that habitat
4 quality is the best predictor of the distance travelled by hunter-gatherers between camps. Likewise,
5 large groups move often, rather than far (idea of costly strategy and high-mobility when necessary).
6 We are not capable of identifying group size or social units, or the duration of occupations on the
7 basis of the size of our lithic series, which may possibly result from recurrent passages and
8 palimpsests (Powell et al., 2009; Bocquet-Appel and Degioanni, 2013; Collard et al., 2013; Derex et
9 al., 2013; Vaesen et al., 2016; Dominguez-Rodrigo et al., 2019). Nor do we have any idea of the
10 original surface area of the occupations. If we admit that the smallest series, if relevant, evidenced
11 short-term camps or ephemeral passages, while the largest ones indicate large camps or palimpsests
12 of regular short-term occupations, the technology used everywhere indicates expert knappers and
13 the smaller lithic series are neither more homogeneous nor heterogeneous than the larger ones. The
14 quantity of bifaces tends to demonstrate how hominins managed their toolkits in the landscape and
15 between occupations. Our series attest that LCTs were regularly abandoned while still usable.

16

17 **Conclusion**

18 The different localities of the Frosinone-Ceprano basin with lithic artefacts are located in a small-
19 sized area. Radio-isotopic constraints as well as geological and archaeological information indicate
20 the contemporaneity or penecontemporaneity of these localities and related series allow for
21 comparisons of the technological behaviours of hominins during the MIS 11-10 time period in this
22 region. Core technologies were simple and similar for all localities. Bifaces and bifacial tools were
23 managed in the same way with minor differences between localities. Tools are generally large in size
24 (ranging in length from 50 to 260 mm). Common trends in toolkits could indicate networks of
25 connected sites along fluvial systems in the basin and record the use of locally available raw
26 materials.

27 Future analyses of bone assemblages will help to understand which types of sites existed and which
28 subsistence activities were performed and how this network worked in relation to the exploitation of
29 carcasses of elephants and other large herbivores. This could explain why core technologies show
30 minimal flaking aimed at rapidly producing flakes in sporadic occupations. A heavy-duty component
31 is present at some sites, and this cannot be explained by stone availability. The quantity of flakes
32 indicating an *in situ* shaping process is often small and it is consequently difficult to investigate the
33 status of these large tools. They could be mobile pieces transported between sites or expedient tools
34 prepared for the activities performed on the site.

1 Current genetic data and anthropological analyses show that Neanderthal features emerged in the
2 European population between 600 and 400 ka, and various hominins may have co-existed at that
3 time (Hublin, 2009; Stringer, 2012). Our analysis supports the hypothesis of early evidence of
4 regional traditions at the end of this transitional morphological period with an increase in the
5 number of sites, which cannot solely be attributed to better preservation as a result of tectonic and
6 climatic factors. This regionalization is not only discernible for the predecessors of Neanderthal
7 populations, but also for the archaic hominin of Ceprano. This process of regionalization is not
8 restricted to the Italian Peninsula but is a broader process that is also apparent in the Northwest and
9 South of Europe (Ashton et al., 2008; Rightmire, 2008; Scott and Ashton, 2011; Ashton and Lewis,
10 2012; Moncel et al., 2011, 2012, 2018; Fontana et al., 2013; Blain et al., 2014, 2015; Brenet et al.,
11 2014; Wisniewski, 2014; Garcia-Medrano et al., 2015; Ashton and Scott, 2016).

12

13 **Acknowledgements**

14 The analyses of the lithic series were financially supported by funding from the National Museum of
15 Natural History (Paris, France) (Action Transversale du Museum, "Acheulean in volcanic areas in
16 Italy"). Surveys were performed by I. Biddittu et al. The excavations and paleoenvironmental analyses
17 at Campogrande were performed within the framework of the project "The Ceprano calvarium and
18 its environment", with financial backing from various grants attributed to one of us (G.M.), by the
19 Sapienza University of Rome. This work has been presented at the USIPP congress 2018 Paris in the
20 session "MIS 13-11: a major transformation in the European Lower Palaeolithic" organized by James
21 Cole and Rob Hosfield. The text has been edited by Louise Byrne, English native speaker and official
22 translator. We would like to thank the two reviewers and the editor for their comments, which
23 helped us to deeply enhance the manuscript.

24

25 **References**

- 26 Antoine P, Moncel M-H, Locht J-L Bahain J-J., Voinchet P, Herisson D, Hurel A (2019) The earliest
27 record of Acheulean human occupation in North-West Europe. *Nature Scientific Reports* 9:13091
- 28 Anzidei AP, Bulgarelli GM, Catalano P, Cerilli E, Gallotti R, Lemorini C, Milli S, Palombo MR, Pantano
29 W, Santucci E (2012) Ongoing research at the late Middle Pleistocene site of La Polledrara di
30 Cecanibbio (central Italy), with emphasis on human-elephant relationships. *Quaternary International*
31 255: 171–187
- 32 Akhilesh K, Pappu S, Rajapara HM, Gunnell Y, Shukla AD, Singhvi AK (2018) Early Middle Palaeolithic
33 culture in India around 385-172 ka reframes out of Africa models. *Nature* 554: 97–101
- 34 Ascenzi A, Biddittu I, Cassoli PF, Segre AG, Segre Naldini E (1996) A calvarium of late *Homo erectus*
35 from Ceprano, Italy. *Journal of Human Evolution* 31: 409-423

- 1 Ascenzi A, Segre AG (1997) Discovery of a Homo erectus calvarium at Ceprano, Central Italy.
2 Anthropologie XXXV: 241–246
- 3 Ascenzi A, Mallegni F, Manzi G, Segre AG, Segre Naldini E (2000) A re-appraisal of Ceprano calvaria
4 affinities with Homo erectus, after the new reconstruction. Journal of Human Evolution 39: 443-450
- 5 Ashton N, Lewis SG, Parfitt SA, Penkman KEH, Russell Coope G (2008) New evidence for complex
6 climate change in MIS 11 from Hoxne, Suffolk, UK. Quaternary science reviews 27: 652–668
- 7 Ashton N, Lewis S G (2012) The environmental contexts of early human occupation of northwest
8 Europe: The British Lower Palaeolithic record. Quaternary International 271: 50-64
- 9 Ashton N, Scott B (2016) The British Middle Palaeolithic. Quaternary International 411: 62-76
- 10 Basalone P, Civetta L (1975) Datazione K-Ar dell'attività vulcanica dei Mt Ernici (Latina). Società
11 Italiana Mineralogia e Petrologia 31(1): 175-179
- 12 Biddittu I (1974) Giacimento pleistocenico ad amigdale acheuleane nel territorio di Ceprano,
13 Frosinone. Memoria dell'Istituto Italiano di Paleontologia Umana 2: 61–67
- 14 Biddittu I, Cassoli PF, Di Brozolo R, Segre AG, Segre NE, Villa I (1979) Anagni, a K-Ar dated Lower and
15 Middle Pleistocene site, Central Italy : preliminary report. Quaternaria 21: 53–71
- 16 Biddittu I, Segre AG (1984) Industria su scheggia e bifacciali: nuovi reperti del paleolitico inferiore ad
17 Anagni-Fontana Ranuccio, Frosinone. Atti della XXIV riunione scientifica dell'Istituto Italiano di
18 Preistoria e Protostoria 24: 105–112
- 19 Biddittu I, Piperno M, Segre AG (1984) Rosaneto, Calabria. I primi abitanti d'Europa, mostra Museo
20 Nazionale Luigi Pigorini, marzo-luglio 1984, De Luca Ed, pp 151-154
- 21 Biddittu I (2004) Guida del Museo Preistorico di Pofi I. Quad di ARGIL 1, 157: 49-58 and 114-119
- 22 Biddittu I, Canetri E, Comerci V, Germani M, Picchi G (2012) Nuove ricerche nel giacimento del
23 Paleolitico inferiore di Lademagne, S Giovanni Incarico (Frosinone) In: (Ghini, G, Mari, Z, eds) Lazio e
24 Sabina Edizioni Quasar 9, pp 437-443
- 25 Biddittu I, Moncel M-H, Milli S, Bellucci L, Ruffo M, Saracino B, Manzi G (2019) Stratigraphy,
26 sedimentology, and archaeology of Middle Pleistocene localities near Ceprano, Campogrande area,
27 Italy. Quaternary Research 93(1): 155-171.
- 28 Binford LR (1980) Willow smoke and dogs' tails – hunter–gatherer settlement systems and
29 archaeological site formation. American Antiquity 45: 4–20
- 30 Binford LR (1982) The archaeology of place. Journal of Anthropological Archaeology 1: 5–31
- 31 Blain H-A, Santonja M, Pérez-González A, Panera J, Rubio-Jara S (2014) Climate and environments
32 during Marine Isotope Stage 11 in the central Iberian Peninsula: the herpetofaunal assemblage from
33 the Acheulean site of Áridos-1, Madrid. Quaternary Science Reviews 94: 7–21

- 1 Blain H-A, Lozano-Fernández I, Ollé A, Rodríguez J, Santonja M, Pérez-González A (2015) The
2 continental record of Marine Isotope Stage 11 (Middle Pleistocene) on the Iberian Peninsula
3 characterized by herpetofaunal assemblages. *Journal of Quaternary Science* 30: 667–678
- 4 Boari E, Tommasini S, Laurenzi MA, Conticelli S (2009) Transition from Ultrapotassic Kamafugitic to
5 Sub-alkaline Magmas: Sr, Nd, and Pb Isotope, Trace Element and ^{40}Ar - ^{39}Ar Age Data from the
6 Middle Latin Valley Volcanic Field, Roman Magmatic Province, Central Italy. *Journal of Petrology*
7 50(7): 1327-1357
- 8 Bocquet-Appel J P, Tuffreau A (2009) Technological responses of Neanderthals to macroclimatic
9 variations (240,000–40,000 BP). *Human Biology* 81(3): 287-307
- 10 Bocquet-Appel J-P, Degioanni A (2013) Neanderthal demographic estimates. *Current Anthropology*
11 54 : S202-S213
- 12 Boëda E, Geneste J-M, Meignen L (1990) Identification de chaînes opératoires lithiques du
13 Paléolithique ancien et moyen. *Paleo* 2: 43-80
- 14 Boschian G, Saccà D (2015) In the elephant, everything is good: Carcass use and re-use at Castel di
15 Guido (Italy). *Quaternary International* 361: 288-296
- 16 Brenet M, Bourguignon L, Colonges D, Folgado M, Jarry M, Lelouvier L-A, Mourre V, Turq A (2014) Les
17 techno-complexes au début du Paléolithique moyen en Aquitaine septentrionale. In: *Transitions,*
18 *Ruptures et Continuité En Préhistoire*, pp 81–101
- 19 Burke A, Ebert D, Cardille J, Dauth D (2008) Paleoethology as a tool for the development of
20 archaeological models of land-use: the Crimean Middle Palaeolithic. *Journal of Archaeological*
21 *Science* 35(4): 894-904
- 22 Carrara C, Frezzotti M, Giraudi C (1995) Stratigrafia plio-quadernaria In: Carrara, C (Ed), *Lazio*
23 *Meridionale, Sintesi delle ricerche geologiche multidisciplinari ENEA*, pp 62–85
- 24 Carrión J S, Walker M J (2019) Background to Neanderthal presence in Western Mediterranean
25 Europe. *Quaternary Science Reviews* 217: 7-44
- 26 Conard N J, Serangeli J, Böhner U, Starkovich B M, Miller C E, Urban B, Van Kolfschoten T (2015)
27 Excavations at Schöningen and paradigm shifts in human evolution. *Journal of Human Evolution* 89:
28 1-17
- 29 Centamore E, Di Manna P, Rossi D (2007) Kinematic evolution of the Volsci Range: a new overview.
30 *Bollettino della Società Geologica Italiana* 126 (2): 159-172
- 31 Centamore E, Dramis F (2010) Note illustrative della Carta geologica d'Italia alla scala 1:50000, Foglio
32 402 Ceccano Regione Lazio, Roma
- 33 Chazan M (2009) Assessing the Lower to Middle Paleolithic Transition In: Camps M, Chauhan P (eds),
34 *Sourcebook of Paleolithic Transitions* Springer, pp 237–243

- 1 Civetta L, Innocenti F, Lirer L, Manetti P, Munno R, Peccirillo A, Poli G, Serri G (1979) Serie potassica
2 ed alta in potassio dei Monti Ernici (Lazio Meridionale): considerazioni petrologiche e geochemiche.
3 Rend Soc Ital Mineral Petrol 35: 227-249
- 4 Cliquet D, Lautridou J P, Antoine P, Lamothe M, Leroyer M, Limondin-Lozouet N, Mercier N (2009) La
5 séquence loessique de Saint-Pierre-lès-Elbeuf (Normandie, France): nouvelles données
6 archéologiques, géochronologiques et paléontologiques. Quaternaire 20(3): 321-343
- 7 Collard M, Buchanan B, O'Brien M J (2013) Population size as an explanation for patterns in the
8 Paleolithic archaeological record: more caution is needed. Current Anthropology 54(S8): S388-S396
- 9 Comerci V, Biddittu I, Di Manna P, Germani M, Piccardi L, Ventura G, Vittori E (2015) Tectonic
10 evidence in the Palaeolithic site of Lademagne (San Giovanni Incarico – FR, Southern Latium, Italy)
11 6th International INQUA Meeting on Paleoseismology, Active Tectonics and Archaeoseismology 19-
12 24 April 2015, Pescina, Fucino Basin, Italy INQUA Focus Group on Paleoseismology and Active
13 Tectonics
- 14 Daujeard C, Moncel M-H (2010) On Neanderthal subsistence strategies and land-use: a regional focus
15 on the Rhône Valley area in southeastern France. Journal of Anthropological Archaeology 29: 368-
16 391
- 17 Deino A L, Behrensmeyer A K, Brooks A S, Yellen J E, Sharp W D, Potts R (2018) Chronology of the
18 Acheulean to Middle Stone Age transition in eastern Africa. Science 360(6384): 95-98
- 19 Degeai JP Villa V, Chaussée C, Pereira A, Nomade S, Aureli D, Pagli M, Nicoud E (2017), Chemical
20 weathering of paleosols associated to Lower Palaeolithic settlements at Valle Giumentina (Abruzzes,
21 Italy). Quaternary Science Reviews 183: 88-109
- 22 Dennell R W, Martínón-Torres M, de Castro J M B (2011) Hominin variability, climatic instability and
23 population demography in Middle Pleistocene Europe. Quaternary Science Reviews 30(11-12): 1511-
24 1524
- 25 Derex M, Beugin M P, Godelle B, Raymond M (2013) Experimental evidence for the influence of
26 group size on cultural complexity. Nature 503(7476): 389
- 27 Devoto G (1965) Lacustrine Pleistocene in the lower Liri Valley (southern Latium). Geologica Romana
28 4: 291–368
- 29 D'Errico F, Banks W, et al (2017) Identifying early modern human ecological niche expansions and
30 associated cultural dynamics in the South African Middle Stone Age. Proceedings of the National
31 Academy of Sciences 114(30): 7869-7876
- 32 Di Vincenzo F, Profico A, Bernardini F, Cerroni V, Dreossi D, Schlager S, Zaio P, Benazzi S, Biddittu I,
33 Rubini M, Tuniz C, Manzi G (2017) Digital reconstruction of the Ceprano calvarium (Italy) and
34 implications for its interpretation. Scientific Reports 7 (13974): 1-11
- 35 Domínguez-Rodrigo M, Cobo-Sánchez L, Aramendi J, Gidna A (2019) The meta-group social network
36 of early humans: A temporal–spatial assessment of group size at FLK Zinj (Olduvai Gorge, Tanzania).
37 Journal of human evolution 127: 54-66.

- 1 Faivre J P, Gravina B, Bourguignon L, Discamps E, Turq A (2017) Late Middle Palaeolithic lithic
2 technocomplexes (MIS 5–3) in the northeastern Aquitaine Basin: Advances and challenges.
3 Quaternary International 433: 116-131
- 4 Fontana F, Moncel M-H, Nenzioni G, Onorevoli G, Peretto C, Combler J (2013) Widespread diffusion
5 of technical innovations around 300,000 years ago in Europe as a reflection of anthropological and
6 social transformations? New comparative data from the western Mediterranean sites of Orgnac
7 (France) and Cave dall’Olio (Italy). *Journal of Anthropological Archaeology* 32: 478–498
- 8 García-Medrano P, Ollé A, Mosquera M, Cáceres I, Carbonell E (2015) The nature of technological
9 changes: The Middle Pleistocene stone tool assemblages from Galería and Gran Dolina-subunit
10 TD101 (Atapuerca, Spain). *Quaternary International* 368: 92–111
- 11 Geneste JM (1991) Systèmes techniques de production lithique : variations technoéconomiques dans
12 les processus de réalisation des outillages paléolithiques. *Technology and Culture* 17 (18): 1–35
- 13 Giannetti B (2001) Origin of the calderas and evolution of the Roccamonfina volcano (Roman region,
14 Italy). *Journal of Volcanology and Geothermal Research* 106: 301-319
- 15 Girone A, Maiorano P, Marino M, Kucera M (2013) Calcareous plankton response to orbital and
16 millennial scale climate changes across the Middle Pleistocene in the western Mediterranean.
17 *Palaeogeogr. Palaeoclimatol. Palaeoecol* 392: 105-116. doi:10.1016/j.palaeo.2013.09.005
- 18 Goren-Inbar N, Sharon G (2006) *Axe Age: Acheulian Tool-making from Quarry to Discard*, Equinox
19 Publishing Ltd, London
- 20 Grove M (2009) Hunter–gatherer movement patterns: causes and constraints. *Journal of*
21 *Anthropological Archaeology* 28(2): 222-233
- 22 Grove M (2016) Population density, mobility, and cultural transmission. *Journal of Archaeological*
23 *Science* 74: 75-84
- 24 Guilfoyle D R (2005) Socializing stone artifact assemblages: Regionalization and raw material
25 availability in northern Queensland Australian. *Archaeology* 60(1): 34-40
- 26 Hamilton MJ, Milne BT, Walker RS, Burger O, Brown JH (2007) The complex structure of hunter–
27 gatherer social networks. *Proceedings of the Royal Society B-Biological Sciences* 274: 2195–2202
- 28 Henry D O, Belmaker M, Bergin S M (2017) The effect of terrain on Neanderthal ecology in the
29 Levant. *Quaternary International* 435: 94-105
- 30 Hérisson D, Airvaux J, Lenoble A, Richter D, Claud E, Primaud, J (2016) Between the northern and
31 southern regions of western Europe: the acheulean site of La Grande Vallée (Colombiers, Vienne,
32 France). *Quaternary International* 411: 108-131
- 33 Herzlinger, G, Goren-Inbar, N (2019) Do a few tools necessarily mean a few people? A techno-
34 morphological approach to the question of group size at Gesher Benot Ya'aqov, Israel *Journal of*
35 *Human Evolution* 128: 45-58

- 1 Hill K R, Walker R S, Božičević M, Eder J, Headland T, Hewlett B, Wood B (2011) Co-residence patterns
2 in hunter-gatherer societies show unique human social structure. *Science* 331(6022): 1286-1289
- 3 Hublin J J (2009) The origin of Neandertals. *Proceedings of the National Academy of Sciences* 106(38):
4 16022-16027
- 5 Iovita R, Tuvi-Arad I, Moncel M-H, Despriée J, Voinchet P, Bahain J-J (2017) High handaxe symmetry
6 at the beginning of the European Acheulian: the data from la Noira (France) in context. *Plos One*
7 12(5): e0177063
- 8 Jagher R (2011) Nadaouiyeh Aïn Askar - Acheulian variability in the Central Syrian Desert In: Le
9 Tensorer, J-M, Jagher, R, Otte, M (eds) *The Lower and Middle Palaeolithic in the Middle East and*
10 *Neighbouring ERAUL* Université de Liège, University of Basel, pp 209-225
- 11 Karner DB, Renne PR (1998) ⁴⁰Ar/³⁹Ar geochronology of Roman volcanic province tephra in the
12 Tiber River Valley: age calibration of Middle Pleistocene sea-levels changes. *Geological Society of*
13 *America Bulletin* 110: 140-147
- 14 Karner DB, Marra F (2003) ⁴⁰Ar/³⁹Ar dating of Glacial Termination V and duration of the Stage 11
15 highstand. In: *Earth's Climate and Orbital Eccentricity: The Marine Isotope Stage 11 Question*,
16 American Geophysical Union, Geophysical Monograph, 137, pp. 61-66.
- 17 Key A J, Proffitt T, Stefani E, Lycett S J (2016) Looking at handaxes from another angle: Assessing the
18 ergonomic and functional importance of edge form in Acheulean bifaces. *Journal of Anthropological*
19 *Archaeology* 44: 43-55
- 20 Kleindienst MR (1961) Variability within the late Acheulian assemblage in Eastern Africa. *South*
21 *African Archaeological Bulletin* 16: 35-52
- 22 Kleinen T, Hildebrandt S, Prange M, Rachmayani R, Müller S, Bezrukova E, Brovkin V, Tarasov PE
23 (2014) The climate and vegetation of Marine Isotope Stage 11 – Model results and proxy-based
24 reconstructions at global and regional scale. *Quaternary International* 348: 247–265
- 25 Kuhn S L (1992) On planning and curated technologies in the Middle Paleolithic. *Journal of*
26 *Anthropological Research* 48(3): 185-214
- 27 Leakey MD (1971) *Olduvai Gorge: Excavations in Bed I and Bed II, 1960-1963*, Cambridge University
28 Press
- 29 Limondin-Lozouet N, Nicoud E, Antoine P, Auguste P, Bahain J J, Dabkowski J, Jolly-Saad M C (2010)
30 Oldest evidence of Acheulean occupation in the Upper Seine valley (France) from an MIS 11 tufa at La
31 Celle. *Quaternary international* 223: 299-311
- 32 Limondin-Lozouet N et al (2015) Northwest European MIS 11 malacological successions: a framework
33 for the timing of Acheulean settlements. *Journal of Quaternary Science* 30 (7): 702-712
- 34 De Lumley H (2008) *Terra Amata Nice Alpes-Maritimes, France Tome I*, CNRS Editions

- 1 Lycett S J, Schillinger K, Eren M I, von Cramon-Taubadel N, Mesoudi A (2016) Factors affecting
2 Acheulean handaxe variation: Experimental insights, microevolutionary processes, and
3 macroevolutionary outcomes. *Quaternary International* 411: 386-401
- 4 McNabb J, Cole J, Hoggard C S (2018) From side to side: Symmetry in handaxes in the British Lower
5 and Middle Palaeolithic. *Journal of Archaeological Science: Reports* 17: 293-310
- 6 McPherron S (2006) What typology can tell us about Acheulian handaxe production In: Goren –
7 Inbar, N, Sharon, G (eds), *Axe Age Acheulian Tool-making from Quarry to Discard*, Equinox Publishing
8 Ltd, London, pp 267-287
- 9 Malinsky-Buller, A, Hovers E (2019) One size does not fit all: Group size and the late middle
10 Pleistocene prehistoric archive. *Journal of human evolution* 127: 118-132
- 11 Manzi G (2012) On the trail of the genus *Homo* between archaic and derived morphologies. *Journal*
12 *of Anthropological Sciences* 90: 99-116
- 13 Manzi G (2016) Humans of the Middle Pleistocene: The controversial calvarium from Ceprano (Italy)
14 and its significance for the origin and variability of *Homo heidelbergensis*. *Quaternary International*
15 411: 254-261
- 16 Manzi G, Mallegni F, Ascenzi A (2001) A cranium for the earliest Europeans: Phylogenetic position of
17 the hominid from Ceprano, Italy. *Proceedings of the National Academy of Sciences* 98: 10011-10016
- 18 Manzi G, Magri D, Milli S, Palombo MR, Margari V, Celiberti V, et al (2010) The new chronology of the
19 Ceprano calvarium (Italy). *Journal of Human Evolution* 59: 580-585
- 20 Manzi G, Magri D, Palombo MR (2011) Early-Middle Pleistocene environmental changes and human
21 evolution in the Italian peninsula. *Quaternary Science Reviews* 30: 1420–1438
- 22 Margari V, Roucoux K, Magri D, Manzi G, Tzedakis PC (2018) The MIS 13 interglacial at Ceprano, Italy,
23 in the context of Middle Pleistocene vegetation changes in southern Europe. *Quaternary Science*
24 *Reviews* 199: 144-158
- 25 Marra F, Karner DB, Freda C, Gaeta M, Renne PR (2009) Large mafic eruptions at the Alban Hills
26 Volcanic District (Central Italy): chronostratigraphy, petrography and eruptive behavior. *Journal of*
27 *Volcanology and Geothermal Research* 179: 217–232
- 28 Masson-Delmotte, V., Stenni, B., Pol, K., Braconnot, P., Cattani, O., Falourd, S., Jouzel, J., Landais, A.,
29 Minster, B., Barnola, J.M., Chappellaz, J., Krinner, G., Johnsen, S., Röthlisberger, R., Hansen, J.,
30 Mikolajewicz, U., Otto-Bliesner, B., 2010. EPICA Dome C record of glacial and interglacial intensities.
31 *Quaternary Science Reviews* 29 (1): 113-128.
32
- 33 Milli S, Palombo MR (2005) The high-resolution sequence stratigraphy and the mammal fossil record:
34 A test in the Middle-Upper Pleistocene deposits of the Roman Basin (Latium, Italy). *Quaternary*
35 *International* 126-128: 251-270
- 36 Mithen S (1994) Technology and society during the Middle Pleistocene: hominid group size, social
37 learning and industrial variability. *Cambridge Archaeological Journal* 4(1): 3-32.

- 1 Moigne A-M, Valensi P, Auguste P, García-Solano J, Tuffreau A, Lamotte A, Barroso C, Moncel M-H
2 (2016) Bone retouchers from Lower Palaeolithic sites: Terra Amata, Orgnac 3, Cagny-l'Épinette and
3 Cueva del Angel. *Quaternary International* 409: 195-212
- 4 Moncel M-H, Moigne A-M, Sam Y, Combier J (2011) The Emergence of Neanderthal Technical
5 Behavior: New Evidence from Orgnac 3 (Level 1, MIS 8), Southeastern France. *Current Anthropology*
6 52: 37–75
- 7 Moncel M-H, Moigne A-M, Combier J (2012) Towards the Middle Palaeolithic in Western Europe: the
8 case of Orgnac 3 (southeastern France). *Journal of human evolution* 63: 653–66
- 9 Moncel M-H, Ashton N, Lamotte A, Tuffreau A, Cliquet D, Despriée J (2015) The Early Acheulian in
10 north-western Europe. *Journal of Anthropological Archaeology* 40: 302-331
- 11 Moncel M-H, Arzarello M, Peretto C (2016) Editorial The Holstainian Eldorado. *Quaternary*
12 *International* 409: 1-8
- 13 Moncel M-H, Schreve D (2016) Editorial European Acheuleans The Acheulean in Europe : origins,
14 evolution and dispersal. *Quaternary International* 411 (part B): 1-8
- 15 Moncel M H, Landais A, Lebreton V, Combourieu-Nebout N, Nomade S, Bazin L (2018a) Linking
16 environmental changes with human occupations between 900 and 400 ka in Western Europe.
17 *Quaternary International* 480: 78-94
- 18 Moncel M-H, Arzarello M, Boëda E, Bonilauri S, Chevrier B, Gaillard C, Forestier H, Yinghua Li Sémah,
19 F, Zeitoun V (2018b) The assemblages with bifacial tools in Eurasia (first part) What is going on in the
20 West? Data on Western and Southern Europe and the Levant, *CR Palevol*, special issue Eurasian
21 Pleistocene (Y Coppens and A Vialet eds) 17 (1-2): 45-60
- 22 Moncel M-H, Ashton N (2018c) From 800 to 500 ka in Europe The oldest evidence of Acheuleans in
23 their technological, chronological and geographical framework. In: M Mussi and R Gallotti eds,
24 Chapter 11 Springer edition. *The Emergence of the Acheulean in East Africa*, pp 215-235
- 25 Moncel M-H, Ashton N, Arzarello M, Fontana F, Lamotte A, Scott B, Muttillio B, Berruti B, Nenzioni G,
26 Tuffreau A., Peretto C. (2020a) Early Levallois core technology between MIS 12 and 9 in Western
27 Europe? *Journal of Human Evolution* 139: in press
- 28 Moncel M-H, Despriée J, Courcimaut G, Voinchet P, Bahain J-J (2020b) La Noira site (Centre, France)
29 and the technological behaviours and skills of the earliest Acheulean in Western Europe between 700
30 and 600 kyrs. *Journal of Paleolithic Archeology*: in press
- 31 Muttoni G, Scardia G, Kent D V, Swisher C, Manzi G (2009) Pleistocene Magnetostratigraphy of Early
32 Hominin Sites at Ceprano and Fontana Ranuccio, Italy. *Earth and Planetary Science Letters* 286: 255–268
- 33 Nomade S, Muttoni G, Guillou H, Robin E, Scardia G (2011) First $^{40}\text{Ar}/^{39}\text{Ar}$ age of the Ceprano man
34 (central Italy). *Quaternary Geochronology* 6: 453-457
- 35 Pereira A, Nomade S, Shao Q, Bahain JJ, Arzarello M, Douville E, Falguères C, Frank N, Garcia T,
36 Lembo G, Muttillio B, Peretto C (2016) $^{40}\text{Ar}/^{39}\text{Ar}$ and ESR-U/Th dates for Guado San Nicola, Middle

- 1 Pleistocene key site at the Lower/Middle Palaeolithic transition in Italy. *Quaternary Geochronology*
2 36: 67-75
- 3 Pereira A, Nomade S, Falguères C, Bahain JJ, Tombret O, Garcia T, Voinchet P, Bulgarelli GM, Anzidei
4 AP (2017) New $^{40}\text{Ar}/^{39}\text{Ar}$ and ESR/U-series data for La Polledrara di Cecanibbio archaeological site
5 (Lazio, Italy). *Journal of Archaeological Science: Reports* 15: 20-29
- 6 Pereira A, Nomade S, Moncel M-H, Voinchet P, Bahain J-J, Biddittu I et al (2018) Geochronological
7 evidences of a MIS 11 to MIS 10 age for several pivotal Acheulian sites from the Frosinone province
8 (Latium, Italy): Archaeological implications. *Quaternary Sciences Reviews* 187: 112-129
- 9 Peretto C, Arzarello M, Bahain J-J, Boulbes N, Dolo J-M, Douville E, Falguères C, Frank N, Garcia T,
10 Lembo G, Moigne A-M, Muttillo B, Nomade S, Pereira A, Rufo MA, Sala B, Shao Q, Thun Hohenstein
11 U, Tessari U, Turrini MC, Vaccaro C (2016) The Middle Pleistocene site of Guado San Nicola
12 (Monteroduni, Central Italy) on the Lower/Middle Palaeolithic transition. *Quaternary International*
13 411: 301–315
- 14 Piperno M, Biddittu I (1978) Studio tipologico ed interpretazione dell'industria acheuleana e pre-
15 musteriana dei livelli m e d di Torre in Pietra (Roma) in Torre in Pietra, Roma. *Quaternaria* 20: 441–
16 536
- 17 Powell A, Shennan S, Thomas M G (2009) Late Pleistocene demography and the appearance of
18 modern human behavior. *Science* 324(5932): 1298-1301
- 19 Premo LS, Hublin J-J (2009) Culture, population structure, and low genetic diversity in Pleistocene
20 hominins. *Proceedings of the National Academy of Sciences* 106: 33-37
- 21 Premo L S, Kuhn S L (2010) Modeling effects of local extinctions on culture change and diversity in
22 the Paleolithic. *PLoS One* 5(12): e15582
- 23 Radmilli AM, Boschian G (1996) Gli scavi a Castel di Guido Il più antico giacimento di cacciatori del
24 Paleolitico inferiore nell'Agro Romano Istituto Italiano di Preistoria e Protostoria, Firenze
- 25 Ravon A-L (2017) Originalité et développement du Paléolithique inférieur à l'extrémité occidentale de
26 l'Eurasie : le « Colombanien » de Menez-Dregan (Plouhinec, Finistère, Bretagne) Phd University of
27 Rennes 1
- 28 Rightmire GP (2008) Homo in the Middle Pleistocene: hypodigms, variation, and species recognition.
29 *Evolutionary Anthropology* 17: 8-21
- 30 Richter J (2011) When did the Middle Palaeolithic Begin? In: Conard, NJRJ (Ed), *Neanderthal Lifeways,*
31 *Subsistence and Technology One Hundred Fifty Years of Neanderthal Study*, pp 7–15
- 32 Ríos L, Kivell, T L, Lalueza-Fox, C, Estalrich, A, García-Taberner, A, Huguet, R, Quintino, Y, de la
33 Rasilla M, Rosas A (2019) skeletal Anomalies in the Neandertal Family of el sidrón (Spain) support A
34 Role of Inbreeding in Neandertal extinction. *Nature Scientific Reports* 9(1): 1697
- 35 Roche H (2005) From Simple Flaking to Shaping: Stone-knapping Evolution among Early Hominids. In:
36 Roux, V, Brill, B (eds) *Stone knapping The necessary conditions for a uniquely hominin behavior*
37 *MacDonald Institute Monograph*, pp 35-53

- 1 Roebroeks W, Villa P (2011) On the earliest evidence for habitual use of fire in Europe. *Proceedings of*
2 *the National Academy of Sciences* 108(13): 5209-5214
- 3 Rohling EJ, Foster GL, Grant KM, Marino G, Roberts AP, Tamisie ME, Williams F (2014) Sea-level and
4 deep-sea-temperature variability over the past 5.3 million years. *Nature* 508 (7497): 477-482.
5
- 6 Rouchon V, Gillot PY, Quidelleur X, Chiesa S, Floris B (2008) Temporal evolution of the Roccamonfina
7 volcanic complex (Pleistocene), Central Italy). *Journal of Volcanology and Geothermal Research* 177:
8 500-514
- 9 Saccà D (2012) Taphonomy of *Palaeoxodon antiquus* at Castel di Guido (Rome, Italy): proboscidean
10 carcass exploitation in the Lower Palaeolithic. *Quaternary International* 276: 27-41
- 11 Sadori L, Koutsodendris A, Panagiotopoulos K, Masi A, Bertini A, Combourieu-Nebout N, Francke A,
12 Kouli K, Joannin S, Mercuri AM, Peyron O, Torri P, Wagner B, Zanchetta G, Sinopoli G, Donders TH
13 (2016) Pollen-based paleoenvironmental and paleoclimatic change at Lake Ohrid (south-eastern
14 Europe) during the past 500 ka. *Biogeosciences* 13: 1423–1437
- 15 Santucci E, Marano F, Cerilli E, Fiore I, Lemorini C, Palombo MR, Anzidei AP, Bulgarelli GM (2016)
16 *Palaeoxodon* exploitation at the Middle Pleistocene site of La Polledrara di Cecanibbio (Rome, Italy)
17 *Quaternary International* 406:169-182.
- 18 Scott B, Ashton N (2011) The Early Middle Palaeolithic: The European Context. In: Ashton N, Lewis
19 SG, Stringer C (eds), *The Ancient Human Occupation of Britain*. *Developments in Quaternary Science*,
20 pp 91–112
- 21 Segre GA, Naldini E, Muttoni G, Parenti F, Scardia G, Giacomo Segre A (2009) Nouvelles recherches
22 dans le bassin Plio-Pléistocène d’Anagni (Latium méridional, Italie). *Palevol* 113: 66–77
- 23 Schreve D, Moncel M-H, Bridgland D (2015) Editorial: The early Acheulean occupation of western
24 Europe: chronology, environment and subsistence behaviour In : Schreve D, Moncel M-H, Bridgland
25 D (eds). *Special issue: Chronology, paleoenvironments and subsistence in the Acheulean of western*
26 *Europe*. *Journal of Quaternary Science* 30(7): 585-593
- 27 Shackleton NJ (1987) Oxygen isotopes, ice volume and sea level. *Quaternary Science Reviews* 6 (3-4):
28 183-190.
29
- 30 Shipton C, Petraglia M D, Paddayya K (2009) Inferring aspects of Acheulean sociality and cognition
31 from lithic technology. In: B Adams and B Blades (eds). *Lithic materials and Paleolithic societies*, pp
32 219-231
- 33 Soriano S, Villa P (2017) Early Levallois and the beginning of the Middle Paleolithic in central Italy.
34 *PLoS One* 12(10): 1–28
- 35 Stringer C (2012) The Status of *Homo heidelbergensis* (Schoetensack 1908). *Evolutionary*
36 *Anthropology* 21: 101-107

- 1 Texier P-J, Roche H (1995) The impact of predetermination on the development of some Acheulean
2 chaînes opératoires In: Bermúdez de Castro, J, Arsuaga, JL, Carbonell, E (eds), Evolucion humana en
3 Europa y los yacimientos de la Sierra de Atapuerca Junta de Castilla y Leon, Valladolid, pp 403-420
- 4 Torre de la I (2016) The origins of the Acheulean: past and present perspectives on a major transition
5 in human evolution. *Philosophical Transactions B* 371(1698): 20150245
- 6 Torre de la I, Mora R (2018) Technological behaviour in the early Acheulean of EF-HR (Olduvai Gorge,
7 Tanzania). *Journal of human evolution* 120: 329, 377
- 8 Tuffreau A, Lamotte A, Marcy J L (1997) Land-use and site function in Acheulean complexes of the
9 Somme Valley. *World archaeology* 29(2): 225-241
- 10 Tzedakis PC, McManus JF, Hooghiemstra H, Oppo DW, Wijmstra TA (2003). Comparison of changes in
11 vegetation in northeast Greece with records of climate variability on orbital and suborbital
12 frequencies over the last 450 000 years. *Earth Planet. Sciences Letters* 212 (1): 197-212.
- 13 Tzedakis P C, Hooghiemstra H, Pälike H (2006) The last 1.35 million years at Tenaghi Philippon:
14 revised chrono - stratigraphy and long-term vegetation trends. *Quaternary Science Reviews* 25:
15 3416–3430.
- 16
- 17 Tzedakis PC, Wolff EW, Skinner LC, Brovkin V, Hodell DA, McManus JF, Raynaud D (2012) Can we
18 predict the duration of an interglacial? *Climate of the Past* 8 (5): 1473-1485.
- 19
- 20 Vaesen K, Collard M, Cosgrove R, Roebroeks W (2016) Population size does not explain past changes
21 in cultural complexity. *Proceedings of the National Academy of Sciences* 113(16): E2241-E2247
- 22 Vaesen K, Houkes W (2017) Complexity and technological evolution: What everybody knows? *Biology*
23 *& Philosophy*: 1-24
- 24 Villa P, Soriano S, Grün R, Marra F, Nomade S, Pereira A, Boschian G, Pollarolo L, Fang F, Bahain J-J
25 (2016) The Acheulean and Early Middle Paleolithic in Latium (Italy): Stability and Innovation. *PLoS*
26 *One* 11, e0160516
- 27 Wagner B, Vogel H, Francke A, Friedrich T, Donders T, Lacey JH, Leng, MJ, Regattieri E, Sadori L, Wilke
28 T, Zanchetta G, Albrecht C, Bertini A, Combourieu-Nebout N, Cvetkoska A, Giaccio B, Grazhdani A,
29 Hauffe T, Holtvoeth J, Joannin S, Jovanovska E, Just J, Kouli K, Kousis I, Koutsodendris A, Krastel S,
30 Lagos M, Leicher N, Levkov Z, Lindhorst K, Masi A, Melles M, Mercuri AM, Nomade S, Nowaczyk N,
31 Panagiotopoulos K, Peyron O, Reed JM, Sagnotti L, Sinopoli G, Stelbrink B, Sulpizio R, Timmermann A,
32 Tofilovska S, Torri P, Wagner-Cremer F, Wonik T, Zhang X, (2019) Mediterranean winter rainfall in
33 phase with African monsoons during the past 1.36 million years. *Nature* 573:256-260
- 34 Winton V (2005) An investigation of knapping-skill development in the manufacture of Palaeolithic
35 handaxes In: Roux, V, Brill, B (eds), *Stone Knapping: The Necessary Conditions for a Uniquely Hominin*
36 *Behaviour* McDonald Institute for Archaeological Research, Cambridge, pp 109e116
- 37 White M, Ashton N (2003) Lower Palaeolithic Core Technology and the origins of the Levallois
38 Method in North-Western Europe. *Current Anthropology* 44: 598–609

1 White M, Ashton N, Scott B (2011) The Emergence, Diversity and Significance of Mode 3 (Prepared
2 Core) technologies. In: Ashton N, Lewis SG, Stringer C (eds), The Ancient Human Occupation of
3 Britain. Quaternary Science 14, pp 67–91

4 Wiśniewski A (2014) The beginnings and diversity of Levallois methods in the early Middle
5 Palaeolithic of Central Europe. Quaternary International 326–327: 364–380

6

7

8

9

10

11

12

1 **Figure captions**

2 Figure 1: a) Reference map of the Latina Valley with the location of the archaeological sites of
3 Fontana Ranuccio, Colle Marino, Cava Pompei, Campogrande (CG), Selvotta (Sel), Colle Avarone (CA),
4 Isoletta, Lademagne, Masseria Castellone and Pignataro Interamna (modified from Pereira et al.,
5 2018). The main volcanic centres are also shown (triangles). b) Location of the archaeological sites
6 investigated in the study.

7 Figure 2. Simplified lithostratigraphic columns of the studied sites indicating the sampling position
8 and stratigraphic correlations between the archaeological levels. Campogrande (composite log from
9 Manzi et al., 2001; Manzi, 2016), Isoletta (Pereira et al., 2018), and Lademagne (modified from
10 Biddittu et al., 2012, Comerci et al., 2015) logs are from previous studies (see Pereira et al., 2018 for
11 more details).

12 Figure 3. Campogrande (CG) 10 lower. Core on a flint nodule with small bifacial removals on a part of
13 the periphery.

14 Figure 4. Campogrande 9 intermediate. Biface on a limestone flake.

15 Figure 5 Campogrande 10 upper: n°1: unifacial core on a limestone pebble, n°2: chopper core on a
16 small flint pebble, n°3: chopper or chopper core on limestone pebble, n°4: pointed pebble tool on a
17 small flint pebble

18 Figure 6. Campogrande 10 upper: asymmetrical elongated bifacial tool on limestone with a general
19 shaping. The middle part of the tool is the thickest part. Removals cover the surfaces and edges are
20 sinuous. This piece is an example of the chemical superficial alteration that makes it difficult to read
21 the shaping process. The organization of the removals is however visible.

22 Figure 7. Campogrande 10 upper: partial bifacial tools on limestone flake. The upper part of the tool
23 is shaped by small removals, including the tip. The base is left unworked.

24 Figure 8. Campogrande 10 upper: partial bifacial tool on a limestone flake. Some breccia on one face
25 makes it difficult the reading of the removals.

26 Figure 9. Colle Avarone: n°1: elongated and totally shaped bifaces on limestone, n°2: partial
27 triangular bifacial tool on a limestone flake, n°3: partial triangular bifacial tool on a flint flake

28 Figure 10. Colle Avarone: one or two series of removals on examples of pointed bifaces and tools
29 with a transversal cutting edge.

30 Figure 11. Selvotta: n°1: partial bifacial tool on a limestone flake, n°2: plano-convex bifaces on
31 limestone.

32 Figure 12. Selvotta: n°1, 4: elongated and plano-convex bifacial tool on limestone, n°2: partial bifacial tool
33 on a limestone cortical flake, n°3: bifacial tool on flint.

34 Figure 13. Isoletta 4: n°1: elongated bifaces on limestone, n°2: partial bifacial tool on a limestone
35 flake, n°3: lateral side of a backed bifacial tool with partial shaping.

36 Figure 14. Lademagne lower: Levallois or Levallois-like flake

- 1 Figure 15. Lademagne lower: n°1: elongated biface with a global shaping on limestone, n°2: backed
2 biface on limestone
- 3 Figure 16. Lademagne lower: n°1, 2: elongated biface with a global shaping on limestone, n°3: biface
4 with a global shaping on flint
- 5 Figure 17. Lademagne upper: n°1: partial backed bifacial tool on a limestone pebble, n°2: elongated
6 and plano-convex biface with a global shaping on limestone
- 7 Figure 18. Masseria Castellone: backed biface on limestone
- 8 Figure 19. Boxplot of the length of the bifaces and bifacial tools by locality.
- 9 Figure 20. Correspondence analysis of the assemblages (colored dots) and a set of factor states
10 characterizing technological features (non-colored dots), in particular the presence or absence of
11 cross-sectional or lateral symmetry (black dots) and the number of removals (grey dots).
- 12
- 13
- 14