



**HAL**  
open science

## Casablanca and the Earliest Occupation of North Atlantic Morocco

Jean-Paul Raynal, Fatima-Zohra Sbihi Alaoui, Lionel Magoga, Abderrahim Mohib, Mehdi Zouak

► **To cite this version:**

Jean-Paul Raynal, Fatima-Zohra Sbihi Alaoui, Lionel Magoga, Abderrahim Mohib, Mehdi Zouak. Casablanca and the Earliest Occupation of North Atlantic Morocco. Bulletin de l'Association française pour l'étude du quaternaire, 2002, 13 (1), pp.65-77. halshs-00004042

**HAL Id: halshs-00004042**

**<https://shs.hal.science/halshs-00004042>**

Submitted on 8 Jul 2005

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## CASABLANCA AND THE EARLIEST OCCUPATION OF NORTH ATLANTIC MOROCCO

Jean-Paul RAYNAL\*, Fatima-Zohra SBIHI ALAOUI\*\*, Lionel MAGOGA\*\*\*,  
Abderrahim MOHIB\*\*\*\* et Mehdi ZOUAK\*\*\*\*\*

### ABSTRACT

The Mio-Plio-Pleistocene sequence at Casablanca covers the last six millions years. The age estimates for different phases of this sequence have been established by various methods : lithostratigraphy, biostratigraphy, absolute dating (OSL, ESR), palaeomagnetism and aminochronology. Mio-Pliocene environments are represented by extremely rich palaeontological sites (Lissasfa, Ahl-Al-Oughlam), but these have not yet yielded hominid remains. The oldest lithic assemblages are found in Late Lower Pleistocene deposits, circa 1 Ma, in unit L of Thomas Quarry 1, and consist of Acheulean artefacts made from quartzite and flint. The first human remains discovered in this area were found in younger Middle Pleistocene deposits and cover an important period of human evolution spanning the time between *Homo erectus* and modern *Homo*. They are associated with Acheulean artefacts and rich faunal remains, recovered in caves (Littorines Cave at Sidi Abderrahmane, other caves at Thomas Quarries 1 and 3). The variability and the chronology of the Acheulean sequence is well documented following recent excavations in various sites around the well known locality of Sidi Abderrahmane (Bears Cave, Cap Chatelier, Unit L and Hominid Cave at Thomas Quarry 1, Rhinoceros Cave at Oulad Hamida Quarry 1, Sidi Abderrahmane Extension and Sidi Al Khadir open-air sites). The Casablanca sequence thus offers useful data for comparison with those from other African areas where hominids appeared and developed. They should also be considered in the debate on the earliest occupation of Europe.

**Key-words :** Morocco, Mio-Plio-Pleistocene, Casablanca, geofacts, Acheulean, quartzites, hominids.

### RÉSUMÉ

#### CASABLANCA ET LES PREMIÈRES OCCUPATIONS DU MAROC NORD-ATLANTIQUE

Casablanca est mondialement connu pour son patrimoine préhistorique. Des sites majeurs y ont été découverts dès le début du siècle dernier et certains ont livré des vestiges d'hominidés («Atlanthropes» de Sidi-Abderrahmane et des Carrières Thomas 1 et Oulad Hamida 1). Un programme de recherches y est conduit depuis 1978 et comporte, outre la révision des sites «classiques», de nouveaux travaux générés par l'urbanisation galopante et la nécessité d'exploiter et de sauvegarder au mieux ce patrimoine exceptionnel. La séquence de Casablanca est beaucoup plus longue qu'on ne le pensait il y a encore quelques années : elle a enregistré en détail l'évolution des milieux physiques et biologiques depuis près de 6 Ma et couvre donc - avec des gisements riches et bien datés - la totalité de la période qui a vu se dérouler l'évolution des Australopithèques et des premiers Hommes en Afrique orientale. Les sites paléontologiques de Lis-sasfa (environ 5,5 Ma), découvert en 1995, et d'Ahl Al Oughlam (anciennement carrière Déprez) (environ 2,5 Ma), découvert en 1985, ont livré des faunes mio-pliocènes de macro et micro-vertébrés qui précisent la chronologie régionale et permettent de mieux comprendre l'évolution du milieu animal d'Afrique du Nord et ses liens avec les autres grandes régions africaines. Les premières occupations sont rapportées à l'Acheuléen ancien et situées vers 1 Ma. Les premiers restes humains de cette région sont tous datés de différents moments du Pléistocène moyen et ont été découverts, associés à des outillages acheuléens, dans des cavités occupées également par des carnivores (Grotte des Littorines, grottes des carrières Thomas 1 et Thomas 3). La variabilité des outillages de la séquence acheuléenne et sa chronologie sont bien établies par les fouilles récentes de gisements variés, en grotte et en plein air, dans la zone de Sidi-Abderrahmane/Oulad Hamida (Grotte des Ours, Cap Chatelier, niveau L et Grotte de Thomas 1, Grotte des Rhinocéros, Sidi Abderrahmane Extension, Sidi Al Khadir).

**Mots-clés :** Maroc, Mio-Plio-Pléistocène, Casablanca, géofacts, Acheuléen, quartzites, hominidés.

\*Université de Bordeaux 1, Institut de Préhistoire et de Géologie du Quaternaire, UMR 5808 CNRS, Avenue des Facultés, 33405 TALENCE cedex (France) - Mission «Littoral» et équipe Casablanca. E-mail jpraynal@wanadoo.fr

\*\*Institut National des Sciences de l'Archéologie et du Patrimoine, Avenue John Kennedy, Rabat-Souissi, Maroc et équipe Casablanca

\*\*\*03200 Vichy (France), Mission «Littoral» et équipe Casablanca.

\*\*\*\*Direction du Patrimoine, Inspection des Monuments et des Sites, Musée archéologique de Tétouan (Maroc), équipe Casablanca.

\*\*\*\*\*Direction du Patrimoine, Inspection des Monuments et des Sites, El Jadida (Maroc), équipe Casablanca.

## INTRODUCTION

The city of Casablanca (fig. 1) played an important role in African prehistory as soon as it began to grow and develop at the beginning of the 20th century. Numerous quarries were opened and several major archaeological finds occurred until the sixties, including discovery of human remains (Lecointre, 1926 ; Neuville et Ruhlmann, 1941 ; Biberson, 1956, 1961 ; Ennouchi, 1969, 1972). This heritage is now quickly disappearing under the modern city and archaeologists have to race with builders to protect and excavate the sites (Raynal et Geraads, 1993 ; Raynal, 1998 ; Sbihi-Alaoui et Mohib, 1998). The joint French-Moroccan program «Casablanca» has

been undertaken twenty-four years ago to excavate sites exposed to destruction, to revise the stratigraphical frame of this major zone of African prehistory with modern methods and to reconstruct palaeoenvironments of early hominids.

We have now clearly established that the long sequence of Casablanca covers the last six millions years and constitutes an exceptional record of sea level variations (Raynal *et al.*, 1999). The age estimates for different phases of this sequence have been established by various methods : lithostratigraphy (Texier *et al.*, 1994, 2002), biostratigraphy (Geraads, 2002), absolute dating (OSL, ESR) (Rhodes, 1990 ; Rhodes *et al.*, 1994), palaeomagnetism and aminochronology (Occhietti *et al.*, 1993,




Fig. 1 : Location map.  
Fig. 1 : Carte de localisation.




Fig. 2 : Casablanca, Thomas 1 Quarry, Unit L1, classical trihedron, quartzite.  
 Fig. 2 : Casablanca, carrière Thomas 1, Unité L1, trièdre classique en quartzite.

1996, 2002). Detailed studies have specified the evolution of the palaeoenvironment (El Graoui, 1994 ; Lefèvre *et al.*, 1985, 1994, 1996 ; Lefèvre, 2000). Recent archaeological studies and synthesis have contributed to establish the evolution of human activities in the area (Raynal *et al.*, 1995, 2001, 2002 ; Mohib, 2001). We will consider here only some of these aspects.

The biochronological framework of the Moroccan Quaternary is now much more firmly established than it was ten years ago (Geraads, 2002). It can now be compared to the East African one, and some periods or aspects are even better sampled. It still needs refinements and could indeed be improved for some parts, especially the transition between Middle and Upper Pleistocene. We shall focus below on some of the key-sites, roughly in decreasing ages, and then come back to the characteristics and evolution of lithic assemblages.

Mio-Pliocene environments are characterized within extremely rich palaeontological sites : Lissasfa around 5.5 Ma (Geraads, 1998 ; Raynal *et al.*, 1999 ; Geraads, 2002), Ahl-Al-Oughlam around 2.5 Ma (Raynal *et al.*, 1990 ; Geraads *et al.*, 1998 ; Geraads, 2002). These sites have not yet yielded hominids remains nor artefacts, but only geofacts : up to now, there is no Oldowayan-like industry clearly documented and *in situ* within Pliocene and Lower Pleistocene sediments of Atlantic Morocco.

## 1 - THE PLEISTOCENE KEY-SITES

### 1.1 - THOMAS QUARRY 1 UNIT L

The oldest lithic assemblages were found in Late Lower Pleistocene deposits, circa 1 Ma, in unit L of Thomas Quarry 1, and consist of Acheulean artefacts made of quartzite and flint (Raynal et Texier, 1989). The assemblage described *infra* has been recovered in a 80 m<sup>2</sup> excavation but 1000 m<sup>2</sup> of layer L have been exposed and are now under excavation. The lithic series contains flakes struck from discoidal cores and polyhedrons. Besides chopping-tools, polyhedrons, and some cleavers, trihedrons and bifaces form the most characteristic elements among the tools (fig. 2). They are often only partially flaked and usually display lateral or lateral-distal concavities, which form the point of the bifaces (fig. 3). A comparison of the probable use of the objects identified in our classification with experimental results leads us to conclude that the activities of stone working, hide slitting, heavy-duty butchery, and bone breaking were probably performed in unit L1, while stone working and light-duty butchery characterise unit L5.

The lowermost layers of Thomas 1 quarry have yielded a few large Mammals, left over by *Homo erectus* as food refuse, and rather badly preserved. After Geraads (*in Raynal et al.*, 2001), Hippo, Zebra and Gazella have little




Fig. 3 : Casablanca, Thomas 1 Quarry, Unit L1, partial biface on quartzite flat pebble.  
 Fig. 3 : Casablanca, carrière Thomas 1, Unité L1, biface partiel sur galet plat de quartzite.

biochronological signification, but the discovery in 1996 of a *Kolpochoerus* third molar, suggests a great antiquity. This Suid, widespread in Eastern and Southern Africa, was previously known in North Africa only in Plio-Pleistocene sites, the youngest of them being Aïn Hanech, a locality which is certainly of lower Pleistocene age, perhaps close to 1.2 Ma. *Kolpochoerus* is absent from all other level in Thomas/Oulad Hamida quarries, and from Tighenif, an Algerian locality with an age close to the Lower/Middle Pleistocene boundary (Geraads *et al.*, 1986).

Micromammals (Geraads *in* Raynal *et al.*, 2001), although also rare, tend to confirm the great age of level L. There are two species of *Gerbillus*, the larger one is of the size of *G. grandis*, known in other levels of the Thomas 1 quarry but with a very broad M/1, and it is certainly specially distinct, the smaller species is also distinct from the similar *G. minutus*, *G. jebiletii* or *G. campestris*. A lower M/1 of *Paraethomys* has the anterior lobe quite transversal instead of V-shaped as in all other Middle Pleistocene species, and it looks more like *P. mellahe* AMEUR, from Oued Mellah, a locality older than Tighenif. The absence of *Ellobius*, an Arvicolid present at Tighenif also supports an age greater than that of this locality.

From a global point of view, the fauna from level L demonstrates its lower Pleistocene age, suggesting an age of perhaps 1 Ma or more, in full accordance with typology and technology of the artefacts, with the paleomagnetic data obtained by Sevket Sen (*in* Raynal *et al.*, 1996), and with the OSL dates for unit L5 (Rhodes *et al.*, 2002) :  $989 \pm 208$  ka,  $1683 \pm 473$  ka,  $1037 \pm 1204$  ka, even if they have very large uncertainties...). The presence of regular polyhedrons (spheroids), as well as irregular ones, both similar to those described in the Algerian locality at Aïn Hanech (Sahnouni et Hadjouis, 1987) supports this opinion. In this site, the absence of bifacials could only be a matter of variability related to the surface excavated and has to be confirmed by larger excavations. On the whole, the lithic assemblage from Thomas 1 quarry unit L is contemporaneous and similar to those of Lower Acheulean from East Africa and Middle East.

#### 1.2 - OULAD HAMIDA 1 RHINOCEROS CAVE AND THOMAS QUARRY HOMINID CAVE

Middle Pleistocene levels are represented in the Oulad Hamida 1 quarry, where the Rhino cave has yielded, in the lower part of the filling, a very rich collection of micro and macromammals that indicate a rather open

and dry environment, associated with a rich Acheulean assemblage (Raynal *et al.*, 1993 ; Geraads, 1993, 1994 ; Bernoussi, 1994, 1997). This occupation has been dated by ESR (Rhodes *et al.*, 1994) and new calculations provide older dates :  $435 \pm 85$  ka by early uptake and  $737 \pm 129$  ka by linear uptake (Rhodes *et al.*, 2002). They are in better agreement with biostratigraphical data (Geraads, 2002), and with unpublished lithostratigraphical observations which point towards a 0.6 Ma minimum age. This site is now one of the best reference levels for this period of Middle Acheulean in North-Africa. The abundant remains of white rhinoceros suggest specialised hunting by hominids. Compared to the local early Acheulean, one can observe an increase of discoid cores and of flake production ; cleavers are rare while bifacial pieces are larger - even if one can observe a large variability in sizes - and characterized by convex and/or concave edges that constitute a pointed extremity (figs. 4 and 5). In

total, 3485 artefacts were removed from the lower layer of a 70 m<sup>2</sup> excavation. Tools on flakes represent only 3.5% of the assemblage and notches and denticulates predominate, the others being different types of scrapers and some multiple tools. Stone working was obviously important in this site, as probably were hide cutting, light- and heavy-duty butchery and bone breaking.

The assemblage recovered in a 65 m<sup>2</sup> excavation of the Hominid level at Thomas Quarry I is quite different from the series of the Rhino Cave bottom unit, as it is dominated by flaked pebbles. It is, however, comparable to the (too) small series from the Rhino Cave top unit and to series collected at the time of the discovery of the *Homo* jaw in 1969. Geological studies by Texier demonstrate that this assemblage was not in primary position but has been secondarily introduced into the cave, presumably removed from the entrance and transported by run-off, which mixed fauna remains and artefacts. The macro-




Fig. 4 : Casablanca, Oulad Hamida 1 Quarry, Rhinoceros cave, cleaver, quartzite.  
Fig. 4 : Casablanca, Oulad Hamida 1 Quarry, Grotte des Rhinocéros, hachereau en quartzite.




Fig. 5 : Casablanca, Oulad Hamida 1 Quarry, Rhinoceros cave, small lanceolate bifacial, quartzite.  
 Fig. 5 : Casablanca, Oulad Hamida 1 Quarry, Grotte des Rhinocéros, petit biface lancéolé en quartzite.

fauna is similar in composition to that of Rhino Cave, although dominated here by carnivores (bears, hyenas and *Canis*) (Bernoussi, 1997), but it is associated with a similar macrofauna to that from Rhino Cave (Geraads, 1980 ; Geraads *et al.*, 1980 ; Bernoussi, 1997). Three large sized and robust teeth of *Homo* have recently been recovered in this cave, only a few meters from the place where the mandible was discovered in 1969. They are very similar to other specimen of *Homo erectus* from North Africa (Zouak, unpublished). This Thomas Quarry 1 Hominid Cave is vast, and some parts of the stratigraphy are still under study (fig. 6). The preliminary dates, as well as biostratigraphy, and lithostratigraphical data point towards a greater antiquity than was previously estimated, with a minimum age of 0.6 Ma.

After Geraads (*in* Raynal *et al.*, 2001), the relatively great antiquity of these two faunal sets is shown by their similarities with Tighenif, and by the occurrence of several extinct species. Among the taxa shared with the

Algerian site are probably the zebra *Equus mauritanicus*, the giant baboon *Theropithecus oswaldi* and several species of poor biochronological significance. The OHI-Th1 faunal unit no longer has some lower Pleistocene survivors still present at Tighenif, such as the Suid *Metridiochoerus*, the antelope *Hippotragus gigas*, or the sabre-tooth cat *Homotherium*, but *Canis*, the ratel *Mellivora*, the lynx, the antelope *Parmularius* and the hare *Serengetilagus* are all of extinct (mostly new) species. Micro-mammals allow refinement of the biochronological succession within this faunal unit. The best collection is from the Rhinoceros Cave (Geraads, 1994) and samples have also been collected in several levels or spots of the Thomas 1 quarry. The Rodent fauna of this cave system, which is rather sharply distinct from that of level L, is dominated by the Gerbillids, of which there are at least 4 species, confirming the aridity suggested by the abundance of Alcelaphines, and gazelles among bovids. Murids (*Paraethomys*, *Praomys* and *Mus*), Arvicolid

(*Ellobius*) and Glirids (*Eliomys*) are less common. The earliest level is the pink breccia at the base of the Thomas 1 cave, from where most probably comes the *Homo erectus* mandible. It has an *Eliomys* smaller than at OH1-GDR, and also a small, perhaps new, species of the Gerbillid genus *Meriones*.

Stratigraphic elements and absolute datings thus allow to place these levels circa 0.6/0.7 Ma. This gives a new palaeoanthropological perspective for the human remains of evolved *Homo erectus* or archaic *Homo sapiens* (Hublin, 1991), which were recovered in Thomas 1 quarry cave and in Thomas III quarry cave, unhappily destroyed. Since they are the oldest discovered in Casablanca : they cannot be linked with more recent fossils like those from Sidi Abderrahmane Littorines Cave, potentially contemporaneous of the end of isotopic stage 11 at most, and moreover with the Salé skull, which was not discovered during excavations, and which stratigraphic position has never been clearly established.

### 1.3 - OTHER SITES

At Sidi Al Khadir-Hélaoui Quarry, an open-air site has been excavated. Flake production was the main activity on this site, may be in relationship with light-duty butchery. This assemblage represents a facies of Middle Acheulean without bifacials, with an isotopic stage 16 minimum age (Lefèvre, 2000).

In Bears Cave at Sidi Abderrahmane, a recent stage of the Middle Acheulean is illustrated, dating of the boundary of isotopic stages 12 and 11. A 64 m<sup>2</sup> recent excavation in the cave deposits has given a large series of stone objects (2976) and a limited series of faunal remains (91), mainly represented by *Ursus* bones fragments. Despite the high degree of marine reworking, the lithic assemblage in secondary position still reveals every step of the operative chains, from the raw material (quartzite pebbles and blocks form 97% of the lithic resources), to the

ultimate retouched tools (Mohib, 2001). Flaking, retouch and reduction processes are illustrated by different types of flakes, fragments and various cores : besides polyedric ones, we have identified Clactonian chopper-cores, cores on flakes and very large discoid cores. Bifaces are often made on flakes and show an asymmetric profile.

The upper part of the Acheulean sequence is well illustrated at Cap Chatelier (Sidi-Abderrahmane Cunette) and at Sidi-Abderrahmane-Extension, and possibly for its basal part at Sidi Abderrahmane Littorines Cave, although it's assemblage has not been re-examined yet and it's exact stratigraphic position remains unclear, despite Biberson's explanations (Biberson, 1956).

The assemblage of Cap Chatelier demonstrates production of predetermined flakes and thin small bifaces, a diverse set of tools on flakes and a very few cleavers. It is older than isotopic stage 9 after new OSL dates obtained at the top of Cap Chatelier section in «small dune» of Biberson (1961) : 376 ± 34 ka and 317 ± 64 ka (Rhodes *et al.*, 2002).

At Sidi-Abderrahmane Extension we observe an important use of block-fragments and frequent recycling of rolled artefacts with multiple scars. Flakes are mainly produced from discoid cores and polyhedral forms are quite rare. So they are predetermined flakes, but these coexist with a diverse toolkit on flakes. Bifacial pieces are generally made on flakes and mostly display convex sides and tend towards ovate forms, even to discoid ones. Cleavers are rare here, as if they had become useless along with the increase of new bifacial patterns. Stone working was probably important at this site, along with hide cutting, light-duty butchery and bone breaking. This assemblage is neatly younger than isotopic stage 9 and can be situated either in isotopic stage 8 or 7, more likely 7 and then contemporaneous of the Bir Feghloul member of the Kef El Haroun Formation if we follow the stratigraphic relationships proposed by Lefèvre (2000).




Fig. 6 : Thomas Quarry 1, Hominid Cave stratigraphy.  
Fig. 6 : Casablanca, carrière Thomas 1, stratigraphie de la Grotte à Hominidés.



This Upper Acheulean is the technical reservoir which, far before the Last Interglacial, developed into pre-Mousterian and Mousterian facies associated with modern humans, as demonstrated at Djebel Irhoud for example, where fauna collected by Ennouchi has recently been revised (Amani, 1991 ; Amani et Geraads, 1993 ; Geraads et Amani, 1998b), and points towards an age close to the Middle / Upper Pleistocene boundary (Geraads, 2002).

## II - THE LITHIC PERSPECTIVE IN THE ACHEULEAN SEQUENCE OF CASABLANCA

The raw material was abundantly available in all sizes in the area and allowed a production of large flakes or voluminous fragments of pebbles and blocks. The only constraint consisted in the transport of heavy objects, making voluminous flaked items rare in the excavated sites. The different types of blanks introduced to the sites are very well recognisable in the bifaces of the various series.

The exploitation of the same raw materials throughout the Acheulean sequence at Casablanca allows a comparison of the technological characteristics of some representative series, presented here along the lines of the classification developed on the basis of a study of the assemblages from Unit L of the Thomas quarry 1, briefly explained below (Raynal *et al.*, 2002). This model is based on the character of the working surfaces and on their disposition and exploitation. It integrates dynamic aspects (sequences of flakes production and shaping, reduction of objects, re-use, etc.) and functional ones (specific morphology, transformation by use, etc.). Seven main groups are discerned :

- Group 1 : flaking carried out by using cortical striking platforms ;
- Group 2 : flaking from one non-cortical striking platform, possibly re-adjusted ;
- Group 3 : flaking using two non-cortical striking platforms for one and the same working surface ;
- Group 4 : flaking using three to five non-cortical striking platforms for one and the same working surface. It contains most objects with multiple flake removals. Starting with flaking from cortical striking platforms and followed by an increase of flaked

surfaces, this group contains the majority of complex and/or typical objects. The most complete bifaces as well as the best-exploited cores occur within this group ;

- Group 5 : flaking from non-cortical striking platforms belonging to various working surfaces ;
- Group 6 : usage/shaping of flakes and fragments ;
- Group 7 : objects transformed by utilisation. With cores on flakes and fragments and tools on flakes, this testifies to the final stages of the knapping process. Items transformed by usage or re-utilisation are at the origin of part of the observed assemblage variability.

Pebbles and flakes are subdivided in the followings :

- M0 : intact pebble.
- M1 : naturally broken pebble, splitted pebble, pebble with a single flake removal, pebble with several non-adjacent flake removals.
- M2 : block or blocks parts.
- M3 : flaking accidents : pebbles broken during flaking or use, splitted sub-spheroids, broken hammer-stones.
- M4 : flakes, chunks, pebble fragments, core fragments.

Among flakes, different technical categories are identified, corresponding to operative chains for bifacial production, discoid core reduction, polyhedral flaking...

The group subdivisions, (expressed by letters – A, B, C, D... in table 1 but not detailed in this paper), thus rest on technological and/or secondary morphological criteria (such as re-use of striking platforms, recurrence of flake removals, length of blanks both artificial and natural, surface size of flake removals, and presence/absence of cortex). A comparison can be made with categories defined by Chavaillon and Chavaillon (1981) and demonstrates the technological variability among them (tab. 1).

The variability of lithic assemblages is well documented, following recent excavations in Thomas Quarry 1 units L1 and L5 (Th1/L1, TH1/L5), Thomas Quarry 1 Hominid Cave (TH1/GH), Rhino Cave bottom unit (GDR), Bear Cave (GDO), Cap Chatelier, Sidi Abderrahman Extension (SAE), and Sidi Al Khadir open-air sites (table 2, fig. 7). The series demonstrate an alternation between industries rich or not in bifacial pieces but they are all supposed to belong to the Acheulean technocomplex. This situation is like the one observed everywhere in Africa and in Europe. This variability occurs sometimes during a short duration of sedimentary events, i.e. Unit L at Thomas Quarry 1 (fig. 8).

But on another hand, similarities and differences in the structure of the various assemblages may actually derive from other factors. Among these, one can nevertheless discard the access to raw materials ; sources are the same along the sequence and the composition of assemblages is remarkably stable from this point of view. El Hank quartzites are dominant, available in pebbles or chunks, and flint, only available in small pebbles, is much less flaked. Thus, the stability of technological solutions observed along the sequence derives evidently from raw

J. and N. Chavaillon (1981) :	Classification in use
Pebble tools	1A, 2A, 2B1, 3, 4C, M1, M3
Cores	1A1, 2B1, 3, 4B, 4C, 5A, 6A, M4
Choppers	1A, 2A, 2B1, 3, 4C, M1
Hammerstones	7B, 7C, 7D
Broken pebbles	M1, M3, 7C
Polyhedrons	5A even 5B, 7A
Bolas	7A

Tab. 1 : Comparison between the classification used in this paper and the one of J. Chavaillon.

Tab. 1 : Comparaison entre la classification utilisée dans cet article et celle de J. Chavaillon.




Fig. 7 : Variability of acheulian assemblages of Casablanca.  
 Fig. 7 : Variabilité des outillages acheuléens de Casablanca.




Fig. 8 : Distribution of technical groups in archaeological units L1 and L5 at Thomas Quarry 1.  
 Fig. 8 : Répartition des groupes techniques dans les unités archéologiques L1 et L5 de la carrière Thomas 1.

Sites Age OIS*	TH1/L1 ± 1.0 Ma 23 ?		TH1/L5 ± 1.0 Ma 23 ?		TH1/GH 0.6/0.7 Ma ? >15		GDR base 0.6/0.7 Ma ? > 15		GDO ± 0.4 Ma 11		SAE ≥ 0.2 Ma ≥ 7	
	n	%	n	%	n	%	n	%	n	%	n	%
1	18	6,0	18	4,8	40	54,8	66	11,3	104	51,2	31	5,4
2	56	18,8	79	21,2	17	23,3	82	14,0	15	7,4	62	10,7
3	38	12,8	11	2,9	9	12,3	46	7,8	9	4,4	96	16,6
4	124	41,6	61	16,4	7	9,6	22	3,8	23	11,3	199	34,5
5	18	6,0	66	17,7	0	0,0	59	10,1	27	13,3	5	0,9
6	36	12,1	97	26,0	0	0,0	156	26,6	25	12,3	181	31,4
7	8	2,7	41	11,0	0	0,0	155	26,5	0	0,0	3	0,5
total	298	100,0	373	100,0	73	100,0	586	100,0	203	100,0	577	100,0

\* oxygen isotopic stage

Tab. 2 : Technical composition of the major sites of the acheulian Casablanca sequence after recent excavations, following the classification of Raynal, Magoga et Sbihi-Alaoui (in Bulletin d'Archéologie Marocaine, t. 19). Th1 L : Thomas 1 unit L ; Th1 GH : Thomas 1 Homidid Cave ; GDR : Rhinos Cave Oulad Hamida 1 Quarry ; GDO : Bears Cave at Sidi Abderrahmane ; SAE : Sidi Abderrahmane Extension.

Tab. 2 : Inventaire technique des différents sites majeurs de la séquence acheuléenne de Casablanca (fouilles récentes) selon le classement de Raynal, Magoga et Sbihi-Alaoui (in Bulletin d'Archéologie Marocaine, t. 19, sous presse). Th1 L : carrière Thomas 1 niveau L ; Th1 GH : carrière Thomas 1 grotte à hominidés ; GDR : grotte des Rhinocéros carrière Oulad Hamida 1 ; GDO : grotte des Ours à Sidi Abderrahmane ; SAE : Sidi Abderrahmane Extension.




Fig. 9 : Correlation between the number of flake removals and the number of striking platforms on quartzite artefacts from key-sites of the Casablanca Acheulean sequence.

Fig. 9 : Corrélation entre nombre d'enlèvements et nombre de plans de frappe sur les objets de quelques sites-clés de la séquence acheuléenne de Casablanca.

materials quality and the variability derives from other factors. The correlation between the number of flake removals and the number of striking platforms is a technological constant (a succession of x flakes originating from one platform), verified in experiments and determined by the mechanical characteristics of the raw material (fig. 9).

Bifacial pieces also bear witness of an increasing complexity of elementary modes of reduction and of a continuous technological enrichment in which earlier acquisitions re-appear (the polyhedral proximal part for example) (fig. 10). The changes within the bifaces reflect in our view a morpho-functional evolution, relayed in time by more systematic production of flakes, including predetermined ones. There is a global trend towards

biconvex symmetric shapes which clearly reflects a functional specialization (fig. 11).

Some factors of variability might have had a natural origin such as for instance, the selection of materials by re-depositional actions, which has taken place in some layers. For example, Layer L1 at Thomas Quarry where small flakes and bone fragments have been washed or concentrated and where the fabric of bigger artefacts clearly indicate a water-flow action. On the other hand, the smallest flakes are preserved in aeolian sands of Layer L5.

Another factor is the limited area of some excavations : for instance the area excavated in Layer L5 at Thomas Quarry 1 does not exceed 25 square metres... Finally, possible cultural reasons for the various inter-relationships




Fig. 10 : Morphology of proximal parts of bifacial pieces. 1 : natural or unretouched. 2 : cutting. 3 : polyedric.  
 Fig. 10 : Morphologie des parties proximales des pièces bifaciales. 1 : naturelle ou non retouchée. 2 : tranchante. 3 : polyédrique.




Fig. 11 : Compared morphology of bifacial pieces in key-sites of the Casablanca Acheulean sequence. TH1 L1 : Thomas Quarry 1 unit L1. GDR : Rhinoceros Cave Oulad Hamida 1 Quarry. GDO : Bears Cave at Sidi Abderrahmane. SAE : Sidi Abderrahmane Extension.  
 Fig. 11 : Morphologie comparée des pièces bifaciales des sites-clés de la séquence acheuléenne de Casablanca. TH1 L1 : carrière Thomas 1, niveau L1. GDR : grotte des Rhinocéros, carrière Oulad Hamida 1. GDO : grotte des Ours à Sidi Abderrahmane. SAE : Sidi Abderrahmane Extension

between the industries must of course be considered. Whenever the use of raw materials is not in question, there may be a certain variability connected with peculiar functional situations, such as adaptive reactions to environmental and/or to microenvironmental changes resulting from coincidence with limited or more global climatic changes. Evidence for such variability in assemblages according to climatic fluctuations within a short time span is illustrated by Layers L5 and L1 of Unit L at Thomas Quarry 1.

Variability in assemblages from sub-contemporaneous cave deposits at Thomas Quarry 1 and Oulad Hamida Quarry 1, may be considered from another point of view : different carnivoras have acted in these sites as hunters and scavengers, different meat resources were offered, different tool kits were employed...

At the end, among the progressive transformations of assemblages, some thresholds appear, which could be linked with human evolutionary steps or reflect different biomechanical solutions. They support the subdivisions of local Acheulean in several main stages, which is nevertheless subject to re-examination with further informations.

### CONCLUSION

There is still a lot to investigate : Lower Pleistocene sites beyond 1.0 Ma are still unknown in Morocco and the detailed chronological frame needs more refinements. Nevertheless, the Casablanca Acheulean sequence is now better documented with main subdivisions founded on recent excavations and clearly contemporaneous of those known from East Africa. The main differences with Biberson's proposals (1963) are :

- Stades I and IV of Moroccan «Pebble-culture» have been respectively defined at Casablanca on the basis of geofacts and of reworked recent assemblages rich in pebble tools (Raynal et Texier, 1989). Up to now, there is no Oldowayan site identified in stratigraphic position below the Acheulean sequence,

- «Lower Acheulean» assemblages are now considered as Middle Acheulean,

- «Upper Acheulean» is much older at Cap Chatelier than supposed before and on the contrary, much more recent at Sidi Abderrahmane Extension.

The succession of archaeological layers, from the oldest to the youngest, appears today as following :

- Lower Acheulean, isotopic stage 23 (?) : Thomas Quarry 1 L1, Oulad Hamida 1 unit L, Thomas Quarry 1 L5,

- Middle Acheulean, isotopic stage 16 as a minimum age : Thomas Quarry 1 Hominid Cave, Oulad Hamida 1 Rhinos Cave, Sidi Abderrahmane unit M, STIC Quarry, Sidi Abderrahmane Bears Cave at isotopic stages 12 to 11 transition,

- Upper Acheulean, from isotopic stage 11 to isotopic stage 8 or 7 at least : Sidi Abderrahmane Littorines Cave (?), Sidi Abderrahmane Cap Chatelier, Sidi Abderrahmane Extension.

Nevertheless, this Acheulean sequence offers useful data for detailed comparisons with other African areas, where hominids appeared and developed, as for example

the Oldoway succession from Bed III to Nduku Beds (Leakey, 1951, 1971 ; Leakey and Roe, 1994). Indeed, within these beds of the Oldoway sequence, or even in the Melka Kunture series (Chavaillon *et al.*, 1978), we can observe the same variability between Acheulean and Developed Oldowayan facies, according to site function and abundance/scarcity of artefacts.

North African Acheulean should now be reconsidered in the debate about the first occupation of Europe and especially the question of multiple «Out of Africa» hypothesis through Mediterranean straits and isthmus. One million years ago, *Homo* supposed *erectus* - but we really don't know yet - who manufactured Lower Acheulean lithic assemblages, was facing Southern Europe and nobody can firmly exclude its possible desire and ability for crossing the waters of narrow straits...

### ACKNOWLEDGEMENTS

The authors thanks Peter Bindon of the Australian Ethnographic Institute, who revised the English version of this paper, the *Institut National des Sciences de l'Archéologie et du Patrimoine du Royaume du Maroc*, *Ministère des Affaires Etrangères* and the *Région Aquitaine* for supporting and funding the *Mission archéologique française «Littoral» Maroc*, and especially the Casablanca program.

Autorisation de publier n° 03/11-02 du 20 Novembre 2001.

### REFERENCES

- AMANI, F., 1991 - *La faune de la grotte à Hominidé du Jebel Irhoud (Maroc)*. Thèse, Université de Rabat, 229 p.
- AMANI, F. et GERAADS, D., 1993 - Le gisement moustérien du Jebel Irhoud, Maroc : précisions sur la faune et la biochronologie, et description d'un nouveau reste humain. *Comptes Rendus de l'Académie des Sciences*, Paris, II, 316, 847-852.
- BERNOUSSI, R., 1994 - *Etude d'une taphocénose Pléistocène : exemple de la Grotte des Rhinocéros (Carrière Oulad Hamida I, Casablanca)*. DEA, Université de Bordeaux I, 101 p.
- BERNOUSSI, R., 1997 - *Contribution à l'étude paléontologique et observations archéozoologiques pour deux sites du Pléistocène moyen du Maroc atlantique : grotte à Hominidés de la carrière Thomas 1 et de la grotte des Rhinocéros de la carrière Oulad Hamida I (Casablanca, Maroc)*. Thèse de l'Université de Bordeaux I, n° 1711, 263 p.
- BIBERSON, P., 1956 - Le gisement de l'Atlantrophe de Sidi-Abderrahmane. *Bulletin d'Archéologie Marocaine*, 1, 38-92.
- BIBERSON, P., 1961 - *Le cadre paléogéographique de la Préhistoire du Maroc atlantique et Le Paléolithique inférieur du Maroc atlantique*. Publications du Service des Antiquités du Maroc, Rabat, fasc. 16 et 17.
- BIBERSON, P., 1963 - Human evolution in Morocco in the framework of the atlantic Pleistocene. *African ecology and human evolution*, Chicago, 417-447.
- CHAVAILLON, J. et N., 1981 - Galets aménagés et nucléus du Paléolithique inférieur. *Mélanges offerts au Doyen L. Balout*, ADPF, Paris, 283-292.
- CHAVAILLON, J., CHAVAILLON, N., HOURS, F. et PIPERNO, M., 1978 - Le début et la fin de l'Acheuléen à Melka-Kunturé : méthodologie pour l'étude des changements de civilisation. *Bulletin de la Société préhistorique française*, 75, 4, 105-115.
- EL GRAOUI, M., 1994 - *Contribution à l'étude des formations littorales quaternaires de la région de Casablanca (Maroc) : sédimentologie, microfaciès et minéraux lourds*. Thèse de l'Université de Bordeaux I, n° 1100, 263 p.
- ENNOUCHI, E., 1969 - Découverte d'un Pithécantropien au Maroc. *C. R. Acad. Sci. Paris*, série D, p. 763.
- ENNOUCHI, E., 1972 - Nouvelle découverte d'un Archanthropien au Maroc. *C. R. Acad. Sci. Paris*, t. 274, série D, 3088-3090.

- GERAADS, D., 1980 - La faune des sites à *Homo erectus* des carrières Thomas (Casablanca, Maroc). *Quaternaria*, 22, 65-94.
- GERAADS, D., HUBLIN, J.J., JAEGER, J.J., TONG, H., SEN, S. & TOUBEAU, P., 1986 - The Pleistocene Hominid of ternifine, Algeria : New results on the environment, age and industries. *Quaternary Research*, 25, 380-386.
- GERAADS, D., BERIRO, P. et ROCHE, H., 1980 - La faune et l'industrie des sites à *Homo erectus* des carrières Thomas (Maroc). Précisions sur l'âge de ces hominidés. *C. R. Acad. Sci. Paris*, t. 291, série II, 195-198.
- GERAADS, D., 1993 - Middle Pleistocene *Crocidura* (*Mammalia*, *Insectivora*) from Oulad Hamida 1, Morocco, and their phylogenetic relationships. *Proc. Kon. Ned. Akad. v. Wetensch.* 96 (3), 281-294.
- GERAADS, D., 1994 - Rongeurs et Lagomorphes du Pléistocène moyen de la «Grotte des Rhinocéros», Carrière Oulad Hamida 1 à Casablanca, Maroc. *N. Jb. Paläont. Abh.*, 191, 2, 147-172.
- GERAADS, D., 1998 - Rongeurs du Mio-Pliocène de Lissasfa (Casablanca, Maroc). *Geobios*, 31, n°2, 229-245.
- GERAADS, D., AMANI, F., RAYNAL, J.P. et SBIHI-ALAOUI, F.Z., 1998 - La faune de mammifères du Pliocène terminal d'Ahl al Oughlam, Casablanca, Maroc. *C. R. Acad. Sci. Paris*, t. 326, 671-676.
- GERAADS, D. et AMANI, F., 1998 - Le gisement moustérien du Djebel Irhoud, Maroc : précisions sur la faune et la paléoécologie. *Bulletin d'Archéologie marocaine*, Rabat, 18, 11-18.
- GERAADS, D., 2002 - Quaternary Mammalian Biostratigraphy of Atlantic Morocco, *Quaternaire*, ce volume.
- HUBLIN, J.J., 1991 - *L'émergence des Homo sapiens archaïques : Afrique du Nord-Ouest et Europe occidentale*. Thèse d'Etat de l'Université de Bordeaux 1.
- LEAKEY, L.S.B., 1951 - *Olduvai Gorge*. Cambridge University Press, 164 p.
- LEAKEY, M.D., 1971 - *Olduvai Gorge. Volume 3. Excavations in Beds I and II, 1960-1963*. Cambridge University Press, 306 p.
- LEAKEY, M. and ROE, D., 1994 - *Olduvai Gorge, excavations in beds III, IV and the Masek beds 1968-1971*. Olduvai Gorge, vol. 4, Cambridge University Press, 323 p.
- LECOINTRE, G., 1926 - *Recherches géologiques dans la Meseta marocaine*. Mémoire de la Société des Sciences naturelles du Maroc, n° XIV.
- LEFEVRE, D., RAYNAL, J.P. et TEXIER, J.P., 1985 - De la fin du Villafranchien au début du Soltanien : exemple d'évolution des paléoenvironnements du Maroc occidental et oriental. *Colloque de l'Association des Géographes français*, Paris, novembre 1985, diffusion restreinte.
- LEFEVRE, D., TEXIER, J.P., RAYNAL, J.P., OCCHIETTI, S. et EVIN, J., 1994 - Enregistrements-réponses des variations climatiques du Pléistocène supérieur et de l'Holocène sur le littoral de Casablanca (Maroc). *Quaternaire*, 5, (3-4), 173-180.
- LEFEVRE, D., RAYNAL, J.P., TEXIER, J.P., GERAADS, D., OCCHIETTI, S. et EL GRAOUI, M., 1996 - Littoraux pliocènes et pléistocènes de Casablanca (Maroc). *Colloque INQUA/PICG 367 Lignes de rivages et zones côtières au Quaternaire*, Perpignan, décembre 1996, résumé.
- LEFEVRE, D., 2000 - *Du continent à l'océan. Morphostratigraphie et paléogéographie du Quaternaire du Maroc atlantique. IIe partie : le modèle casablançais*. Thèse d'Habilitation à diriger des recherches, volume 3, p. 100 à 308, 47 fig., 2 tab., Université de Montpellier III.
- MOHIB, A., 2001 - *L'Acheuléen de la grotte des Ours à Sidi Abderrahmane (Casablanca, Maroc) dans son contexte régional (fouilles anciennes et récentes)*. Thèse de l'Institut National des Sciences de l'Archéologie et du Patrimoine, Rabat, 348 p.
- NEUVILLE, R. et RUHLMANN, A., 1941 - *La place du Paléolithique ancien dans le Quaternaire marocain*. Hespéris, n° VIII, Casablanca, 156 p.
- OCCHIETTI, S., RAYNAL, J.P., PICHET, P. et TEXIER, J.P., 1993 - Aminostratigraphie du dernier cycle climatique au Maroc atlantique, de Casablanca à Tanger. *C. R. Acad. Sc. Paris*, t. 317, série II, 1625-1632.
- OCCHIETTI, S. et RAYNAL, J.P., 1996 - La méthode de datation par les acides aminés appliquée à la préhistoire du Maroc. *XIII International Congress of Prehistoric and Protohistoric Sciences*, Forli, Italia, 8-14 septembre 1996, Abstracts 1, 25.
- OCCHIETTI, S., RAYNAL, J.P., PICHET, P. et LEFEVRE, D., 2002 - Aminostratigraphy of Pleistocene and Holocene littoral deposits, Casablanca, Morocco, and correlation with the Mediterranean paleoshorelines, *Quaternaire*, ce volume.
- RAYNAL, J.P. et TEXIER, J.P., 1989 - Découverte d'Acheuléen ancien dans la carrière Thomas I à Casablanca et problème de l'ancienneté de la présence humaine au Maroc. *C. R. Acad. Sci. Paris*, t. 308, série II, 1743-1749.
- RAYNAL, J.P., TEXIER, J.P., GERAADS, D. et SBIHI-ALAOUI, F.Z., 1990 - Un nouveau gisement plio-pléistocène en Afrique du Nord : Ahl Al Oughlam (ancienne carrière Dépérez) à Casablanca (Maroc). *C. R. Acad. Sci. Paris*, t. 310, série II, 315-320.
- RAYNAL, J.P., MAGOGA, L. et SBIHI-ALAOUI, F.Z., 1992 - Quelques caractères des industries du niveau L de la carrière Thomas I à Casablanca, Maroc (fouilles 88-91). *First International Meeting on Technical Systems to Configure Lithic Objects of Scarce Elaboration*, Montblanc (Tarragona), 27-30 avril 1992, résumé.
- RAYNAL, J.P., GERAADS, D., MAGOGA, L., ELHAJRAOUI, A., TEXIER, J.P., LEFEVRE, D. et SBIHI-ALAOUI, F.Z., 1993 - La grotte des Rhinocéros (Carrière Oulad Hamida 1, anciennement Thomas III, Casablanca), nouveau site acheuléen du Maroc atlantique. *C. R. Acad. Sc. Paris*, t. 316, série II, 1477-1483.
- RAYNAL, J.P. et GERAADS, D., 1993 - Problème patrimonial qui se pose à Casablanca au Maroc. In «*L'objet archéologique africain et son devenir*», CNRS Ed., 49-55.
- RAYNAL, J.P., MAGOGA, L., SBIHI-ALAOUI, F.Z. et GERAADS, D., 1995 - The Earliest Occupation of Atlantic Morocco : The Casablanca Evidence. in *The earliest occupation of Europe*, W. Roebroeks & T. van Kolfschoten Ed., University of Leiden, 1996, 255-262.
- RAYNAL, J.P., SBIHI-ALAOUI, F.Z., AMANI, F., BERNOUSSI, R., EL GRAOUI, M., GERAADS, D., HUBLIN, J.J., LEFEVRE, D., MAGOGA, L., MOHIB, A., OCCHIETTI, S., RHODES, E., SEN, S., TEXIER, J.P. et ZOUAK, M., 1996 - Premiers peuplements du Maroc atlantique : l'exemple de Casablanca. *XIII International Congress of Prehistoric and Protohistoric Sciences*, Forli, Italia, 8-14 septembre 1996, Abstracts 1, 126.
- RAYNAL, J.P., 1998 - Les racines oubliées de Casablanca. *Zellige*, n° 8, juin 1998, 35-36.
- RAYNAL, J.P., LEFEVRE, D., GERAADS, D. et EL GRAOUI, M., 1999 - Contribution du site paléontologique de Lissasfa (Casablanca, Maroc) à une nouvelle interprétation du Mio-Pliocène de la Méseta. *Comptes rendus de l'Académie des Sciences Paris*, Sciences de la terre et des planètes, t. 329, 617-622.
- RAYNAL, J.P., SBIHI ALAOUI, F.Z., GERAADS, D., MAGOGA, L. & MOHIB, A., 2001 - The earliest occupation of North-Africa: the moroccan perspective, *Quaternary International*, 75, 65-75.
- RAYNAL, J.P., MAGOGA, L. et SBIHI ALAOUI, F.Z., 2002 - Quelques caractères des industries acheuléennes du niveau L de la carrière thomas I (Casablanca, Maroc) - fouilles 1988-1991. *Bulletin d'Archéologie Marocaine*, t. 19, sous presse.
- RHODES, E., 1990 - *Optical Dating of Quartz from Sediments*. Thèse, Univ. d'Oxford, 153 p.
- RHODES, E., RAYNAL, J.P., GERAADS, D. et SBIHI-ALAOUI, F.Z., 1994 - Premières dates RPE pour l'Acheuléen du Maroc atlantique (Grotte des Rhinocéros, Casablanca). *C. R. Acad. Sc. Paris*, série II, t. 319, 1109-1115.
- RHODES, E.J., SINGARAYER, J., RAYNAL, J.P. & WESTAWAY, K., 2002 - New luminescence dating for the Palaeolithic assemblages and Pleistocene succession of Casablanca, Morocco. A paraître.
- SAHNOUNI, M. & HADJOUIS, D., 1987 - Découverte de nouveaux documents osseux et lithiques à l'Ain Hanech (Sétif, Algérie Orientale). *Bulletin de la Société Préhistorique de France*, Paris, 84, 133-134.
- SBIHI-ALAOUI, F.Z. et MOHIB, A., 1998 - Aperçu sur le patrimoine préhistorique de Casablanca. Nouvelles archéologiques et patrimoniales, novembre 1998, n°3, 4-7.
- TEXIER, J.P., LEFEVRE, D. et RAYNAL, J.P., 1994 - Contribution pour un nouveau cadre stratigraphique des formations littorales quaternaires de la région de Casablanca. *C. R. Acad. Sci. Paris*, série II, t. 318, n°9, 1247-1253.
- TEXIER, J.P., LEFEVRE, D., RAYNAL, J.P. & EL GRAOUI, M., 2002 - Lithostratigraphy of the littoral deposits of the last one million years in the Casablanca region (Morocco), *Quaternaire*, ce volume.