



HAL
open science

Hominid Cave at Thomas Quarry I (Casablanca, Morocco): recent findings and their context.

Jean-Paul Raynal, Fatima-Zohra Sbihi Alaoui, Abderrahim Mohib, Mohssine El Graoui, David Lefevre, Pierre-Jean Texier, Denis Geraads, Jean-Jacques Hublin, Tanya Smith, Paul Tafforeau, et al.

► **To cite this version:**

Jean-Paul Raynal, Fatima-Zohra Sbihi Alaoui, Abderrahim Mohib, Mohssine El Graoui, David Lefevre, et al.. Hominid Cave at Thomas Quarry I (Casablanca, Morocco): recent findings and their context.. The oldest human expansions in Eurasia Favouing and limiting factors, international Congress,, Nov 2008, Paris, France. halshs-00442647

HAL Id: halshs-00442647

<https://shs.hal.science/halshs-00442647v1>

Submitted on 21 Dec 2009

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Centre National de la Recherche Scientifique



Muséum National d'Histoire Naturelle



**Les premières expansions humaines en Eurasie à partir de l'Afrique
Facteurs limitant ou favorisant**

***The oldest human expansions in Eurasia
Favouring and limiting factors***

***Colloque international organisé au Muséum National d'Histoire Naturelle, Paris
Département de Préhistoire
Auditorium de la Grande Galerie et de l'Evolution
26, 27, 28 novembre 2008***

***International Congress organised in the Muséum National d'Histoire Naturelle, Paris
Auditorium de la Grande Galerie
november 26, 27, 28 - 2008***

Abstract

Hominid Cave at Thomas Quarry I (Casablanca, Morocco): recent findings and their context

**Jean-Paul Raynal, Fatima-Zohra Sbihi-Alaoui, Abderrahim Mohib, Mosshine El Graoui, David Lefèvre
Jean-Pierre Texier, Denis Geraads, Jean-Jacques Hublin, Tanya Smith, Paul Tafforeau, Mehdi Zouak
Rainer Grün, Edward J. Rhodes, Stephen Eggins, Camille Daujeard, Paul Fernandes
Rosalia Gallotti, Saïda Hossini, Alain Queffelec**

Hominid Cave at Thomas Quarry I (Casablanca, Morocco): recent findings and their context

Jean-Paul Raynal¹, Fatima-Zohra Sbihi-Alaoui², Abderrahim Mohib², Mosshine El Graoui²
David Lefèvre³, Jean-Pierre Texier¹, Denis Geraads⁴, Jean-Jacques Hublin⁵, Tanya Smith⁶
Paul Tafforeau⁷, Mehdi Zouak², Rainer Grün⁸, Edward J. Rhodes⁹, Stephen Eggins⁸
Camille Daujeard¹⁰, Paul Fernandes¹, Rosalia Gallotti¹¹, Saïda Hossini¹², Alain Queffelec¹.

- 1 - Université Bordeaux 1 Sciences et technologies, UMR 5199 PACEA, IPGQ, Bât. B18, av. des Facultés, F-33405 Talence
- 2 - Direction du Patrimoine, Institut National des Sciences de l'Archéologie et du Patrimoine, Rabat, Maroc
- 3 - Université Paul Valéry Montpellier III, UMR 5140 « Archéologie des sociétés méditerranéennes », route de Mende, F-34199, Montpellier
- 4 - CNRS UPR 2147, 44 rue de l'Amiral Mouchez, F-75014 Paris
- 5 - Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, D-04103 Leipzig
- 6 - Department of Anthropology, Harvard University, 11 Divinity Avenue, Cambridge, MA 02138, USA
- 7 - Laboratoire de Géobiologie, Biochronologie et Paléontologie Humaine, UMR 6046, CNRS, Université de Poitiers, 40 Avenue du Recteur Pineau, F-86022 Poitiers Cedex ; European Synchrotron Radiation Facility, 6 rue Jules Horowitz, BP 220, F38046 Grenoble Cedex
- 8 - Research School of Earth Sciences, The Australian National University, Canberra ACT 0200, Australia
- 9 - Department of Environmental & Geographical Sciences, Manchester Metropolitan University, Chester Street, Manchester, M1 5GD, UK
- 10 - 6 grande rue de Vaise, F-69009 LYON
- 11 - Largo T. B. Barra, 4, 71022 Ascoli Satriano (FG), Italie
- 12 - Université Moulay Ismaïl, Faculté des Sciences, Département de Géologie, BP 11201, Zitoune, Meknès, Maroc

The Thomas Quarry I, first investigated by P. Biberson (1961a and b), was made famous in 1969 with the discovery by Ph. Beriro of a half-mandible, first attributed to *Atlanthropus mauritanicus* by Ennouchi (1969), but more recently considered as a representative of *Homo rhodesiensis* (Hublin, 2001). In 1985, further investigations revealed the presence of a *Lower Acheulian* assemblage in marly limestones at the bottom of the section (Raynal et Texier, 1989), but it was only from 1988 onwards that modern controlled excavations have been undertaken, within the framework of the Franco-Moroccan co-operative project *Casablanca*.

The stratigraphy of the quarry is complex. It represents several major episodes of coastal sedimentation that are dated to the Lower and Middle Pleistocene on the basis of a detailed regional lithostratigraphical and microfaciological study (El Graoui, 1994; Lefèvre, 2000; Lefèvre et Raynal, 2002; Texier *et al.*, 1994, 2002). Moreover, a continental sediments preserved in a cave belonging to a polyphase paleo-shoreline date to the Middle and Upper Pleistocene on lithostratigraphical and biochronological evidence (Geraads, 1980, 2002; Hossini, 2002). In the Hominid Cave (GH), the general stratigraphy shows at the top a red complex with abundant microfauna (stratigraphic unit 1). Below this, a multilayer dripstone floor (stratigraphic unit 2) caps the lower stratigraphic units (4 and 5). Stratigraphic unit 4 contains artefacts, fauna and hominid fossils and rests on collapsed eolianite blocks imbedded in coarse sands which form stratigraphic unit 5; this lowermost intertidal unit fossilize the notch of the polyphase shoreline (Raynal *et al.*, 1995, 2001, 2002).

The lithic assemblage recovered by recent excavations in GH stratigraphic unit 4 is similar to the series collected at the time of the discovery of the first *Homo* fossil in 1969 (Geraads *et al.*, 1980). It is manufactured mainly on various quartzites available close to the site as blocks and pebbles and a few flint nodules collected from beach deposits. It consists of pebble-tools (mainly unifacial choppers with a few removals) and core-tools, semi-cortical flakes obtained by direct and bipolar flaking, along with rare small handaxes/handaxe-like cores, hammerstones and anvils. As do the upper occupations of the nearby Rhinoceros Cave (GDR) in Oulad Hamida 1 quarry, the GH series represents a facies of the *Acheulian*.

Besides a few reptiles (*Testudo* sp.) and birds, the associated mammalian macrofauna is rich. Primates: *Theropithecus oswaldi*, *Homo* sp. Carnivores: *Herpestes ichneumon*, *Hyaena hyaena*, *Felis* cf. *libyca*, *Panthera* cf. *leo*, cf. *Nyctereutes* sp., *Vulpes* sp., *Lycaon* cf. *magnus*, *Mellivora capensis*,

Lutra sp., *Ursus biberoni*, *Monachus* sp. Artiodactyla: *Kolpochoerus* sp., *Phacochoerus africanus*, Bovini, *Gazella* cf. *atlantica*, Hippotragini, Alcelaphini cf. *Parmularius* sp., *Connochaetes taurinus progna* Perissodactyla: *Equus* cf. *mauritanicus* *Ceratotherium mauritanicum* Rodents: *Hystrix cristata*, *Paraethomys tighennifae*, *Praomys darelbeidae*, *Mus* sp., *Gerbillus* cf. *campestris*, *Gerbillus grandis*, *Meriones* cf. *maghrebianus*, *Meriones* cf. *maximus*, *Ellobius* cf. *atlanticus*, *Eliomys* sp. Lagomorpha: Lagomorpha indet. Insectivores: *Crociodura* sp. Chiroptera: Chiroptera indet.

This faunal assemblage clearly belongs to the same faunal set as the one from GDR whose age lies between 435 ± 85 and 737 ± 129 ka (Rhodes *et al.*, 2006), but some elements show that the assemblage from Thomas GH is earlier. Among large mammals, the suid *Kolpochoerus* is represented by scrappy remains but whose identification is almost certain. It is a widespread genus in East African sites, in the Late Pliocene of Ahl al Oughlam, and is also present in Thomas Quarry I level L; based on the dates of Hominid Cave (see below), this site records its last appearance in North Africa. Some rodents are also distinct from those of GDR. *Meriones* is represented by the same two species, but they are smaller, suggesting an earlier age. *Eliomys* is smaller than *E. darelbeidae* from GDR. Lastly, a lower molar of *Ellobius* has a notched anterior lobe of m1, reminiscent of Tighenif, whereas those from GDR are rounded.

In the macrofauna collected in the central part of the cave during excavations prior to 2005, carnivores were abundant (43% of NISP) while antelopes (25% of NISP) and gazelles (20% of NISP) were dominant among herbivores. Carnivores were considered as the first agent for bone accumulation (Bernoussi, 1997). In the faunal spectrum from recent excavations in the south-eastern part of the cave (2005-2008), cf. *Nyctereutes* sp. is dominant among carnivores (34 % of NISP), and gazelles and other antelopes among herbivores (respectively 31.9 % and 17.1 % of NISP). Preliminary taphonomical analysis of the megafauna shows that the carcasses were processed by carnivores and that cut-marks are absent, which raises the question of the human role in the bone accumulations. Moreover, geological studies have demonstrated that this unit has been partly re-deposited by run-off which mixed human artefacts with the bones accumulated by predators.

Four teeth of *Homo* have been recovered in stratigraphic unit 4 between 1994 and 2005: a right upper premolar (Th1 94 OA 23-24), another right upper premolar (Th1 95 SA 26 n° 89), a first left upper incisor (Th1 95 SA 26 n° 90) and a left upper premolar (Th1 2005 PA 24 n° 107). The teeth are large relative to living humans, and show moderate to heavy wear. It was not possible to quantify crown formation or enamel thickness due to the degree of attrition. Phase contrast synchrotron microtomographic imaging revealed enamel microstructure in one premolar.

The first right upper premolar discovered in 1994 has been dated using laser ablation ICP-MS. Dating procedures followed those applied to the Banyoles mandible (Grün *et al.* 2006). Combining the ESR and U-series data for the modelling of the U-uptake (Grün *et al.* 1988), an US/ESR age of $501^{+94} -76$ ka is obtained (with associated p-values of 1.43 ± 0.59 and 0.75 ± 0.45 for enamel and dentine, respectively). Sediment samples were used for OSL dating. The OSL measurement followed the procedures applied in previous studies of the Pleistocene successions of Casablanca (Rhodes *et al.* 2006). The sample above the tooth yielded an age of 420 ± 34 ka and the one below 391 ± 32 ka. These results are in agreement with an earlier OSL study, which bracketed the age between 360 and 470 ka. Nevertheless, lithostratigraphy and biochronology suggest that this is a minimum age.

Hominid Cave at Thomas Quarry I is vast, and large parts of its deposit are still under study. The preliminary dates, as well as biostratigraphy and lithostratigraphy point towards a greater antiquity for the human fossils than was previously estimated. Thomas Quarry I Hominid Cave is one of the very rare African key-sites for studying individuals and techno-economic behaviors related to a crucial period of human evolution during which North African and European populations started to diverge

biologically.

References:

- Biberson P., 1961a - *Le cadre paléogéographique de la Préhistoire du Maroc atlantique*. Publications du Service des Antiquités du Maroc, Rabat, fascicule 16, 235 p.
- Biberson P., 1961b - *Le Paléolithique inférieur du Maroc atlantique*. Publications du Service des Antiquités du Maroc, Rabat, fascicule 17, 544 p.
- Bernoussi R., 1997 - *Contribution à l'étude paléontologique et observations archéozoologiques pour deux sites du Pléistocène moyen du Maroc atlantique : grotte à Hominidés de la carrière Thomas I et grotte des Rhinocéros de la carrière Oulad Hamida 1 (Casablanca, Maroc)*. Thèse de l'Université de Bordeaux 1, n° 1711, 263 p.
- El Graoui M., 1994 - *Contribution à l'étude des formations littorales Quaternaires de la région de Casablanca (Maroc) : sédimentologie, microfaciès et minéraux lourds*. Thèse de l'Université de Bordeaux 1, n° 1100, 263 p.
- Ennouchi E., 1969 - Découverte d'un Pithécanthropien au Maroc. *C. R. Acad. Sci. Paris*, series D, p. 763.
- Geraads, D., 1980 - La faune des sites à *Homo erectus* des carrières Thomas (Casablanca, Maroc). *Quaternaria*, Roma, 22, 65-94.
- Geraads D., 2002 – Quaternary Mammalian Biostratigraphy of Atlantic Morocco. *Quaternaire*, volume 13, n° 1, 43-54.
- Geraads, D., Beriro, P. & Roche, H. 1980 - La faune et l'industrie des sites à *Homo erectus* des carrières Thomas (Casablanca, Maroc). Précisions sur l'âge de ces Hominidés. *C.R. Acad. Sc. Paris*, D, 291, 195-198.
- Hossini S., 2002 - *La faune d'Anoures marocains du Miocène au Pléistocène et ses rapports avec celle de la même époque au sud-ouest européen : hypothèse sur l'origine des Anoures au Maroc*. Thèse d'Etat. Université Moulay Ismail, 243p
- Grün R., Schwarcz H.P. and Chadam J.M., 1988 - ESR dating of tooth enamel: coupled correction for U-uptake and U-series disequilibrium, *Nucl. Tracks Radiat. Meas.* 14, pp. 237–241.
- Grün R., Maroto J., Eggins S., Stringer C., Robertson S., Taylor L., Mortimer G. and McCulloch M., 2006 - ESR and U-series analyses of enamel and dentine fragments of the Banyoles mandible, *J. Hum. Evol.* 50, pp. 347–358.
- Hublin, J.-J. 2001 - Northwestern African Middle Pleistocene hominids and their bearing on the emergence of *Homo sapiens*. In: L. Barham and K. Robson-Brown (eds.), *Human Roots. Africa and Asia in the Middle Pleistocene*. CHERUB, Western Academic and Specialist Press Lmted, Bristol, pp. 99-121.
- Lefèvre D., 2000 - *Du continent à l'océan. Morphostratigraphie et paléogéographie du Quaternaire du Maroc atlantique. Le modèle casablançais*. Mémoire d'Habilitation à diriger des recherches, volume 3, 11ème partie, p 99- 299, 45 fig., 2 pl., Université de Montpellier III.
- Lefèvre D. et Raynal J.P., 2002 - Les formations plio-pléistocènes de Casablanca et la chronostratigraphie du Quaternaire marin du Maroc revisitées, in *Paléorivages de Casablanca. Stratigraphie et Préhistoire ancienne au Maroc atlantique*, *Quaternaire*, 13, 1, 9-21.
- Raynal J.P. et Texier J.P., 1989 - Découverte d'Acheuléen ancien dans la carrière Thomas I à Casablanca et problème de l'ancienneté de la présence humaine au Maroc. *C. R. Acad. Sci. Paris*, t. 308, series II, 1743-1749.
- Raynal J.P., Magoga L., Sbihi-Alaoui F.Z., Geraads D., 1995 - The Earliest Occupation of Atlantic Morocco: The Casablanca Evidence. in *The earliest occupation of Europe*, W. Roebroeks & T. van Kolfschoten Ed., University of Leiden, 1996, 255-262.
- Raynal J.P., Sbihi Alaoui F.Z., Geraads D., Magoga L., Mohib A., 2001 - The earliest occupation of North-Africa: the moroccan perspective, *Quaternary International*, 75, 65-75.
- Raynal J.P., Sbihi-Alaoui F.Z., Magoga L., Mohib A., Zouak M., 2002 - Casablanca and the early occupation of north-atlantic Morocco. *Quaternaire* 2002, volume 13, 1, 65-77.
- Rhodes, E.J., Singarayer, J.S., Raynal, J.P., Westaway, K.E., Sbihi-Alaoui, F.Z. (2006). New age estimations for the Palaeolithic assemblages and Pleistocene succession of Casablanca, Morocco. *Quaternary Science Reviews* 25, 25698-2585.
- Texier J.P., Lefevre D., Raynal J.P. (1994) - Contribution pour un nouveau cadre stratigraphique des formations littorales Quaternaires de la région de Casablanca (Maroc). *C. R. Acad. Sci. Paris*, t. 318, series II, 1247-1253.
- Texier J.P., Lefevre D., Raynal J.P., El Graoui M., 2002 - Lithostratigraphy of the littoral deposits of the last one million years in the Casablanca region (Maroc). *Quaternaire*, 2002, volume 13, 1, 23-41.