



HAL
open science

LOCALLY CONFORMALLY SYMPLECTIC STRUCTURES ON COMPACT NON- AHLER COMPLEX SURFACES

Vestislav Apostolov, Georges Dloussky

► **To cite this version:**

Vestislav Apostolov, Georges Dloussky. LOCALLY CONFORMALLY SYMPLECTIC STRUCTURES ON COMPACT NON- AHLER COMPLEX SURFACES. 2015. halshs-01108138v1

HAL Id: halshs-01108138

<https://shs.hal.science/halshs-01108138v1>

Preprint submitted on 23 Jan 2015 (v1), last revised 10 Nov 2015 (v2)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LOCALLY CONFORMALLY SYMPLECTIC STRUCTURES ON COMPACT NON-KÄHLER COMPLEX SURFACES

VESTISLAV APOSTOLOV AND GEORGES DLOUSSKY

ABSTRACT. We prove that every compact complex surface with odd first Betti number admits a locally conformally symplectic 2-form which tames the underlying almost complex structure.

1. INTRODUCTION

It is a well-known result [43, 47, 11, 30] that a compact complex surface $S = (M, J)$ admits a Kähler metric if and only if its first Betti number $b_1(M)$ is even. A cornerstone for the proof of this result is the fact, proved independently in [18, Lemme II.3] and [22, p. 185] (see also [43, p. 143, Prop. 1.6] for the case of a K3 surface), that $b_1(M)$ is even if and only if M admits a symplectic form ω which *tames* J , in the sense that the $(1, 1)$ part of ω is positive definite. This and the methods of proof in [11, 30] inspired the so-called “*tamed to compatible*” conjecture in symplectic geometry, which asks whether an almost complex structure on M which is tamed by a symplectic form admits a compatible symplectic form, see [15, 44].

A natural extension of the theory of Kähler manifolds to the non-Kählerian complex case can be obtained through the notion of *locally conformally Kähler* metrics, introduced and studied in foundational work by F. Tricerri and I. Vaisman, see e.g. [14, 36] for an overview. Recall that a locally conformally Kähler (or *lcK*) metric on a complex manifold $X = (M, J)$ is defined by a positive-definite $(1, 1)$ -form F satisfying $dF = \theta \wedge F$ for a closed 1-form θ . The 1-form θ is uniquely determined and is referred to as *the Lee form* of F . The corresponding Hermitian metric $g(\cdot, \cdot) = F(\cdot, J\cdot)$ defines a conformal class c on M . Changing the Hermitian metric $\tilde{g} = e^f g$ within c amounts to transform the Lee form by $\tilde{\theta} = \theta + df$, so that the de Rham class $[\theta]$ is an invariant of the conformal class c .

Of particular interest is the case of compact complex surfaces, where recent works [6, 9, 10, 19] showed that lcK metric exists for all known examples of compact complex surfaces with odd first Betti number, with the exception of the complex surfaces obtained by blowing up points of certain Inoue surfaces with zero second Betti number, described in [6]. However, a general existence result is still to come.

Date: January 13, 2015.

V.A. was supported in part by an NSERC discovery grant and is grateful to the Université Aix-Marseille and the Institute of Mathematics and Informatics of the Bulgarian Academy of Sciences where a part of this project was realized. G.D. is grateful to the UQAM for the hospitality during the preparation of the work. V.A. and G. D. were supported in part by the ANR grant MNGNK-ANR-10-BLAN-0118. The authors are very grateful to Dimiter Vassilev for his invaluable help with the spectral theory of strongly elliptic operators, and to Dimitry Jakobson, François Hamel and Frédéric Rochon for sharing with us their expertise on spectral analysis and the theory of elliptic PDE's. The authors are also grateful to Liviu Ornea and Misha Verbitsky to clarifying their results and giving an access to the unpublished work [40].

In this paper we study, on a compact complex surface $S = (M, J)$ with odd first Betti number, the problem of existence of *locally conformally symplectic forms* ω which tame J , i.e. 2-forms ω satisfying $d\omega = \theta \wedge \omega$ for a closed 1-form θ (called Lee form of ω), and such that the $(1, 1)$ -part of ω is positive-definite. This is, in general, a weaker condition than the existence of lcK metrics, which turns out to be related to the theory of *bihermitian* conformal structures [2, 41] in the case when the ω -conjugate of J determines another integrable almost-complex structure J^ω on M , see [1].

We establish the following general existence result, which we believe is an important step towards the resolution of the existence problems for both lcK and bihermitian conformal structures on a non-Kähler complex surface, and which answers in positive (in the case of complex surfaces) a question raised in [36, Open Problem 1].

Theorem 1.1. *Any compact complex surface $S = (M, J)$ with odd first Betti number admits a locally conformally symplectic form ω which tames J .*

The above theorem is derived from another existence result concerning a conformal class of Hermitian metrics on $S = (M, J)$, which can be regarded as a twisted version of Gauduchon’s celebrated theorem [17], and can be stated as follows. Let $a \in H_{dR}^1(M)$ be a de Rham cohomology class and $\alpha \in a$ a closed 1-form in a . Denote by $d_\alpha := d - \alpha \wedge \cdot$ the twisted differential operator defining the *Lichnerowicz–Novikov* complex, and let $d_\alpha^c := JdJ^{-1}$.

Theorem 1.2. *Let $S = (M, J)$ be a compact complex surface with odd first Betti number, and c a conformal class of Hermitian metrics on S . Then, there exists a non-zero de Rham class $a \in H_{dR}^1(M)$ such that for any metric $g \in c$, there exists a representative $\alpha \in a$ such that the fundamental 2-form F of g satisfies*

$$d_\alpha d_\alpha^c F = 0.$$

The de Rham class a appearing in Theorem 1.2 determines, via the exponential map

$$(1) \quad H_{dR}^1(S, \mathbb{C}) \xrightarrow{\exp} H^1(S, \mathbb{C}^*) \longrightarrow \text{Pic}_0(S),$$

a flat holomorphic line bundle \mathcal{L}_a , and we derive Theorem 1.1 from Theorem 1.2 by showing that, when $H^2(S, \mathcal{L}_a) = \{0\}$, S also admits a locally conformally symplectic form with a Lee form α . The proofs of Theorems 1.1 and Theorem 1.2 are presented in Section 2 of the paper, whereas the necessary analytical tools are collected in the Appendix A.

In the light of the “tamed to compatible” conjecture mentioned above, it is natural to compare the existence on S of lcK metrics and of locally conformally symplectic forms taming J , through the corresponding de Rham classes of their Lee forms. We thus introduce in Section 3 the subset $\mathcal{C}(S)$ (resp. $\mathcal{T}(S)$) in $H_{dR}^1(M)$ of classes a for which there exists a lcK metric on S with Lee form $\theta \in a$ (resp. for which there exists a locally conformally symplectic form which tames J , with Lee form in a). We obviously have the inclusion $\mathcal{C}(S) \subseteq \mathcal{T}(S)$, and one may ask (see also [36], [9, Rem. 9]):

Problem 1.3. Let $S = (M, J)$ be a compact complex surface with odd first Betti number. Determine the set $\mathcal{T}(S) \subset H_{dR}^1(M)$ of classes a for which there exists a locally conformally symplectic form ω which tames J and has a Lee form $\theta \in a$. Is $\mathcal{T}(S)$ strictly bigger than $\mathcal{C}(S)$?

Our initial motivation to study the above problem came from the theory of bihermitian conformal structures developed in [1], where the existence of the latter was

reduced to answering the question of whether certain classes $a \in H_{dR}^1(M)$ belong to $\mathcal{T}(S)$ and $\mathcal{C}(S)$.

A number of partial results concerning Problem 1.3 are obtained in Section 4, where we specialise to the case of a compact complex surface with first Betti number equal to 1.

In the last Section 5 of the paper, we consider some examples of non-Kähler complex surfaces in the Kodaira class VII (i.e. satisfying $H^0(S, \mathcal{K}_S^\ell) = \{0\}$ for all $\ell \geq 1$, where \mathcal{K}_S stands for the canonical line bundle of S , see [5]), for which a complete answer to the above problem can be given. It is known that in this case, the first Betti number equals to 1 (see e.g. [5]), and that the degree with respect to some Gauduchon metric on S of the holomorphic line bundles determined via (1) induces an orientation on $H_{dR}^1(S) \cong (\mathbb{R}, >)$, which turns out to be independent of the choice of a Gauduchon metric (see [46, Rem. 2.4] or Lemma 4.1). Thus, for any compact complex surface in the Kodaira class VII, one can naturally identify $H_{dR}^1(S)$ with the oriented real line $(-\infty, +\infty)$. In this notation, a combination of Propositions 5.1 and 5.2 gives

Theorem 1.4. *Let S be a compact complex surface with a minimal model S_0 .*

- (i) *If S_0 is a Hopf surface, then $\mathcal{T}(S) = \mathcal{C}(S) = (-\infty, 0)$.*
- (ii) *If S is an Inoue surface of the type $S_{N,p,q,r;u}^+$ with $u \in \mathbb{C} \setminus \mathbb{R}$, then $\mathcal{C}(S) = \emptyset$ and $\mathcal{T}(S) = \{a_0\}$.*
- (iii) *If S is an Inoue surface of the type $S_{N,p,q,r;u}^+$ with $u \in \mathbb{R}$, then $\mathcal{C}(S) = \mathcal{T}(S) = \{a_0\}$,*

where $a_0 \in H_{dR}^1(S)$ denotes the de Rham class for which the holomorphic line bundle determined by (1) is isomorphic to the anti-canonical line bundle \mathcal{K}_S^* .

2. EXISTENCE OF LOCALLY CONFORMALLY SYMPLECTIC FORMS TAMING THE COMPLEX STRUCTURE

Let α be a closed 1-form on M , representing de Rham class $a = [\alpha]$. We denote by $L = L_a$ the flat real line bundle over M determined by a via (1), and by $L^* = L_{-a}$ its dual. The corresponding holomorphic line bundles over S will be denoted by \mathcal{L} and \mathcal{L}^* , respectively.

The differential operator $d_\alpha = d - \alpha \wedge \cdot$ defines the Lichnerowicz–Novikov complex

$$(2) \quad \dots \xrightarrow{d_\alpha} \Omega^{k-1}(M) \xrightarrow{d_\alpha} \Omega^k(M) \xrightarrow{d_\alpha} \dots$$

which is isomorphic to the de Rham complex of differential forms with values in L^*

$$(3) \quad \dots \xrightarrow{d_{L^*}} \Omega^{k-1}(M, L^*) \xrightarrow{d_{L^*}} \Omega^k(M, L^*) \xrightarrow{d_{L^*}} \dots$$

In particular, we have an isomorphism between the cohomology groups

$$H_\alpha^k(M) \simeq H_{d_{L^*}}^k(M, L^*).$$

Considering the Dolbeault cohomology groups of S with values in the flat holomorphic line bundle \mathcal{L}^* , we have

$$d_{L^*} = \partial_{\mathcal{L}^*} + \bar{\partial}_{\mathcal{L}^*}, \quad \text{and} \quad d_\alpha = \partial_\alpha + \bar{\partial}_\alpha$$

with

$$\partial_\alpha = \partial - \alpha^{1,0} \wedge \quad \text{and} \quad \bar{\partial}_\alpha = \bar{\partial} - \alpha^{0,1} \wedge,$$

giving rise to the isomorphisms

$$(4) \quad H_{\bar{\partial}_\alpha}^{p,q}(S) \simeq H_{\bar{\partial}_{\mathcal{L}^*}}^{p,q}(S, \mathcal{L}^*).$$

Definition 2.1. Let $X = (M, J)$ be a complex manifold. We shall say that a differentiable 2-form ω is a locally conformally symplectic form taming J if there exists a closed differentiable 1-form α such that $d_\alpha\omega = 0$, and the $(1, 1)$ -part $\omega^{1,1}$ of ω is positive definite. If, furthermore, ω is of type $(1, 1)$, it defines a locally conformally Kähler structure on X . The 1-form α is called the Lee form α of ω .

Remark 2.2. In terms of the isomorphism between (2) and (3), if we write $\alpha|_{U_i} = df_i$ on an open covering $\mathfrak{U} = (U_i)$ of M , then $\omega_i|_{U_i} := e^{-f_i}\omega$ defines a d_{L^*} -closed 2-form with values in L^* , whose $(1, 1)$ -part is positive definite (for the latter we use the fact $L^* = L_{-a}$ is defined by the co-cycle $(e^{f_j - f_i}, U_{ij} = U_i \cap U_j)$ which consist of positive constant real functions). Similarly, if ω is a locally conformally symplectic form taming J with $d_\alpha\omega = 0$, then $\tilde{\omega} = e^f\omega$ is a locally conformally symplectic form taming J which satisfies $d_{\tilde{\alpha}}\tilde{\omega} = 0$ with $\tilde{\alpha} = \alpha + df$. It follows that the existence of a locally conformally symplectic form taming J with Lee form α merely depend upon the de Rham class $a = [\alpha]$.

Our first observation is the following

Lemma 2.3. *Let ω be a locally conformally symplectic form on $X = (M, J)$ with Lee form α . Denote by $F := \omega^{1,1}$ the $(1, 1)$ -part of ω . Then,*

$$d_\alpha d_\alpha^c F = 0,$$

where $d_\alpha^c = Jd_\alpha J^{-1} = i(\bar{\partial}_\alpha - \partial_\alpha)$.

Proof. Writing

$$\omega = F + \omega^{2,0} + \omega^{0,2}$$

where $\omega^{2,0}$ and $\omega^{0,2}$ denote the $(2, 0)$ and $(0, 2)$ -part of ω , respectively. As $d_\alpha = \partial_\alpha + \bar{\partial}_\alpha$, one has

$$d_\alpha\omega = 0 \iff \begin{cases} \partial_\alpha F + \bar{\partial}_\alpha\omega^{2,0} &= 0 \\ \bar{\partial}_\alpha F + \partial_\alpha\omega^{0,2} &= 0, \end{cases}$$

hence $2i\partial_\alpha\bar{\partial}_\alpha F = d_\alpha d_\alpha^c F = 0$. □

Lemma 2.4. *Let $S = (M, J)$ be a complex surface and g a Hermitian metric with fundamental 2-form $F(\cdot, \cdot) = g(J\cdot, \cdot)$. Then,*

$$d_\alpha d_\alpha^c F = 0 \iff \delta(\theta - \alpha) + g(\theta - \alpha, \alpha) = 0,$$

where δ is the co-differential with respect to g and $\theta = J\delta F$ is the Lee form of g .

Proof. Using $dF = \theta \wedge F$ (which, as F is a self-dual 2-form with respect to g is equivalent to the relation $\theta = J\delta F$), one gets

$$d_\alpha d_\alpha^c F = \left[d(J(\theta - \alpha)) + (\theta - \alpha) \wedge J(\theta - \alpha) \right] \wedge F.$$

As F is self-dual, i.e. $*F = F$ where $*$ stands for the Hodge operator with respect to g , $d_\alpha d_\alpha^c F = 0$ is equivalent to

$$\begin{aligned} g\left(d(J(\theta - \alpha)) + (\theta - \alpha) \wedge J(\theta - \alpha), F\right) &= \sum_{i=1}^4 \left(D_{e_i}(J(\theta - \alpha)) \right) (Je_i) + |\theta - \alpha|_g^2 \\ &= -\delta(\theta - \alpha) + g(\delta F, J(\theta - \alpha)) + |\theta - \alpha|_g^2 \\ &= -\delta(\theta - \alpha) - g(\theta, \theta - \alpha) + g(\theta - \alpha, \theta - \alpha) \\ &= -\delta(\theta - \alpha) - g(\theta - \alpha, \alpha) = 0, \end{aligned}$$

where D is the riemannian connection of g and $\{e_i, i = 1, \dots, 4\}$ is any J -adapted orthonormal frame. □

Remark 2.5. In the light of Remark 2.2, it is easily seen that the condition

$$(5) \quad \delta(\theta - \alpha) + g(\theta - \alpha, \alpha) = 0$$

is also conformally invariant. More precisely, it is straightforward to check that if $\tilde{g} = e^f g$ and $\tilde{\alpha} = \alpha + df$, then (5) is satisfied for (g, α) if and only if it is satisfied for $(\tilde{g}, \tilde{\alpha})$.

Recall the fundamental result of Gauduchon [17] which affirms that if $X = (M, J)$ is an m -dimensional compact complex manifold endowed with a Hermitian metric g , then there exists (a unique up to scale) conformal metric $\tilde{g} = e^f g$ whose fundamental form \tilde{F} satisfies $dd^c \tilde{F}^{m-1} = 0$, or equivalently, for which $\tilde{\delta} J \tilde{\delta} \tilde{F} = 0$. Such Hermitian metric is referred to as *Gauduchon metric*. By Lemma 2.4 and Remark 2.5, we then obtain

Proposition 2.6. *Let $S = (M, J)$ be a compact complex surface, $a \in H_{dR}^1(M)$ a de Rham class and $c = [g]$ a conformal class of Hermitian metrics on M , with g a Gauduchon metric in c . Then the following conditions are equivalent:*

- (i) *for any $\tilde{g} \in c$, the fundamental 2-form \tilde{F} satisfies $d_{\tilde{\alpha}} d_{\tilde{\alpha}}^c \tilde{F} = 0$ for some $\tilde{\alpha} \in a$;*
- (ii) *there exists a positive smooth function ψ on M which satisfies the equation*

$$(6) \quad \mathbb{L}_{g,a}(\psi) = \Delta_g(\psi) - g(\theta^g - 2a_h^g, d\psi) + g(\theta^g - a_h^g, a_h^g)\psi = 0,$$

where Δ_g is the Riemannian Laplace operator of g , θ^g is the corresponding co-closed Lee form, and a_h^g is the harmonic representative for a with respect to g .

Proof. By Remark 2.5, if for some metric $\tilde{g} \in c$ the corresponding fundamental form \tilde{F} satisfies $d_{\tilde{\alpha}} d_{\tilde{\alpha}}^c \tilde{F} = 0$, for any other metric $g = e^{-f} \tilde{g}$ in c , the fundamental 2-form F satisfies $d_{\alpha} d_{\alpha}^c F = 0$ with $\alpha = \tilde{\alpha} - df$. It follows that the condition (i) is equivalent to the existence of a 1-form $\alpha \in a$ such that the fundamental 2-form of a Gauduchon metric $g \in c$ satisfies $d_{\alpha} d_{\alpha}^c F = 0$. Writing $\alpha = a_h^g - d \log \psi$ with $\psi > 0$, Lemma 2.4 reads

$$\begin{aligned} 0 &= \delta(\theta^g - \alpha) + g(\theta^g - \alpha, \alpha) = -\delta\alpha + g(\theta^g - \alpha, \alpha) \\ &= -\delta(a_h^g - d \log \psi) + g(\theta^g - a_h^g + d \log \psi, a_h^g - d \log \psi) \\ &= \Delta_g(\log \psi) - g(\theta^g - 2a_h^g, d \log \psi) + g(\theta^g - a_h^g, a_h^g) - g(d \log \psi, d \log \psi) \\ &= \frac{1}{\psi} \left(\Delta_g \psi - g(\theta^g - 2a_h^g, d\psi) + g(\theta^g - a_h^g, a_h^g)\psi \right), \end{aligned}$$

and the claim follows. \square

The general theory for the existence of positive solutions of the elliptic linear second-order PDE $\mathbb{L}_{g,a}(\psi) = 0$ is reviewed in the appendix A to this paper. We recollect below the following variational characterization.

Proposition 2.7. *The PDE (6) has a positive solution ψ if and only if*

$$(7) \quad \lambda_a(g) := \sup_{u \in C^\infty(M), u > 0} \left\{ \min_M \frac{\mathbb{L}_{g,a}(u)}{u} \right\} = 0.$$

Furthermore, $\lambda_a(g)$ is a finite number which varies analytically with respect to linear variations $g_t = (1-t)g + t\tilde{g}$ of Gauduchon metrics, or linear variations $a_t = ta$ of de Rham classes.

Proof of Theorem 1.2. By Propositions 2.6 and 2.7, it is enough to fix a Gauduchon metric $g \in c$ and show that $\lambda_a(g) = 0$ for a suitable choice of $a \neq 0$. Let θ_h^g denote the harmonic part of the Lee form of g . It is well-known (see e.g. [3, Prop. 1] or Proposition 3.5 below) that $[\theta_h^g] \neq 0$ under the assumption that the first Betti number is odd. We let $a_t = \frac{t}{2}[\theta_h^g]$ with $t > 0$ and denote by $a_h^t = \frac{t}{2}\theta_h^g$ the harmonic representative of a_t with respect to g . We are going to show that $\lambda_{a_t}(g) < 0$ when $t \geq 2$ and $\lambda_{a_t}(g) > 0$ for t close to 0; by the second part of Proposition 2.7, this would imply that $\lambda_{a_t}(g) = 0$ for some $t \in (0, 2)$.

Let $u_t > 0$ be the eigenfunction of \mathbb{L}_{g, a_t} , corresponding to the principal eigenvalue $\lambda(t) := \lambda_{a_t}(g)$, normalized by $\int_M u_t^2 v_g = 1$ (see Theorem A.1 in the Appendix A). We then have

$$\begin{aligned}
 \lambda(t) \text{vol}_g(M) &= \int_M \frac{\mathbb{L}_{g, a_t}(u_t)}{u_t} v_g \\
 &= \int_M (\Delta_g u_t) \frac{1}{u_t} v_g - \int_M g(\theta^g - 2a_h^t, d \log u_t) v_g + \int_M g(\theta^g - a_h^t, a_h^t) v_g \\
 (8) \quad &= - \int_M \frac{1}{u_t^2} \|du_t\|_g^2 v_g + \int_M g(\theta_h^g - a_h^t, a_h^t) v_g \\
 &= - \int_M \frac{1}{u_t^2} \|du_t\|_g^2 v_g + \frac{t(2-t)}{4} \int_M \|\theta_h^g\|_g^2 v_g,
 \end{aligned}$$

where we have used that g is a Gauduchon metric (i.e. $\delta\theta^g = 0$) to go from the second to the third line. Taking $t \geq 2$ in (8) yields $\lambda(t) \leq 0$. We are going to show that $\lambda(t) > 0$ for positive t close to 0. Indeed, as for each t $\lambda(t)$ is a simple eigenvalue (see Theorem A.1), by the Kato–Rellich theory [26, 42] (see Theorem A.3 in the Appendix A), $\lambda(t)$ and u_t vary analytically with respect to t . As $\mathbb{L}_{g, 0}$ does not have a zero order term, by the Hopf maximum principle (see e.g. [4, III, Sect. 8, 3.71]) $\lambda(0) = 0$ and $u_0 = 1/\text{vol}_g(M)$. It thus follows that the function $t \mapsto \int_M \frac{\|du_t\|_g^2}{u_t^2} v_g$ has a global minimum 0 at $t = 0$, so differentiating (8) at $t = 0$ we get

$$\lambda'(0) \text{vol}_g(M) = \frac{1}{2} \int_M \|\theta_h^g\|_g^2 v_g > 0.$$

□

In order to obtain a converse of Lemma 2.3, we first notice the following

Lemma 2.8. *Let \mathcal{L} be a flat holomorphic line bundle over a compact complex surface S , such that $H^0(S, \mathcal{L}) = H^2(S, \mathcal{L}) = \{0\}$. Then, $H_{\partial\mathcal{L}^*}^{2,1}(S, \mathcal{L}^*) = \{0\}$.*

Proof. As $H_{\partial\mathcal{L}^*}^{2,1}(S, \mathcal{L}^*) \cong H^1(S, \mathcal{K}_S \otimes \mathcal{L}^*)$, by Serre duality

$$\dim H_{\partial\mathcal{L}^*}^{2,1}(S, \mathcal{L}^*) = h^1(S, \mathcal{K}_S \otimes \mathcal{L}^*) = h^1(\mathcal{L}).$$

Using the vanishing of the cohomology groups from the hypothesis of the Lemma and the Riemann–Roch formula, one concludes $\dim H_{\partial\mathcal{L}^*}^{2,1}(S, \mathcal{L}^*) = 0$. □

Proposition 2.9. *Let g be a Gauduchon Hermitian metric on a compact complex surface S , whose fundamental 2-form F satisfies $d_\alpha d_\alpha^c F = 0$ for some closed but not exact 1-form α . Then the flat holomorphic line bundle \mathcal{L} determined by $a = [\alpha] \in H_{dR}^1(M)$ via (1) satisfies $H^0(S, \mathcal{L}) = \{0\}$. If, moreover, $H^2(S, \mathcal{L}) = \{0\}$, then there exists a locally conformally symplectic 2-form ω on S with Lee form α and whose $(1, 1)$ -part is F .*

Proof. Suppose g is a Gauduchon metric whose fundamental 2-form F satisfies $d_\alpha d_\alpha^c F = 0$. By (6), integrating $\frac{1}{\psi} \mathbb{L}_{g,a}(\psi)$ over the manifold (M, g) , implies

$$(9) \quad \int_M g(\theta^g, a_h^g) v_g > 0.$$

Let \mathcal{L} be the flat holomorphic line bundle corresponding to $a = [\alpha]$. Its degree with respect to the Gauduchon metric g is defined to be

$$\deg_g(\mathcal{L}) = \frac{1}{2\pi} \int_M \rho_{\mathcal{L}} \wedge F,$$

where $\frac{1}{2\pi} \rho_{\mathcal{L}}$ is a pluriharmonic representative of the first Chern class of \mathcal{L} . Writing $\alpha|_{U_i} = df_i$ on an open covering $\mathfrak{U} = (U_i)$ of M , \mathcal{L} is the topologically trivial complex line bundle over M with $s_0 = (U_i, e^{f_i})$ being a nowhere vanishing smooth section, whereas $h|_{U_i} = e^{-2f_i}(\cdot, \cdot)$ introduces a hermitian metric on \mathcal{L} with $h(s_0, s_0) = 1$. It follows that $\rho = -\frac{1}{2} dd^c \log e^{-2f_i} = -d^c \alpha$ is the curvature of the Chern connection of (\mathcal{L}, h) , so that

$$(10) \quad \begin{aligned} \deg_g(\mathcal{L}) &= -\frac{1}{2\pi} \int_M d^c \alpha \wedge F = -\frac{1}{2\pi} \int_M g(d^c \alpha, F) v_g \\ &= -\frac{1}{2\pi} \int_M g(\theta^g, \alpha) v_g = -\frac{1}{2\pi} \int_M g(\theta^g, a_h^g) v_g < 0, \end{aligned}$$

where for the last line we have used that $d\alpha = 0$, $\delta\theta^g = 0$ and (9). By the properties of the degree (see e.g. [18]), it follows that $H^0(S, \mathcal{L}^\ell) = \{0\}$ for all $\ell \geq 1$.

If, moreover, $H^2(S, \mathcal{L}) = \{0\}$, by Lemma 2.8 and the isomorphism (4), we have $H_{\partial_{\mathcal{L}^*}}^{2,1}(S, \mathcal{L}^*) \cong H_{\bar{\partial}_\alpha}^{2,1}(S) = \{0\}$. As $\bar{\partial}_\alpha \partial_\alpha F = \frac{i}{2} d_\alpha d_\alpha^c F = 0$, it follows that

$$\partial_\alpha F + \bar{\partial}_\alpha \beta = 0,$$

for a $(2, 0)$ -form β . Letting

$$\omega = F + \beta + \bar{\beta},$$

one has

$$\begin{aligned} d_\alpha \omega &= (\partial_\alpha + \bar{\partial}_\alpha)(F + \beta + \bar{\beta}) \\ &= \partial_\alpha F + \bar{\partial}_\alpha \beta + \bar{\partial}_\alpha F + \partial_\alpha \bar{\beta} = 0, \end{aligned}$$

which completes the proof. \square

Proof of Theorem 1.1. Let $S = (M, J)$ be a compact complex surface with odd first Betti number $b_1(M)$. If the Kodaira dimension of S is non-negative, it follows from [6] and [48] that S admits a locally conformally Kähler metric. We therefore suppose that the Kodaira dimension of S is negative, i.e. S belongs to class VII of the Kodaira list, see e.g. [5]. Denote by S_0 the minimal model of S .

We first suppose that the second Betti number of S_0 is greater or equal to 1, i.e. S_0 is in the Kodaira class VII_0^+ (for which a complete classification is still to come). In this case, Theorem 1.1 follows from Theorem 1.2, Proposition 2.9 and the following vanishing result.

Lemma 2.10. *Let S be a compact complex surface with $b_1(S) = 1$ and negative Kodaira dimension. Suppose that the second Betti number of its minimal model S_0 is > 0 . Then, for any topologically trivial line bundle $\mathcal{L} \in \text{Pic}_0(S)$, we have*

$$H^2(S, \mathcal{L}) = 0.$$

Proof. By Serre duality, we have to show that for any $\mathcal{L} \in \text{Pic}_0(S)$, $H^0(S, \mathcal{K}_S \otimes \mathcal{L}) = 0$. Suppose that there exists a non trivial section $\sigma \in H^0(S, \mathcal{K}_S \otimes \mathcal{L})$. The line bundle $\mathcal{K}_S \otimes \mathcal{L}$ is not trivial as otherwise \mathcal{K}_S^{-1} must be flat, and therefore $0 = c_1^2(S) = -b_2(S) = -c_2(S)$ (see [5]), a contradiction. It follows that σ must vanish along an effective divisor D with $[D] = \mathcal{K}_S \otimes \mathcal{L}$. Therefore in $H^2(S, \mathbb{Z})$,

$$(11) \quad 0 = (\mathcal{K}_S + \mathcal{L} - D) \cdot \mathcal{K}_S = \mathcal{K}_S^2 - \mathcal{K}_S \cdot D.$$

We show the assertion by induction on the number p of blowing ups.

If $p = 0$, the surface is minimal and by [34, p. 399], an irreducible curve C of D is either

- a rational curve such that $C^2 \leq -2$, therefore by the adjunction formula, $0 = \pi(C) = 1 + \frac{\mathcal{K}_S \cdot C + C^2}{2}$ and $\mathcal{K}_S \cdot C \geq 0$, or
- an elliptic or a rational curve C with a double point. By the same formula, $\mathcal{K}_S \cdot C + C^2 = 0$, hence $\mathcal{K}_S \cdot C \geq 0$,

We deduce that for any effective divisor D , $\mathcal{K}_S \cdot D \geq 0$. Moreover, as $b_2(S) = -c_1(S)^2 = -\mathcal{K}_S^2 > 0$, we have a contradiction in (11).

If $p \geq 1$, there is an exceptional curve of the first kind E and let $B_x : S \rightarrow \check{S}$ be the blowing down of E to a point $x \in \check{S}$. Let \check{U} be a ball centred at x and $U = B_x^{-1}(\check{U})$ a simply connected neighbourhood of E . The line bundle \mathcal{L} is holomorphically trivial on U and, therefore, the coherent sheaf $\check{\mathcal{L}} = (B_x)_* \mathcal{L}$ is in fact a topologically trivial line bundle on \check{S} . As $(B_x)_* \mathcal{K}_S = \mathcal{K}_{\check{S}}$, any non-trivial section $\sigma \in H^0(S, \mathcal{K}_S \otimes \mathcal{L})$ gives, via the biholomorphism $B_x : S \setminus E \cong \check{S} \setminus \{x\}$ a non-trivial section $\check{\sigma} \in H^0(\check{S} \setminus \{x\}, \mathcal{K}_{\check{S}} \otimes \check{\mathcal{L}})$. By Hartogs' theorem, this section extends to \check{S} . As $b_2(\check{S}) \geq b_2(S_0) > 0$, by the induction hypothesis $\check{\sigma} = 0$, a contradiction. \square

Let us now consider the case when $b_2(S_0) = 0$. According to [8, 45, 33], S_0 is either an Inoue surface (see [24]) or a Hopf surface (see [25]). The arguments in [48, 49] can be used without any change to show that the blow-up S of S_0 admits a locally conformally symplectic form taming J if S_0 does. We thus consider the case $S = S_0$.

The Inoue surfaces with second Betti number equal to zero are classified by Inoue [24] who shows that they do not admit curves. Thus, if S is such a surface, the condition $H^2(S, \mathcal{L}) \cong H^0(S, \mathcal{K}_S \otimes \mathcal{L}^*) = \{0\}$ is equivalent to $\mathcal{L} = \mathcal{K}_S$, where \mathcal{K}_S denotes the canonical bundle of S . Let g be a Gauduchon metric whose fundamental 2-form F satisfies $d_\alpha d_\alpha^c F = 0$ for a 1-form α with $a = [\alpha] \neq 0$, given by Theorem 1.2. The corresponding line bundle $\mathcal{L} = \mathcal{L}_a$ has negative degree with respect to g by (10) whereas it is shown in [46, Remark 4.2] that the degree of \mathcal{K}_S is positive with respect to any Gauduchon metric. Thus, $\mathcal{L} \neq \mathcal{K}_S$, showing that $H^2(S, \mathcal{L}) = \{0\}$, and therefore S admits a locally conformally symplectic 2-form taming J with Lee form α by Proposition 2.9.

If S is a Hopf surface, Lemma 2.10 generally fails, so we cannot directly use Theorem 1.2 in this case. However, according to [6, 19], S admits a lcK metric. \square

3. COHOMOLOGICAL INVARIANTS OF A NON-KÄHLER COMPLEX MANIFOLD

An important and well-studied invariant associated to a compact complex manifold $X = (M, J)$ which admits Kähler metrics is its *Kähler cone*, $\mathcal{K}(X)$, defined to be the subset of classes $\Omega \in H_{dR}^{1,1}(X, \mathbb{R})$ for which there exists a Kähler metric on X whose fundamental 2-form belongs to Ω . A characterization of $\mathcal{K}(X)$ in terms of the intersection form of the cohomology ring of X , the Hodge structure and homology of analytic cycles has been obtained by N. Buchdahl [11, 12] and Lamari [30, 31] when X is a compact complex surface, and Demailly–Paun [13] in general. These results

imply in particular that the classes in $H_{dR}^{1,1}(X)$ which contain symplectic forms taming J coincide with $\mathcal{K}(X)$.

In order to introduce similarly designed cohomological invariants in the non-Kähler lcK case, one can consider the sets (see [36, 9]):

Definition 3.1. Let $X = (M, J)$ be a compact complex manifold. We introduce the following subsets of $H_{dR}^1(M)$:

- The subset of classes of Lee forms of lcK metrics:

$$\mathcal{C}(X) = \{[\alpha] \mid \exists F \in \Omega^{1,1}(X), F > 0, d_\alpha F = 0\}.$$

- The subset of classes of Lee forms of locally conformally symplectic forms taming J :

$$\mathcal{T}(X) = \{[\alpha] \mid \exists \omega \in \Omega^2(X), \omega^{1,1} > 0, d_\alpha \omega = 0\}.$$

- The subset of classes of the harmonic parts of the Lee forms of Gauduchon metrics

$$\mathcal{G}(X) = \{[\theta_h^g] \mid \exists F \in \Omega^{1,1}(X), F > 0, dd^c F^{m-1} = 0, d_\theta F^{m-1} = 0\},$$

where θ_h^g denotes the harmonic part of θ with respect to the riemannian metric $g(\cdot, \cdot) = F(\cdot, J\cdot)$.

Remark 3.2. It follows from the definition that

$$\mathcal{C}(X) \subseteq \mathcal{T}(X), \quad \mathcal{C}(X) \subseteq \mathcal{G}(X),$$

$\mathcal{G}(X)$ is connected while $\mathcal{T}(X)$ is invariant under small deformations of X . Using similar techniques as in [20], one can show that $a \in \mathcal{T}(X)$ is an interior point, provided that $H_{d_L^*}^3(M, L^*) = \{0\}$, where $L = L_a$ is the real flat line bundle determined by a .

We now consider the blow-up \hat{X} of X at a point x and denote $B_x \hat{X} \rightarrow X$ the blow-down map which is a biholomorphism between $\hat{X} \setminus E \rightarrow X \setminus \{x\}$, where $E \cong \mathbb{C}P^{m-1}$ is the exceptional divisor. The following is well-known:

Lemma 3.3. *Let $B_x : \hat{X} \rightarrow X$ be the blow-down map which contracts a divisor $E \cong \mathbb{C}P^{m-1} \subset \hat{X}$ with normal bundle $N_E \cong \mathcal{O}(-1)$ to a point $x \in X$. For any $a \in H_{dR}^1(X)$ with generator $\alpha \in a$, denote by $\hat{\alpha} = B_x^*(\alpha)$. Then $B_x^* : H_\alpha^k(X) \rightarrow H_{\hat{\alpha}}^k(\hat{X})$ is surjective for any positive $k \neq 2(m-1)$, and is injective for any positive $k \neq 2m-1$.*

Proof. As $H_\alpha^k(X)$ does not depend on the choice of $\alpha \in a$ (see the discussion in Sec. 2), we can choose α such that it identically vanishes on a open ball U centred at x . It follows that $\hat{\alpha} = B_x^*(\alpha)$ vanishes on $\hat{U} = B_x^{-1}(U)$.

We shall first prove that $B_x^* : H_\alpha^k(X) \rightarrow H_{\hat{\alpha}}^k(\hat{X})$ is surjective. With our choice for α , any $d_{\hat{\alpha}}$ -closed k -form $\hat{\varphi}$ on \hat{X} is closed over \hat{U} . As $H_{dR}^k(\hat{U}) \cong H_{dR}^k(\mathbb{C}P^{m-1}) = \{0\}$ for $k \neq 2(m-1)$, we can write $\hat{\varphi}|_{\hat{U}} = d(\hat{\xi}|_{\hat{U}})$. Multiplying $\hat{\xi}|_{\hat{U}}$ by the pull-back via B_x of a bump function centred at x and support in U , we can assume $\hat{\xi}$ is globally defined on \hat{X} and $\hat{\varphi} = \hat{\varphi} - d_{\hat{\alpha}} \hat{\xi}$ is another form representing $[\hat{\varphi}] \in H_{\hat{\alpha}}^k(\hat{X})$ which vanishes identically on a tubular neighbourhood of E . Then, the diffeomorphism $(B_x^{-1}) : \hat{X} \setminus E \rightarrow X \setminus \{x\}$ allows to define a smooth k -form $\phi = (B_x^{-1})^*(\hat{\varphi})$ on X with $d_\alpha \phi = 0$ and $B_x^*(\phi) = \hat{\varphi}$.

We now prove that $B_x^* : H_\alpha^k(X) \rightarrow H_{\hat{\alpha}}^k(\hat{X})$ is injective. Suppose φ is a d_α -closed k -form on X , such that $\hat{\varphi} = B_x^*(\varphi) = d_{\hat{\alpha}} \hat{\xi}$. As $H_{dR}^k(U) = \{0\}$, we can modify φ with a d_α -exact form (as we did above with $\hat{\varphi}$) and assume without loss that $\varphi|_U \equiv 0$. It follows

that the $(k-1)$ -form $\hat{\xi}$ satisfies $d\hat{\xi}|_{\hat{U}} \equiv 0$. If $k = 1$, $\hat{\xi}$ is a smooth function on \hat{X} which is constant on \hat{U} and, therefore, is the pull back to \hat{X} of a smooth function ξ on X (which is constant on U). It follows that $\varphi = d_\alpha \xi$. If $k > 1$, $H_{dR}^{k-1}(\hat{U}) \cong H^{k-1}(\mathbb{C}P^{m-1}) = \{0\}$, so we conclude that $\hat{\xi}|_{\hat{U}}$ is exact, i.e. $\hat{\xi}|_{\hat{U}} = d\hat{\eta}|_{\hat{U}}$. Multiplying $\hat{\eta}$ by a bump-function, we obtain that $\hat{\xi} - d_\alpha \hat{\eta}$ is identically zero in a neighbourhood of E , so that it descends to X to define a $(k-1)$ -form ξ with $d_\alpha \xi = \varphi$. \square

We recall the following result established in [48, 49]. The argument in [49] applies without change to the case of locally conformally symplectic structures taming J :

Proposition 3.4. *Let \hat{X} be the blow-up of X at a point x . Then*

- (a) *if $a \in \mathcal{T}(X)$, then $\hat{a} = B_x^*(a) \in \mathcal{T}(\hat{X})$;*
- (b) *$a \in \mathcal{C}(X)$ if and only if $\hat{a} = B_x^*(a) \in \mathcal{C}(\hat{X})$.*

When $X = S$ is a compact complex surface, the following result is well-known:

Proposition 3.5. *On a compact complex surface $S = (M, J)$ the following conditions are equivalent:*

- (i) $0 \in \mathcal{C}(S)$, i.e. S is Kähler;
- (ii) $0 \in \mathcal{T}(S)$;
- (iii) $0 \in \mathcal{G}(S)$;
- (iv) $\mathcal{C}(S) = \mathcal{T}(S) = \mathcal{G}(S) = \{0\}$.

Proof. ‘(i) \Rightarrow (ii)’ and ‘(iv) \Rightarrow (i)’ are obvious. In order to prove ‘(ii) \Rightarrow (iii)’, let ω be a symplectic 2-form on M which tames J . Letting $F = \omega^{1,1}$ be the positive definite $(1, 1)$ -part, it defines a hermitian metric $g(\cdot, \cdot) = F(\cdot, J\cdot)$ which is Gauduchon (see e.g. Lemma 2.3 with $\alpha = 0$). Furthermore, ω is a closed self-dual 2-form with respect to g , and is therefore co-closed. Writing $\omega = F + \Psi$, where Ψ is of type $(2, 0) + (0, 2)$, we get

$$0 = \delta\omega = \delta F + \delta\Psi = -J\theta^g - J(\delta\mathcal{J}\Psi),$$

where \mathcal{J} stands for the natural action of the almost-complex structure J on the bundle of real 2-forms of type $(2, 0) + (0, 2)$ by $\mathcal{J}\Psi(X, Y) := -\Psi(JX, Y)$. This shows that the Lee form θ of g is co-exact and, therefore, $\theta_h^g = 0$.

It remains to establish ‘(iii) \Rightarrow (iv)’. To this end, by [3, Prop. 1], $b_1(M)$ is even and $\mathcal{G}(S) = \{0\}$. It follows that $\mathcal{C}(S) = \{0\}$ as $\mathcal{C}(S) \subseteq \mathcal{G}(S)$ and $\mathcal{C}(S) \neq \emptyset$ by the characterization of Kähler surfaces [43, 47, 11, 30].

In order to prove $\mathcal{T}(S) = \{0\}$, we use Proposition 2.6: integrating $\frac{1}{\psi} \mathbb{L}_{g,a}(\psi) = 0$ over (M, g) implies (by using that the harmonic part θ_h^g of the Lee form vanishes)

$$0 \leq \int_M g(\theta^g - a_h^g, a_h^g) v_g = \int_M g(\theta_h^g - a_h^g, a_h^g) v_g = - \int_M \|a_h^g\|_g^2 v_g,$$

showing $a_h^g = 0$. \square

Recall the following definition from [37]

Definition 3.6. A *lcK metric with potential g* on $X = (M, J)$ is a lcK metric such that the pull-back \tilde{F} of its fundamental 2-form F to the universal covering space \tilde{X} of X is of the form $\tilde{F} = \frac{dd^c \tilde{f}}{\tilde{f}}$, where $\tilde{f} > 0$ is a positive plurisubharmonic function on \tilde{X} which satisfies $\gamma^* \tilde{f} = e^{c_\gamma} \tilde{f}$ ($c_\gamma \in \mathbb{R}$) for any deck-transform $\gamma \in \pi_1(X)$.

Examples of lcK metrics with potential include the *Vaisman* lcK metrics (i.e. lcK metrics for which the Lee form θ is parallel) or more generally, *pluricanonical* lcK metrics, introduced and studied by G. Kokarev in [29], for which the covariant derivative $D\theta$ of the Lee form is of type $(2, 0) + (0, 2)$ with respect to J , see [37, Claim 3.3] and [40].¹ The following observation is taken from [36].

Lemma 3.7. *Let $X = (M, J)$ be a compact complex manifold endowed with a lcK metric with potential, g , and Lee form θ . Then for any $t \geq 1$, $t\theta$ is the Lee form of a lcK metric with potential on X . If, furthermore, g is a pluricanonical lcK metric, then $t\theta$ is the Lee form of a pluricanonical lcK metric for any $t > 0$.*

Proof. Writing $\tilde{F} = \frac{dd^c \tilde{f}}{\tilde{f}}$ on \tilde{X} , the pull-back of the Lee form is $\tilde{\theta} = -\frac{d\tilde{f}}{\tilde{f}}$. For any $t \geq 1$ put

$$\tilde{F}_t := \frac{dd^c \tilde{f}^t}{\tilde{f}^t} = t \frac{dd^c \tilde{f}}{\tilde{f}} + t(t-1) \frac{d\tilde{f} \wedge d^c \tilde{f}}{\tilde{f}^2}.$$

For $t \geq 1$, \tilde{F}_t defines a positive definite $(1, 1)$ -form satisfying $d\tilde{F}_t = \tilde{\theta}_t \wedge \tilde{F}_t$ with $\tilde{\theta}_t = -t \frac{d\tilde{f}}{\tilde{f}} = t\tilde{\theta}$. As \tilde{F}_t is invariant under any deck transformation, it defines a lcK metric with potential, g_t , on X , whose Lee form is $\theta_t = t\theta$.

It is not hard to see that the pluricanonical condition $(D_X^g \theta)(Y) = -(D_{JX}^g \theta)(JY)$ is equivalent to

$$(12) \quad dJ\theta = -|\theta|_g^2 F + \theta \wedge J\theta.$$

This is essentially the formula appearing in [37, p. 724], by noting that there is a sign error in [37] in deriving the formula for $d(I\theta)$ from the previous one, and an omission of a factor $|\theta|_g^2$ before g in the formula expressing $\nabla\theta - D\theta$; the precise statement appears in [40]. For convenience of the reader we supply here a brief argument for (12): it is well-known (see e.g. [27, II, Prop. 4.2]) that when J is integrable

$$(D_X F)(Y, Z) = -\frac{1}{2} \left(dF(X, JY, JZ) - dF(X, Y, Z) \right).$$

The lcK condition implies $dF = \theta \wedge F$, which allows to re-write the above equality as

$$(13) \quad D_X F = \frac{1}{2} \left(X^\flat \wedge J\theta + JX^\flat \wedge \theta \right),$$

where X^\flat denotes the g -dual 1-form to X . Thus, using (13),

$$\begin{aligned} (D_X J\theta)(Y) &= -\theta((D_X J)(Y)) - (D_X \theta)(JY) \\ &= \frac{1}{2} \left(\theta(X)J\theta(Y) - J\theta(X)\theta(Y) - |\theta|_g^2 F(X, Y) \right) - (D_X \theta)(JY). \end{aligned}$$

It then follows

$$dJ\theta = -|\theta|_g^2 F + \theta \wedge J\theta + (JD\theta)^{\text{anti}},$$

where $(JD\theta)^{\text{anti}}$ denotes the anti-symmetrization of the $(2, 0)$ -tensor $(JD\theta)(X, Y) := -D\theta(X, JY)$; as $d\theta = 0$ (i.e. $D\theta$ is symmetric), $JD\theta^{\text{anti}}(X, Y) = -(D\theta)^{1,1}(X, JY)$ where $(D\theta)^{1,1}(X, Y) := (D_X \theta)(Y) + (D_{JX} \theta)(JY)$. Thus, for a lcK metric g , the equation (12) is equivalent to $(D\theta)^{1,1} = 0$, i.e. to g being pluricanonical.

Differentiating (12) one more time yields

$$d|\theta|_g^2 \wedge F = 0,$$

¹In [37], the authors claim that a lcK metric admits a potential if and only if it is pluricanonical, but a proof is given only in one direction; see [40] for the precise link between the two notions.

showing that $|\theta|_g^2$ is a constant. Thus

$$F_t := F + \frac{t}{|\theta|_g^2} \theta \wedge J\theta, \quad t > -1$$

defines a family of positive definite $(1,1)$ -forms with $dF_t = (1+t)\theta \wedge F_t$. Clearly, F_t give rise to a family of lcK metrics which verify the pluricanonical condition (12). \square

Remark 3.8. If g is a pluricanonical lcK (non-Kähler) metric on $X = (M, J)$ with Lee form θ , writing the pull-back metric as $\tilde{g} = e^{\tilde{\varphi}} \tilde{g}_K$ where \tilde{g}_K is a Kähler metric on the universal cover \tilde{X} conformal to \tilde{g} (or, equivalently, writing the pull-back $\tilde{\theta}$ of the Lee form as $\tilde{\theta} = d\tilde{\varphi}$) gives rise to a potential function $\tilde{f} := e^{-\tilde{\varphi}}$ for the fundamental 2-form \tilde{F} of \tilde{g} , i.e. $\tilde{F} = \frac{d\tilde{f}}{\tilde{f}}$ by (12).

4. COMPACT COMPLEX SURFACES WITH $b_1(S) = 1$

We start with the following easy consequence of Proposition 3.5.

Lemma 4.1. *Let $S = (M, J)$ be a compact complex surface with $b_1(M) = 1$. For any $a \in H_{dR}^1(M)$, denote by \mathcal{L}_a the corresponding flat holomorphic line bundle defined via (1). Then, the sign of $\deg_g(\mathcal{L})$ does not depend on the choice of a Gauduchon metric on S and is zero if and only if $a = 0$.*

Proof. Let g be a Gauduchon metric on S . The computation in (10) shows

$$\deg_g(\mathcal{L}_a) = -\frac{1}{2\pi} \int_M g(\theta^g, a_h^g) v_g = -\frac{1}{2\pi} \int_M g(\theta_h^g, a_h^g) v_g.$$

It follows that $\deg_g(\mathcal{L}_a) = 0$ for any Gauduchon metric if $a = 0$. On the other hand, if $a \neq 0$, $\theta_h^g = \mu_a(g) a_h^g$ for a constant $\mu_a(g)$, so that the sign of $\deg_g(\mathcal{L}_a)$ is equal to the sign of $-\mu_a(g)$. We know by Proposition 3.5 that $\mu_a(g) \neq 0$. As the space of Gauduchon metrics is convex (and therefore connected), it follows that the sign of $\mu_a(g)$ is non-zero and is independent of g . \square

The above Lemma allows for the following

Definition 4.2. Let $S = (M, J)$ be a compact complex surface with $b_1(M) = 1$. For $a, b \in H_{dR}^1(M)$ we will say that $a > b$ if

$$\deg_g(\mathcal{L}_a \otimes \mathcal{L}_b^*) = \deg_g(\mathcal{L}_{(a-b)}) = \deg_g(\mathcal{L}_a) - \deg_g(\mathcal{L}_b) > 0,$$

for some (and hence any) Gauduchon metric g .

We will thus identify the ordered set $H_{dR}^1(S) \cong (\mathbb{R}, >)$.

Proposition 4.3. *Let $S = (M, J)$ be a compact complex surface with $b_1(M) = 1$. Then $\mathcal{G}(S) \subseteq (-\infty, 0)$ and $\mathcal{T}(S) \subseteq (-\infty, 0)$. In particular, for any $a \in \mathcal{T}(S)$, $H^0(S, \mathcal{L}_a) = \{0\}$. Furthermore, for each $a \in \mathcal{T}(S)$, there exists a class $b \leq a$ which belongs to $\mathcal{G}(S)$.*

Proof. Let g be Gauduchon metric on S and θ_h^g the harmonic part of the corresponding Lee form θ^g . Applying (10) for $a = [\theta_h^g]$ yields

$$\deg_g(\mathcal{L}_a) = -\frac{1}{2\pi} \int_M g(\theta^g, \theta_h^g) v_g = -\frac{1}{2\pi} \int_M \|\theta_h^g\|_g^2 v_g < 0,$$

according to Proposition 3.5, thus showing the first inclusion.

For the second one, one integrates the identity $\frac{1}{\psi}\mathbb{L}_{g,a}(\psi) = 0$ in Proposition 2.6(ii) in order to obtain

$$(14) \quad 0 \leq \int_M g(\theta^g - a_h^g, a_h^g)v_g = \int_M g(\theta^g, a_h^g)v_g - \int_M \|a_h^g\|_g^2 v_g.$$

It follows that

$$\deg_g(\mathcal{L}_a) = -\frac{1}{2\pi} \int_M g(\theta^g, a_h^g) \leq -\frac{1}{2\pi} \int_M \|a_h^g\|_g^2 v_g < 0,$$

using again Proposition 3.5. The vanishing $H^0(S, \mathcal{L}_a) = \{0\}$ follows from the properties of the degree.

Inequality (14) and the fact that a and $b = [\theta_h^g]$ are both negative (so that $\theta_h^g = \mu a_h^g$ with $\mu > 0$) show $\deg_g(\mathcal{L}_{a-b}) \geq 0$, i.e. the de Rham class $b \leq a$. \square

Proposition 4.4. *Suppose $S = (M, J)$ is a compact complex surface with $b_1(M) = 1$. If $b \in \mathcal{C}(S)$, $c \in \mathcal{G}(S)$ with $b \leq c$, then any $a \in [b, c] \subset H_{dR}^1(S)$, such that the corresponding flat line bundle \mathcal{L}_a satisfies $H^2(S, \mathcal{L}_a) = \{0\}$, belongs to $\mathcal{T}(S)$.*

Proof. Let g be a lcK metric on S whose closed Lee form θ^g belongs to b , and suppose (without loss) that g is a Gauduchon metric: thus, the Lee form of g is harmonic, i.e. $\theta^g = b_h^g$. For any $a \in H_{dR}^1(M)$, let a_h^g denote the harmonic representative of a with respect to g : thus, writing $a_h^g = \mu b_h^g$ for a constant μ , we have $a \geq b$ iff

$$\deg_g(\mathcal{L}_{b-a}) = -\frac{1}{2\pi} \int_M g(b_h^g, b_h^g - a_h^g)v_g = \frac{(\mu - 1)}{2\pi} \int_M \|b_h^g\|_g^2 v_g \leq 0,$$

i.e. iff $\mu \in [0, 1]$. It then follows that for any $a \in [b, c]$,

$$(15) \quad \mathbb{L}_{g,a}(\psi) = \Delta_g \psi + (2\mu - 1)g(b_h^g, d\psi) + (1 - \mu)\mu \|b_h^g\|_g^2 \psi.$$

We claim that $\lambda_a(g) \geq 0$. Indeed,

$$\lambda_a(g) \geq \min_M \mathbb{L}_{g,a}(1) = (1 - \mu)\mu \min_M \|b_h^g\|_g^2 \geq 0.$$

Similarly, let \tilde{g} be a Gauduchon metric on S for which the de Rham class $c = [\tilde{\theta}_h^g]$ of the harmonic part of its the Lee form satisfies $c \geq b$. For any $a \leq c$, we have $\tilde{\theta}_h^g = \nu a_h^{\tilde{g}}$ for a real constant $\nu \in [0, 1]$. It then follows that

$$\int_M \tilde{g}(\tilde{\theta}_h^g - a_h^{\tilde{g}}, a_h^{\tilde{g}})v_{\tilde{g}} = (\nu - 1) \int_M \|a_h^{\tilde{g}}\|_{\tilde{g}}^2 v_{\tilde{g}} \leq 0.$$

As we saw in the proof of Theorem 1.2, this implies $\lambda_a(\tilde{g}) \leq 0$.

Considering the linear path $g_t = (1 - t)g + t\tilde{g}$, $t \in [0, 1]$ of Gauduchon metrics and using the continuity of $\lambda_a(g_t)$ (see Proposition 2.7), one concludes that there exists a Gauduchon metric g' on S with $\lambda_a(g') = 0$. Our claim then follows from Proposition 2.9. \square

5. EXAMPLES

In this section we illustrate the previous discussion on the known compact complex surfaces S in the Kodaira class VII [5]. Note that in this case $b_1(S) = 1$.

5.1. Hopf surfaces. These are, by definition, compact complex surfaces with universal covering space $\mathbb{C}^2 \setminus \{(0,0)\}$. It is shown by Kodaira [28] that the fundamental group Γ of such a surface is a finite extension of the infinite cyclic group \mathbb{Z} . The list of concrete realizations of Γ as a group of automorphisms of \mathbb{C}^2 can be found in [25], and we summarize this classification in the following rough form: $\Gamma = H \ltimes \langle \gamma_0 \rangle$, where $\langle \gamma_0 \rangle$ denotes the infinite cyclic group generated by the contraction γ_0 of \mathbb{C}^2

$$\gamma_0(z_1, z_2) = (\alpha z_1 + \lambda z_2^m, \beta z_2),$$

where the complex numbers α, β, λ satisfy $0 < |\alpha| \leq |\beta| < 1$, $\lambda(\alpha - \beta^m) = 0$ for an integer $m \in \mathbb{N}^*$. Furthermore, it follows by the classification in [25] that when $\lambda \neq 0$, $H \subset U(1) \times U(1)$ is abelian and commutes with γ_0 . We denote by $S_{\alpha, \beta, \lambda} = \mathbb{C}^2 \setminus \{(0,0)\} / \langle \gamma_0 \rangle$ the corresponding *primary* Hopf surface and by $S_{\alpha, \beta, \lambda; H}$ the further quotient by H , called *secondary* Hopf surface.

Hopf surfaces with $\lambda = 0$ are called *diagonal* (or, confusingly, of class 1). Belgun shown [6, Thm. 1] that any such surface admits a Vaisman lcK metric. By Lemma 3.7 and Proposition 4.3, in this case we have $\mathcal{T}(S) = \mathcal{C}(S) = (-\infty, 0)$.

Hopf surfaces with $\lambda \neq 0$ are called *resonant* (also called, even more confusingly, of class 0). Let $S_{\beta^m, \beta, \lambda; H}$ be a resonant Hopf surface. The analytic family $S_\lambda := S_{\beta^m, \beta, \lambda; H}$, $\lambda \in \mathbb{C}$ has as central fibre the diagonal Hopf surface $S_0 = S_{\beta^m, \beta, 0; H}$, while for $\lambda \neq 0$ the surfaces S_λ are isomorphic (see [19]). As S_0 admits a taming conformally symplectic form with Lee form in any $a \in (-\infty, 0)$ so does S_λ , just by continuity using that the taming condition is open. It follows that $\mathcal{T}(S_\lambda) = (-\infty, 0)$.

Similarly, as S_0 admits a Vaisman metric by [6, 19], it has a Vaisman lcK metric g_a with fundamental 2-form F_a and Lee form in a for any $a \in (-\infty, 0)$, by Lemma 3.7. (Recall that any Vaisman lcK metric is pluriccanonical.) As the corresponding H commutes with γ_0 in this case, the potential \tilde{f} for F_a can be chosen to be H -invariant (by averaging over H). By the argument in [19], the \tilde{f} can be deformed to define an H -invariant potential for a lcK metric on S_λ , for λ small enough, with the same constant c_{γ_0} (see Definition 3.6). Using the isomorphism $\mathbb{C}^* = H^1(S, \mathbb{C}^*) \cong \text{Pic}_0(S)$ established for class VII surfaces in [28, I, p. 756], this shows that the induced lcK metrics on S_λ will have Lee forms in the same de Rham class a . We thus see that $\mathcal{C}(S_\lambda) = (-\infty, 0)$ too.

Using Proposition 3.4 (and Lemma 3.3 with $a = 0$ and $k = 1$), we conclude

Proposition 5.1. *For any compact complex surface S whose minimal model is a Hopf surface, $\mathcal{T}(S) = \mathcal{C}(S) = (-\infty, 0)$.*

5.2. Inoue surfaces with $b_2 = 0$. The Inoue surfaces S_0 with second Betti number equal to zero are classified by Inoue [24] into three types, $S_M, S_{N; p, q, r}^-$ and $S_{N; p, q, r; u}^+$ where the parameters M, N are matrices with integer coefficients, p, q, r are integers and u is a complex number. Any such surface is the quotient of $\mathbb{C} \times \mathbb{H}$ (where \mathbb{H} denotes the upper half-plane in \mathbb{C}) under a discrete subgroup Γ of the group $A(2, \mathbb{C})$ of affine transformations of \mathbb{C}^2 , leaving $\mathbb{C} \times \mathbb{H}$ invariant. The specific description of Γ in each case is given in [24], but we shall not make use of this. The relevant information for the discussion below is the fact, shown by Tricerri in [48], that all Inoue surfaces admit lcK metrics, except the surfaces of the type $S_u := S_{N; p, q, r; u}^+$ for which the complex parameter u is not real. In the later case, Belgun [6] shows that there are no lcK metrics at all. Nevertheless, we proved in Theorem 1.1 that S_u always admits timing locally conformally symplectic structures.

Let us now consider in a little more detail the analytic family S_u of Inoue surfaces of the third type. We shall show, by using an argument from [6], that in this case $\mathcal{T}(S_u)$ is a single point.

It is known [6, p. 35], [23, Thm. 1], [50] that $S_u = (\mathbb{C} \times \mathbb{H})/\Gamma_u$ where Γ_u is a lattice in the solvable Lie group

$$\mathrm{Sol}_1^4 = \left\{ \begin{pmatrix} 1 & a & b + i \log \gamma \\ 0 & \gamma & c \\ 0 & 0 & 1 \end{pmatrix}, \gamma > 0, a, b, c \in \mathbb{R} \right\} \subset \mathrm{GL}(3, \mathbb{C}).$$

The group Sol_1^4 acts itself simply-transitively (and holomorphically) on $\mathbb{C} \times \mathbb{H}$. This allows to identify S_u with the quotient $\mathrm{Sol}_1^4/\Gamma_u$, endowed with a (fixed) left-invariant integrable almost complex structure J , by the left action of the lattice Γ_u . In explicit terms, let

$$Y = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad Z = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad T = \begin{pmatrix} 0 & 0 & i \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad U = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix}$$

be the generators of the Lie algebra of Sol_1^4 with

$$Z \text{ central, } [Y, T] = Y, [T, U] = U, [Y, U] = Z$$

and denote with the same letters the induced left-invariant vector fields on Sol_1^4 . Then the left-invariant complex structure J on Sol_1^4 is given by [6, (35)]

$$JY = -Z, JZ = Y, JT = -U - Z, JU = T - Y.$$

Furthermore, Sol_1^4 admits an ad-invariant 1-form α_0 , defined by $\alpha_0(T) = 1, \alpha_0(U) = \alpha_0(Y) = \alpha_0(Z) = 0$, which descends to define a closed but not exact 1-form (still denoted α_0) on S_u . As $b_1(S_u) = 1$, by Remark 2.2 we can assume without loss that S_u admits a locally conformally symplectic 2-form F which tames J , and whose Lee form equals $k\alpha_0$ for a non-zero real constant k . As Sol_1^4 has a bi-invariant volume form v [20, Lemma 4] (which defines a volume form on the quotient S_u , still denoted by v), for any left-invariant vector fields U and V on Sol_1^4 (which define vector fields on the quotient S_u , still denoted by U and V), one can consider the average of F over S_u :

$$F_0(U, V) := \int_{S_u} F(U, V)v.$$

It can be shown, as in the proof of [6, Thm. 7], that F_0 defines a *left-invariant* 2-form on (Sol_1^4, J) which tames J and satisfies $dF_0 = k\alpha_0 \wedge F_0$. Evaluating the later equality over Y, Z, T yields (see also [6, (36)]) $F_0(Y, Z) = kF(Y, Z)$; as $Y = JZ$ and the $(1, 1)$ -part of F_0 is positive definite, it follows that $k = 1$. This shows that $\mathcal{T}(S_u) = \{a_0\}$ with $a_0 = [\alpha_0]$. It is easy to check (see [20, Sect. 7.1]), using the explicit description of S_u of [24] and the isomorphism $\mathbb{C}^* = H^1(S, \mathbb{C}^*) \cong \mathrm{Pic}_0(S)$ of [28, I, p. 756], that the corresponding holomorphic line bundle \mathcal{L}_{a_0} is isomorphic to the anticanonical line bundle $\mathcal{K}_{S_u}^*$.

Noting finally that the Inoue surfaces of the type $S_{N,p,q,r,s}^-$ are quotients of Inoue surfaces of the type $S_{N,p,q,r,0}^+$ by an involution [24], we obtain the following

Proposition 5.2. *Let S be an Inoue surface with $b_2(S) = 0$ in one of the types $S_{N,p,q,r,u}^+$ or $S_{N,p,q,r}^-$. Then $\mathcal{C}(S)$ and $\mathcal{T}(S)$ are given as follows:*

- $\mathcal{C}(S) = \mathcal{T}(S) = \{a_0\}$ iff S is of the type $S_{N,p,q,r}^-$ or $S_{N,p,q,r,u}^+$ with $u \in \mathbb{R}$;
- $\mathcal{C}(S) = \emptyset, \mathcal{T}(S) = \{a_0\}$ iff S is of the type $S_{N,p,q,r,u}^+$ with $u \in \mathbb{C} \setminus \mathbb{R}$,

where $\mathcal{L}_{a_0} = \overline{\mathcal{K}_{S_u}^*}$.

5.3. Kato surfaces. These are minimal complex surfaces in the Kodaira class VII whose second Betti number is strictly greater than 0, and which have a *global spherical shell* (GSS). Conjecturally, any minimal surface in the class VII should be either a Hopf surface, an Inoue surface, or a Kato surface. However, this conjecture is still far from being solved.

For Kato complex surfaces, Brunella [9, 10] has shown that $\mathcal{C}(S) \neq \emptyset$ and that $\mathcal{C}(S)$ has $-\infty$ as an accumulation point. Note that any Kato surface S is diffeomorphic to $(S^1 \times S^3) \# k \overline{CP^2}$ (where $k := b_2(S)$), see e.g. [34]. As $S^1 \times S^3$, with a complex structure of a Hopf surface of class 1, admits a Vaisman metric, it follows by [32, 39] that $H_{d_L}^3(S^1 \times S^3, L) = \{0\}$ for any non-trivial flat real line bundle L . By Lemma 3.3, $H_{d_L}^3(S, L) = \{0\}$ and, therefore, by Remark 3.2, $\mathcal{T}(S)$ must be an open subset of $(-\infty, 0)$. Similar conclusion holds true for $\mathcal{C}(S) \subseteq \mathcal{T}(S)$, by [20] and Lemma 2.8, together with the vanishing of $H^2(S, \mathcal{L})$ and $H^0(S, \mathcal{L})$ established in Lemma 2.10 and Proposition 4.3, respectively.

Further progress in this case seems to depend on a better understanding of the subset $\mathcal{G}(S)$, as the following result suggests.

Proposition 5.3. *Let S be a Kato surface. Then,*

$$\mathcal{G}(S) \subseteq \mathcal{T}(S).$$

Proof. Let $c \in \mathcal{G}(S)$. Brunella shows (see [9, Rem. 9]) that there exists a lcK metric whose Lee form defines a de Rham class b with $b < c$. By Proposition 4.4 and Lemma 2.10, any $a \in [b, c]$ (in particular $a = c$) belongs to $\mathcal{T}(S)$. \square

APPENDIX A. A PERRON-TYPE THEOREM FOR SECOND-ORDER STRONGLY ELLIPTIC LINEAR OPERATORS ON A COMPACT RIEMANNIAN MANIFOLD

We review here some spectral properties of the second order strongly elliptic linear operators in the form

$$L(u) = \Delta^g u + g(\alpha, du) + cu,$$

where g is a Riemannian metric on a compact manifold M , $\Delta^g = \delta^g d$ is the corresponding Riemannian Laplacian (which we shall consequently denote by Δ), α is a given smooth 1-form and c a given smooth function on M . Note that L need not to be self-adjoint in general, a case where the spectral theory is well-established.

In a local chart U , L takes the form

$$L_U = - \sum_{i,j=1}^n a_U^{ij}(x) \frac{\partial^2}{\partial x_i \partial x_j} + \sum_{k=1}^n b_U^k(x) \frac{\partial}{\partial x_k} + c_U(x),$$

where a_U^{ij}, b_U^k, c_U are smooth real functions such that the symmetric matrix (a_U^{ij}) is uniformly positive definite on U . In this case, the Perron-type Theorem established in [16, p. 360] and [35] states that if $\Omega \subset U$ is compactly supported domain with smooth boundary, the operator L_U taken on smooth functions on $\bar{\Omega}$ with zero boundary value has *real* eigenvalue λ_0 (called *principal eigenvalue*) such that for any other eigenvalue λ , $\operatorname{Re}(\lambda) \geq \lambda_0$; furthermore λ_0 is simple and the corresponding eigenspace is generated by a nowhere vanishing smooth function u_0 on Ω .

Recall that for any strongly elliptic linear operator $L : \mathcal{C}^\infty(M) \rightarrow \mathcal{C}^\infty(M)$, the set of (complex) eigenvalues is discrete, having a limit point only at infinity, see e.g. [7,

p. 465] or [4, p. 126]. We want to establish the following adaptation of the Perron-type Theorem mentioned above to the case of a compact manifold without boundary (as we failed to find a reference to this result in the literature).

Theorem A.1. *Let $L(u) = \Delta u + g(\alpha, du) + cu$ be a linear strongly elliptic linear differential operator of order 2 on a compact Riemannian manifold (M, g) . Then, there exists a real eigenvalue λ_0 for L which admits a smooth everywhere positive eigenfunction u_0 . Furthermore, λ_0 satisfies the following properties:*

- (i) λ_0 is of multiplicity one.
- (ii) If λ is another eigenvalue, then $\operatorname{Re}(\lambda) > \lambda_0$.
- (iii) $\lambda_0 = \sup_{u \in \mathcal{A}} \left(\inf_{x \in M} \frac{L(u)}{u} \right)$, where $\mathcal{A} = \{u \in C^\infty(M) \mid u > 0\}$.

Definition A.2. The eigenvalue λ_0 is called the *principal eigenvalue* of L .

Proof of Theorem A.1. The proof will be decided into four steps, corresponding to the statements in the Theorem A.1 as follows.

Step 1. We shall establish in this Step the existence of a real eigenvalue λ_0 corresponding to an everywhere positive smooth eigenfunction u_0 . To this end, we shall work with the Sobolov spaces $W_k^2(M)$ corresponding to the norm

$$\|f\|_k := \left[\sum_{0 \leq j \leq k} \int_M |D^j f|^2 v_g \right]^{1/2},$$

where $|D^j f|$ is the point-wise norm of the j -th covariant derivative and v_g is the Riemannian volume form. In what follows, we shall choose k sufficiently large so that we have a continuous embedding $W_k^2(M) \subset C^2(M)$. Because M is compact, we also have the continuous embeddings

$$W_k^2(M) \subset W_l^2(M),$$

for any integers $0 \leq l \leq k$.

We shall assume (without loss) in what follows that the smooth function $c \geq 0$ (otherwise we consider the operator $L - (\inf_M c)\operatorname{Id}$ instead of L). Then, by the maximum principle [4, III, Sect. 8, 3.71] (which in the sequel we shall always apply to $-L$ and $-u$), 0 is not an eigenvalue of L , i.e. $\operatorname{Ker} L = \{0\}$. It is a standard fact that $L : W_{k+2}^2(M) \rightarrow W_k^2(M)$ is then invertible, and $L^{-1} : W_k^2(M) \rightarrow W_{k+2}^2(M)$ is bounded. Indeed, as $L = \Delta + T$ with Δ being the self-adjoint Riemannian Laplacian and T of order ≤ 1 , the composition $T : W_{k+2}^2(M) \rightarrow W_{k+1}^2(M) \subset W_k^2(M)$ is a compact operator by the Rellich–Kondrachov theorem (see [7, p. 458]); it follows that $\operatorname{Index}(L) = \operatorname{Index}(\Delta) = 0$ by [21, Cor. 19.1.8] and, as $\operatorname{Ker} L = \{0\}$, L is invertible and bounded (by the standard L^p -estimates, see e.g. [7, p. 463]). Using the compactness of $W_{k+1}^2(M) \subset W_k^2(M)$ again, we conclude that the composition

$$A : W_k^2(M) \xrightarrow{L^{-1}} W_{k+2}^2 \subset W_k^2(M)$$

is a compact operator. We define the cone

$$(16) \quad C := \{u \in W_k^2(M) \mid u \geq 0\}.$$

By the maximum principle [4, III, Sect. 8, 3.71] again, we have that $A(C) \subset C$ and, if $w \in C$ with $w \not\equiv 0$, then $Aw > 0$. Using standard elliptic regularity (see e.g. [7, p. 467]) it will be enough to show that A has a non-trivial eigenfunction $w_0 \in C$,

corresponding to a real positive eigenvalue $1/\lambda_0$ (we will have then that $u_0 := \lambda_0 A(w_0)$ is a smooth strictly positive eigenfunction of L , corresponding to the real positive eigenvalue λ_0).

Let us fix a function $w \in C$, $w \neq 0$, and let $v := Aw$. By continuity, there exists $\mu > 0$ such that

$$\mu v \geq w$$

on M . We claim that if the equation

$$u = \lambda A(u + \epsilon w)$$

with $\epsilon > 0$ and $\lambda > 0$ have an everywhere positive solution u , then necessarily

$$(17) \quad \lambda \leq \mu.$$

Indeed,

$$u = \lambda A(u) + \lambda A(\epsilon w) > \lambda A(\epsilon w) = \lambda \epsilon v \geq \left(\frac{\lambda}{\mu}\right) \epsilon w$$

hence

$$u \geq \lambda A(u) \geq \lambda A\left(\frac{\lambda \epsilon}{\mu} w\right) = \frac{\lambda^2}{\mu} \epsilon v \geq \left(\frac{\lambda}{\mu}\right)^2 \epsilon w.$$

By induction, for $k \geq 1$,

$$u \geq \left(\frac{\lambda}{\mu}\right)^k \epsilon w,$$

which is possible only if $\lambda \leq \mu$.

We are now going to show that for any $\epsilon > 0$, the closed subset of $W_k^2(M)$

$$S_\epsilon := \{u \in C \mid \exists \lambda, 0 \leq \lambda \leq 2\mu, u = \lambda A(u + \epsilon w)\}$$

is unbounded. To this end, we shall use (17) in conjunction with the well-known Schaefer theorem (see [16, p. 540]), which states that if $\tilde{A} : W \rightarrow W$ is a continuous compact (not necessarily linear) mapping of a Banach space W and $C \subset W$ is a convex subset stable by \tilde{A} , then \tilde{A} has a fixed point in C provided that the subset

$$\{u \in C \mid \exists \lambda, 0 \leq \lambda \leq 1 \text{ such that } x = \lambda \tilde{A}(x)\}$$

is bounded.

In our case, $W = W_k^2(M)$,

$$\tilde{A}(u) := 2\mu(A(u) + \epsilon v)$$

which is continuous compact because A is, and $C \subset W_k^2(M)$ is the convex subset introduced in (16). Furthermore, if S_ϵ were bounded, \tilde{A} would satisfy the hypothesis of the Schaefer theorem with respect to C , so \tilde{A} would have a fixed point u in C , i.e. $u = \tilde{A}u = 2\mu A(u + \epsilon w)$, which contradicts (17).

Since S_ϵ is not bounded, there exists $u_\epsilon \in S_\epsilon$ with $\|u_\epsilon\|_k \geq \frac{1}{\epsilon}$. Denote by λ_ϵ ($0 \leq \lambda_\epsilon \leq 2\mu$) the corresponding real number such that

$$u_\epsilon = \lambda_\epsilon A(u_\epsilon + \epsilon w).$$

For any sequence $\epsilon_m \rightarrow 0$ let $\lambda_m := \lambda_{\epsilon_m}$ and $u_m := \frac{u_{\epsilon_m}}{\|u_{\epsilon_m}\|_k} \in C$, so that

$$(18) \quad u_m = \lambda_m A\left(u_m + \epsilon_m \frac{w}{\|u_{\epsilon_m}\|_k}\right).$$

Since A is compact, we can take a sub-sequence if necessary and suppose that (λ_m) and (u_m) are convergent. Let $\lambda_0 := \lim_{m \rightarrow \infty} \lambda_m$ and $u_0 := \lim_{m \rightarrow \infty} u_m \geq 0$. Taking limit in (18), we get

$$u_0 = \lambda_0 A(u_0)$$

The condition $\|u_m\|_k = 1$ prevents λ_0 and u_0 to vanish. Thus, $u_0 = \lambda_0 A(u_0) > 0$, and applying L to the above equality we obtain $L(u_0) = \lambda_0 u_0$ with $\lambda_0 > 0$.

Step 2. We shall now prove that the multiplicity of λ_0 is one.

As L is a real operator, it is enough to consider a smooth non-identically zero real valued function u on M , such that $Lu = \lambda_0 u$. Furthermore, replacing u by $-u$ if necessary, we shall assume that u is somewhere positive. Thus, letting

$$\chi = \sup\{\mu > 0 \mid u_0 - \mu u \geq 0 \text{ on } M\},$$

we then have $\chi > 0$, $v := u_0 - \chi u \geq 0$ on M , and hence $Lv = \lambda_0 v \geq 0$.

By the Hopf maximum principle [4, III, Sect. 8, 3.71], we have that either $v > 0$ on M or $v \equiv 0$. By the definition of χ we conclude that $v \equiv 0$.

Step 3. We now show that for any other (complex) eigenvalue λ of L , $\operatorname{Re}(\lambda) > \lambda_0$.

Let $u \not\equiv 0$ be a complex-valued smooth function on M with $Lu = \lambda u$ and let $v := u/u_0$. We then have

$$\begin{aligned} \lambda u_0 v &= L(u_0 v) = vL(u_0) + u_0 L(v) - cu_0 v - 2g(du_0, dv) \\ &= \lambda_0 u_0 v + u_0 (\Delta v + g(\alpha - 2d \log u_0, dv)) \\ &= \lambda_0 u_0 v + u_0 K(v), \end{aligned}$$

where we have set

$$K(v) := \Delta(v) + g(\alpha - 2d \log u_0, dv).$$

Dividing by u_0 the last equality, we have

$$K(v) = (\lambda - \lambda_0)v,$$

and as K is a real operator

$$K(\bar{v}) = (\bar{\lambda} - \lambda_0)\bar{v}.$$

It follows

$$\begin{aligned} (19) \quad K(|v|^2) &= K(v\bar{v}) = vK(\bar{v}) + \bar{v}K(v) - 2g(dv, d\bar{v}) \\ &= 2(\operatorname{Re}(\lambda) - \lambda_0)|v|^2 - 2g(dv, d\bar{v}) \\ &\leq 2(\operatorname{Re}(\lambda) - \lambda_0)|v|^2. \end{aligned}$$

Suppose for contradiction that $\operatorname{Re}(\lambda) - \lambda_0 \leq 0$. Then, the Hopf maximum principle [4, III, Sect. 8, 3.71] implies $|v| = \text{const}$, and therefore

$$0 = K(|v|^2) \leq 2(\operatorname{Re}(\lambda) - \lambda_0)|v|^2$$

showing that $\operatorname{Re}(\lambda) = \lambda_0$ (as $v = u/u_0 \not\equiv 0$). Going back to (19) (and using again $K(|v|^2) = 0$), we have

$$\begin{aligned} 0 &= K(|v|^2) = -2g(dv, d\bar{v}) + 2(\operatorname{Re}(\lambda) - \lambda_0)|v|^2 \\ &= -2\|dv\|_g^2 \end{aligned}$$

i.e. $v = u/u_0$ is a constant. As $\lambda \neq \lambda_0$ by assumption, this is a contradiction.

Step 4. We finally have to prove that

$$\lambda_0 = \sup_{u \in \mathcal{A}} \inf_{x \in M} \frac{L(u)}{u}$$

where $\mathcal{A} = \{u \in C^\infty(M) \mid u > 0\}$.

To this end, we shall show first that the formal conjugate operator

$$L^*(u) = \Delta u - g(\alpha, du) + (\delta^g \alpha + c)u$$

of L (with respect to the global L_2 product on (M, g)) has the same principal eigenvalue as L . Indeed, let λ_0^* be the principal eigenvalue of L^* and $u_0^* > 0$ an eigenfunction. Then

$$\lambda_0 \int_M u_0 u_0^* v_g = \int_M L(u_0) u_0^* v_g = \int_M u_0 L^*(u_0^*) v_g = \lambda_0^* \int_M u_0 u_0^* v_g.$$

Since $u_0 > 0$ and $u_0^* > 0$, it follows that $\lambda_0 = \lambda_0^*$.

Now, let

$$\mu := \sup_{u \in \mathcal{A}} \inf_{x \in M} \frac{L(u)}{u}.$$

We clearly have $\mu \geq \inf_{x \in M} \frac{L(u_0)}{u_0} = \lambda_0$. To establish the converse inequality, let (u_m) be a maximizing sequence in \mathcal{A} such that

$$(20) \quad \mu - \frac{1}{m} \leq \inf_{x \in M} \frac{L(w_m)}{w_m} \leq \mu.$$

It then follows

$$\lambda_0 \int_M w_m u_0^* v_g = \int_M w_m L^*(u_0^*) v_g = \int_M L(w_m) u_0^* v_g \geq \left(\mu - \frac{1}{m}\right) \int_M w_m u_0^* v_g,$$

thus showing $\lambda_0 \geq \mu$. \square

The following result follows from the general Rellich–Kato theory, see e.g. [26, II, Sect. 1.8; IV, Theorem 3.16, and VII, Thm. 1.7] applied to the bounded operator A constructed in the proof of Theorem A.1, by noting that the principal eigenvalue λ_0 of L is simple and can be separated from the reminder of the spectrum.

Theorem A.3. *Let $L(t)$ be an analytic family of linear strongly elliptic operators as in Theorem A.1. For each t , denote by $\lambda_0(t)$ the principal eigenvalue of $L(t)$ with corresponding eigenfunction $u_t > 0$, normalized by $\int_M u_t^2 v_g = 1$. Then $\lambda_0(t)$ and u_t vary analytically with respect to t .*

REFERENCES

- [1] V. Apostolov, M. Bailey, and G. Dloussky, *From locally conformally Kähler to bihermitian structures on non-Kähler complex surfaces*, to appear in Math. Res. Letters, arxiv:1307.3660v1 (2013).
- [2] V. Apostolov, P. Gauduchon, and G. Grantcharov, *Bihermitian structures on complex surfaces*, Proc. London Math. Soc. (3) **79** (1999), 414–428. *Corrigendum*, **92** (2006), 200–202.
- [3] V. Apostolov, M. Gualtieri, *Generalized Kähler manifolds, commuting complex structures and split tangent bundles*, Commun. Math. Phys. **271** (2007), 561–575.
- [4] Th. Aubin, *Some Nonlinear Problems in Riemannian Geometry*, *Springer Monographs*, 1998.
- [5] W. Barth, K. Hulek, C. Peters, and A. Van de Ven, *Compact Complex Surfaces*, *Springer, Heidelberg*, Second Edition, 2004.
- [6] F. Belgun, *On the metric structure of non-Kähler complex surfaces*, Math. Ann. **317** (2000), 1–40.
- [7] A. L. Besse, *Einstein manifolds*, *Ergeb. Math. Grenzgeb.* **3**, Springer-Verlag, Berlin, Heidelberg, New York, 1987.

- [8] F. Bogomolov, *Classification of surfaces of class VII₀ with $b_2 = 0$* , Math. USSR Izv **10** (1976), 255–269é
- [9] M. Brunella, *Locally conformally Kähler metrics on certain non-Kählerian surfaces*, Math. Ann. (2010) **346** 629–639.
- [10] M. Brunella, *Locally conformally Kähler metrics on Kato surfaces*, Nagoya Math. J. **202**, 77–81.
- [11] N. Buchdahl, *On compact Kähler surfaces*, Ann. Inst. Fourier **49** (1999), 287–302.
- [12] N. Buchdahl, *A Nakai-Moishezon criterion for non-Kähler surfaces*, Ann. Inst. Fourier **50** (2000), 1533–1538.
- [13] J.-P. Demailly, M. Paun, *Numerical characterization of the Kähler cone of a compact Kähler manifold*, Ann. Math. **159** (2004), 1247–1274.
- [14] S. Dragomir, L. Ornea, *Locally coformal Kähler geometry*, Birkhauser, 1998.
- [15] S. K. Donaldson, *Two-forms on four-manifolds and elliptic equations*, in ‘Inspired by S. S. Chern’, 153–172, Nankai Tracts Math., **11**, World Sci. Publ., Hackensack, NJ, 2006.
- [16] L. C. Evans, *Partial differential equations Graduate Studies in Mathematics*, vol 19, AMS Providence, second edition, 2010.
- [17] P. Gauduchon, *Le théorème d’excentricité nulle*, C. R. Acad. Sci. Paris **285** (1977), 387–390.
- [18] P. Gauduchon, *La 1-forme de torsion d’une variété hermitienne* Math. Ann. **267** (1984), 495–518.
- [19] P. Gauduchon, L. Ornea *Locally conformally Kähler metrics on Hopf surfaces*, Ann. Inst. Fourier (Grenoble), **48** (1998), 1107–1127.
- [20] R. Goto, *On the stability of locally conformal Kähler structures*, available at arxiv 1012.2285v1 (2010).
- [21] L. Hormander, *The Analysis of Linear Partial Differential Operators III*, Springer-Verlag, 1994.
- [22] R. Harvey, H. Blaine Lawson, Jr. *An intrinsic characterization of Kähler manifolds*, Invent. math. **74** (1983), 169–198.
- [23] K. Hasegawa, *Complex and Kähler structures on compact solvmanifolds*. J. Symplectic Geom. **3** (2005), 749–767.
- [24] M. Inoue, *On surfaces of class VII₀*, Invent. math. **24** (1974), 269–310.
- [25] M. Kato, *Topology of Hopf surfaces*, J. Math. Soc. Japan **27** (1975), 222–238. Erratum J. Math. Soc. Japan **41** (1989), 173–174.
- [26] T. Kato, *Perturbation Theory for Linear Operators*, Springer-Verlag, 1980.
- [27] S. Kobayashi, K. Nomizu, *Foundations of Differential Geometry. Vol.II*, Newyork: Interscience Publishers, **179**, 1963.
- [28] K. Kodaira, *On the structure of complex analytic surfaces I, II, III*, Am. J. Math **86** (1966), 751–798; Am. J. Math. **88** (1966), 682–721; Am. J. Math. **90** (1968), 55–83.
- [29] G. Kokarev, *On pseudo-harmonic maps in conformal geometry*, Proc. London Math. Soc. **99** (2009), 168–194.
- [30] A. Lamari, *Courants kählériens et surfaces compactes*, Ann. Inst. Fourier **49** (1999), 263–285.
- [31] A. Lamari, *Le cone kählérien d’une surface*, J. Math. Pures Appl. **78** (1999), 249–263.
- [32] M. de León, B. López, J. C. Marrero, E. Padrón, *On the computation of the Lichnerowicz–Jacobi cohomology*, J. Geom. Phys. **44** (2003), 507–522.
- [33] J. Li, S. T. Yau, *Hermitian Yang–Mills connections on non-Kähler manifolds*, Math. aspects of string theory (San Diego, Calif., 1986), Adv. Ser. Math. Phys. 1, 560–573, World Scientific Publishing (1987).
- [34] I. Nakamura, *On surfaces of class VII₀ with curves*, Invent. math. **78** (1984), 393–343.
- [35] L. Ni, *A Perron-type theorem on the principal eigenvalue of nonsymmetric elliptic operator* to appear in The Am. Math. Monthly.
- [36] L. Ornea, M. Verbitsky, *A report on locally conformally Kähler manifolds. Harmonic maps and differential geometry*, 135–149, Contemp. Math. **542**, Amer. Math. Soc., Providence, RI, 2011.
- [37] L. Ornea, M. Verbitsky, *Topology of Locally Conformally Kähler Manifolds with Potential*, IMRN **2010** (2009), 717–726.
- [38] L. Ornea, M. Verbitsky, *Locally conformal Kähler manifolds with potential*, Math. Ann. **348** (2010), 25–33.
- [39] L. Ornea, M. Verbitsky, *Morse-Novikov cohomology of locally conformally Kähler manifolds*, J. Geom. Phys. **59** (2009), 295–305.
- [40] L. Ornea, M. Verbitsky, *Compact pluricanonical manifolds are Vaisman*, in preparation.
- [41] M. Pontecorvo, *Complex structures on Riemannian 4-manifolds*, Math. Ann. **309** (1997), 159–177.

- [42] F. Rellich, Perturbation theory of eigenvalue problems. *New York, Gordon and Breach*, 1969.
- [43] Y.-T. Siu, *Every K3 surface is Kähler*, Invent. Math. **73** (1983), 139–150.
- [44] C. H. Taubes, *Tamed to compatible: Symplectic forms via moduli space integration*, J. Symplectic Geom. **9** (2011), 161–250.
- [45] A. Teleman, *Projectively flat surfaces and Bogomolov’s theorem on class VII₀-surfaces*, Int. J. Math., **5** (1994), 253–264.
- [46] A. Teleman, *The pseudo-effective cone of a non-Kählerian surface and applications*, Math. Ann. **335** (2006), 965–989.
- [47] A. Todorov, *Applications of the Kähler-Einstein-Calabi-Yau metric to moduli of K3 surfaces*, Invent. math. **61** (1980), 251–265.
- [48] F. Tricerri, *Some examples of locally conformal Kähler manifolds*, Rend. Sem. Mat. Univers. Politecn. Torino **40** (1982), 81–92.
- [49] V. Vuletescu, *Blowing-up points on locally conformally Kähler manifolds*, Bull. Math. Soc. Sci. Math. Roumanie **52** (2009), 387–390.
- [50] C. I. C. Wall, *Geometric structures on compact complex analytic surfaces*. Topology **25** (1986), 119–153.

VESTISLAV APOSTOLOV, DÉPARTEMENT DE MATHÉMATIQUES, UQAM, C.P. 8888, SUCC. CENTRE-VILLE, MONTRÉAL (QUÉBEC), H3C 3P8, CANADA
E-mail address: `apostolov.vestislav@uqam.ca`

GEORGES DLOUSSKY, AIX-MARSEILLE UNIVERSITY, CNRS, CENTRALE MARSEILLE, I2M, UMR 7373, 13453, 39 RUE F. JOLIOT-CURIE 13411, MARSEILLE CEDEX 13, FRANCE
E-mail address: `georges.dloussky@univ-amu.fr`