



HAL
open science

Infinite Dimensional Weak Dirichlet Processes and Convolution Type Processes

Giorgio Fabbri, Francesco Russo

► **To cite this version:**

Giorgio Fabbri, Francesco Russo. Infinite Dimensional Weak Dirichlet Processes and Convolution Type Processes. *Stochastic Processes and their Applications*, 2017, 127 (1), pp.325-357. 10.1016/j.spa.2016.06.010 . halshs-01309384

HAL Id: halshs-01309384

<https://shs.hal.science/halshs-01309384>

Submitted on 29 Apr 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Infinite Dimensional Weak Dirichlet Processes and Convolution Type Processes

Giorgio Fabbri
Francesco Russo

WP 2016 - Nr 16

Infinite dimensional weak Dirichlet processes and convolution type processes.

Giorgio Fabbri* and Francesco Russo†

April 20th 2016

Abstract

The present paper continues the study of infinite dimensional calculus via regularization, started by C. Di Girolami and the second named author, introducing the notion of *weak Dirichlet process* in this context. Such a process \mathbb{X} , taking values in a Banach space H , is the sum of a local martingale and a suitable *orthogonal* process.

The concept of weak Dirichlet process fits the notion of *convolution type processes*, a class including mild solutions for stochastic evolution equations on infinite dimensional Hilbert spaces and in particular of several classes of stochastic partial differential equations (SPDEs).

In particular the mentioned decomposition appears to be a substitute of an Itô's type formula applied to $f(t, \mathbb{X}(t))$ where $f : [0, T] \times H \rightarrow \mathbb{R}$ is a $C^{0,1}$ function and \mathbb{X} a convolution type processes.

Key words and phrases: Covariation and Quadratic variation; Calculus via regularization; Infinite dimensional analysis; Tensor analysis; Dirichlet processes; Generalized Fukushima decomposition; Convolution type processes; Stochastic partial differential equations.

2010 AMS Math Classification: 60H05, 60H15, 60H30, 35R60.

1 Introduction

Stochastic calculus via regularization for real processes was initiated in [34] and [35]. It is an efficient calculus for non-semimartingales whose related literature is surveyed in [37]. We briefly recall some essential notions from [34] and [35]. Let $T > 0$ and $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space. Given Y (respectively X) an a.s. bounded (respectively continuous) real process¹ defined on $[0, T]$, the forward integral of Y with respect to X and the covariation between Y and X are defined as follows. Suppose that, for every $t \in [0, T]$, the limit $I(t)$ (respectively $C(t)$) in probability exists:

$$\begin{aligned} I(t) : &= \lim_{\epsilon \rightarrow 0^+} \int_0^t Y(r) \left(\frac{X(r+\epsilon) - X(r)}{\epsilon} \right) dr, \\ C(t) : &= \lim_{\epsilon \rightarrow 0^+} \int_0^t \frac{(X(r+\epsilon) - X(r))(Y(r+\epsilon) - Y(r))}{\epsilon} dr. \end{aligned} \tag{1}$$

*Aix-Marseille University (Aix-Marseille School of Economics), CNRS and EHESS. e-mail: giorgio.fabbri@univ-amu.fr, Centre de la vieille Charité - 2 rue de la Charité - 13002 Marseille - France. E-mail: giorgio.fabbri@univ-amu.fr

†ENSTA ParisTech, Université Paris-Saclay, Unité de Mathématiques appliquées, 828, boulevard des Maréchaux, F-91120 Palaiseau. E-mail: francesco.russo@ensta-paristech.fr

¹In the whole paper, all the considered processes are supposed to be measurable from $[0, T] \times \Omega$ (equipped with the product of the Borel σ -field of $[0, T]$ and the σ -algebra of all events \mathcal{F}) and the value space, for instance \mathbb{R} , endowed with the Borel σ -field.

If the random process I (respectively C) admits a continuous version, this is denoted by $\int_0^\cdot Y d^-X$ (respectively $[X, Y]$). It is the *forward integral* of Y with respect to X (respectively the *covariation* of X and Y). If X is a real continuous semimartingale and Y is a càdlàg process which is progressively measurable (respectively a semimartingale), the integral $\int_0^\cdot Y d^-X$ (respectively the covariation $[X, Y]$) is the same as the classical Itô's integral (respectively covariation).

Real processes X for which $[X, X]$ exists are called *finite quadratic variation processes*. A rich class of finite quadratic variation processes is provided by Dirichlet processes. Let $(\mathcal{F}_t, t \in [0, T])$ be a fixed filtration, fulfilling the usual conditions. A real process X is said to be *Dirichlet* (or *Föllmer-Dirichlet*) if it is the sum of a local martingale M and a *zero quadratic variation process* A , i.e. such that $[A, A] = 0$. Those processes were defined by H. Föllmer [17] using limits of discrete sums. A significant generalization, due to [15, 21], is the notion of *weak Dirichlet process*, extended to the case of jump processes in [3].

Definition 1.1. *A real process $X: [0, T] \times \Omega \rightarrow \mathbb{R}$ is called weak Dirichlet process if it can be written as $X = M + A$, where M is a local martingale and A is a process such that $[A, N] = 0$ for every continuous local martingale N and $A(0) = 0$.*

Obviously a semimartingale is a weak Dirichlet process. It can be proved (see Remark 3.5 of [21]) that the decomposition described in Definition 1.1 is unique.

Elements of calculus via regularization were extended to (real) Banach space valued processes in a series of papers, see e.g. [8, 9, 10, 11]. Two classical notions of stochastic calculus in Banach spaces, which appear in [28] and [13] are the scalar² and tensor quadratic variations. We propose here, following [11], a definition of them using the regularization approach, even though, originally they appeared in a discretization framework.

Definition 1.2. *Consider a real separable Banach spaces B . We say that a process $\mathbb{X}: [0, T] \times \Omega \rightarrow B$, a.s. square integrable, admits a **scalar quadratic variation** if, for any $t \in [0, T]$, the limit, for $\epsilon \searrow 0$ of*

$$[\mathbb{X}, \mathbb{X}]^{\epsilon, \mathbb{R}}(t) := \int_0^t \frac{|\mathbb{X}(r + \epsilon) - \mathbb{X}(r)|_B^2}{\epsilon} dr$$

exists in probability and it admits a continuous version. The limit process is called scalar quadratic variation of \mathbb{X} and it is denoted by $[\mathbb{X}, \mathbb{X}]^{\mathbb{R}}$.

Definition 1.3. *Consider two real separable Banach spaces B_1 and B_2 . Suppose that either B_1 or B_2 is different from \mathbb{R} . Let $\mathbb{X}: [0, T] \times \Omega \rightarrow B_1$ and $\mathbb{Y}: [0, T] \times \Omega \rightarrow B_2$ be two a.s. square integrable processes. We say that (\mathbb{X}, \mathbb{Y}) admits a **tensor covariation** if the limit, for $\epsilon \searrow 0$ of the $B_1 \hat{\otimes}_\pi B_2$ -valued processes*

$$[\mathbb{X}, \mathbb{Y}]^{\otimes, \epsilon} := \int_0^\cdot \frac{(\mathbb{X}(r + \epsilon) - \mathbb{X}(r)) \otimes (\mathbb{Y}(r + \epsilon) - \mathbb{Y}(r))}{\epsilon} dr,$$

*exists ucp³. The limit process is called tensor covariation of (\mathbb{X}, \mathbb{Y}) and is denoted by $[\mathbb{X}, \mathbb{Y}]^{\otimes}$. The tensor covariation $[\mathbb{X}, \mathbb{X}]^{\otimes}$ is called **tensor quadratic variation** of \mathbb{X} and is denoted by $[\mathbb{X}]^{\otimes}$.*

The concepts of scalar and tensor quadratic variation are too strong in certain contexts: several interesting examples of Banach (or even Hilbert) space valued processes have no tensor quadratic variation. For this reason Di Girolami and Russo introduced (see for instance Definition 3.4 of [10]) the notion of χ -covariation. It is recalled in Definition 3.1 and widely use in this work. Their idea was to introduce a suitable space χ continuously embedded into the dual of the projective tensor space $B_1 \hat{\otimes}_\pi B_2$, called

²The two mentioned monographs use the term *real* instead of *scalar*; we changed it to avoid confusion with the quadratic variation of real processes.

³Given a Banach space B and a probability space (Ω, \mathbb{P}) a family of processes $\mathbb{X}^\epsilon: \Omega \times [0, T] \rightarrow B$ is said to converge in the ucp sense to $\mathbb{X}: \Omega \times [0, T] \rightarrow B$, when ϵ goes to zero, if $\lim_{\epsilon \rightarrow 0} \sup_{t \in [0, T]} |\mathbb{X}_t^\epsilon - \mathbb{X}_t|_B = 0$, in probability. Observe that we use the convergence in probability in the definition of covariation when the two processes are real and ucp convergence, when either \mathbb{X} or \mathbb{Y} is not one-dimensional. When $\mathbb{X} = \mathbb{Y}$ the two definitions are equivalent (see Lemma 2.1 of [37]).

Chi-subspace and to introduce the notion of χ -quadratic variation (and of χ -covariation, denoted by $[\cdot, \cdot]_\chi$), recalled in Section 3.1.

An interesting example of Banach space valued process with no tensor quadratic variation is the $C([- \tau, 0])$ -valued process \mathbb{X} defined as the frame (or window) of a standard Brownian motion (see [11]).

A second example, that indeed constitutes a main motivation for the present paper, is given by mild solutions of stochastic evolution equations in infinite dimensions: they have no scalar quadratic variation even if driven by a one-dimensional Brownian motion. We briefly recall their definition.

Consider a stochastic evolution equations of the form

$$\begin{cases} d\mathbb{X}(t) = (A\mathbb{X}(t) + b(t, \mathbb{X}(t))) dt + \sigma(t, \mathbb{X}(t)) d\mathbb{W}_Q(t) \\ \mathbb{X}(0) = x, \end{cases} \quad (2)$$

characterized by a generator of a C_0 -semigroup A , Lipschitz coefficients b and σ and a Q -Wiener process (with respect to some covariance operator Q) \mathbb{W} .

As described for example in Part III of [6], several families of partial differential equations with stochastic forcing terms or coefficients (SPDEs) can be reformulated as stochastic evolution equations and then can be studied in the general abstract setting; of course for any of them the specification of the generator A and of the functions b and σ are different.

There are some different possible way to define what we mean by *solution* of (2), among them there is the notion of *mild solution*. A progressively measurable process $\mathbb{X}(t)$ is a mild solutions of (2) (see [5] Chapter 7 or [19] Chapter 3) if it is the solutions of the following integral equation

$$\mathbb{X}(t) = e^{tA}x + \int_0^t e^{(t-r)A}b(r, \mathbb{X}(r)) dr + \int_0^t e^{(t-r)A}\sigma(r, \mathbb{X}(r)) d\mathbb{W}_Q(r).$$

This concept is widely used in the literature. Previous processes \mathbb{X} appear more generally in the form

$$\mathbb{X}(t) = e^{tA}x + \int_0^t e^{(t-r)A}b(r) dr + \int_0^t e^{(t-r)A}\sigma(r) d\mathbb{W}_Q(r).$$

Those processes will be called *convolution type processes* and will be the object of Section 4.

The contributions of the work

The novelty of the present paper arises both at the level of the stochastic calculus and of the infinite dimensional stochastic differential equations (and more in general convolution type processes).

The stochastic calculus part starts (Sections 2) with a natural extension (Definition 2.2) of the notion of forward integral in Banach spaces introduced in [8] and with the proof of its equivalence with the classical notion of integral when we integrate a predictable process with respect to a local martingale (Theorem 2.3). We also prove (Proposition 2.5) that, under suitable hypotheses, forward integrals also extend Young type integrals.

In Section 3, we extend the notion of Dirichlet process to infinite dimension. According to the literature, an Hilbert space-valued stochastic process can be naturally considered to be an *infinite dimensional Dirichlet process* if it is the sum of a local martingale and a zero energy process. A *zero energy process* (with some light sophistications) is a process such that the expectation of the quantity in Definition 1.2 converges to zero when ε goes to zero. This happens for instance in [7], even though that decomposition also appears in [26] Chapter VI Theorem 2.5, for processes associated with an infinite-dimensional Dirichlet form.

Extending Föllmer's notion of Dirichlet process to infinite dimension, a process \mathbb{X} taking values in a Hilbert space H , could be called *Dirichlet* if it is the sum of a local martingale \mathbb{M} plus a process \mathbb{A} having a zero scalar quadratic variation. However that natural notion is not suitable for an efficient stochastic calculus for infinite dimensional stochastic differential equations. Indeed solutions of SPDEs are in general no of such form, so this prevents to use Itô type formulae or generalized Doob-Meyer decompositions.

Using the notion of χ -finite quadratic variation process introduced in [10] we introduce the notion of χ -Dirichlet process as the sum of a local martingale \mathbb{M} and a process \mathbb{A} having a zero χ -quadratic variation.

A completely new notion in the present paper is the one of Hilbert valued ν -weak Dirichlet process which is again related to a Chi-subspace ν of the dual of the projective tensor product $H \hat{\otimes}_\pi H_1$ where H_1 is another Hilbert space, see Definition 3.14. It is of course an extension of the notion of real-valued weak Dirichlet process, see Definition 1.1. We illustrate that notion in the simple case when $H_1 = \mathbb{R}$, $\nu = \nu_0 \hat{\otimes}_\pi \mathbb{R} \equiv \nu_0$ and ν_0 is a Banach space continuously embedded in H^* : a process \mathbb{X} is called ν -weak Dirichlet process if it is the sum of a local martingale \mathbb{M} and a process \mathbb{A} such that $[\mathbb{A}, N]_\nu = 0$ for every real continuous local martingale N . This happens e.g. under the following assumptions.

- (i) There is a family $(R(\epsilon), \epsilon > 0)$ of non-negative random variables converging in probability, such that $Z(\epsilon) \leq R(\epsilon), \epsilon > 0$, where

$$Z(\epsilon) := \frac{1}{\epsilon} \int_0^T |\mathbb{A}(r + \epsilon) - \mathbb{A}(r)|_{\nu_0^*} |N(r + \epsilon) - N(r)| dr.$$

- (ii) For all $h \in \nu_0$, $\lim_{\epsilon \rightarrow 0^+} \frac{1}{\epsilon} \int_0^t \nu_0 \langle \mathbb{A}(r + \epsilon) - \mathbb{A}(r), h \rangle_{\nu_0^*} (N(r + \epsilon) - N(r)) dr = 0, \forall t \in [0, T]$.

The stochastic calculus theory developed in Sections 2 and 3, allows to prove, in Theorem 4.8, a general Itô's formula for convolution type processes and to show that they are χ -Dirichlet processes and a ν -weak-Dirichlet processes; this is done in Corollary 4.6. The most important result is however Theorem 3.19. It generalizes to the Hilbert values framework, Proposition 3.10 of [21] which states that given $f : [0, T] \times \mathbb{R} \rightarrow \mathbb{R}$ of class $C^{0,1}$ and X is a weak Dirichlet process with finite quadratic variation then $Y(t) = f(t, X(t))$ is a real weak Dirichlet process. Our result is a Fukushima decomposition in the spirit of Dirichlet forms, which is the natural extension of Doob-Meyer decomposition for semimartingales. It can also be seen as a substitution-tool of Itô's formula if f is not smooth. In particular, given some H -valued process \mathbb{X} , it allows to expand $f(t, \mathbb{X}(t))$ if \mathbb{X} is a ν -weak Dirichlet process which also has a χ -quadratic variation. In particular it fits the case when \mathbb{X} is convolution type process.

An important application consists in an application to stochastic control with state equation given by the solution of an infinite dimensional stochastic evolution equation. This is the object of [16] where, denoted by v a solution of the Hamilton-Jacobi-Bellman equation associated to the problem and by \mathbb{X} a mild solution of the state equation, the characterization of the process $t \mapsto v(t, \mathbb{X}(t))$ as a real weak Dirichlet process is used to prove a verification type theorem.

The scheme of the work is the following: in Section 2 we introduce the definition of forward integral with values in Banach spaces and we discuss the relation with the Da Prato-Zabczyk and Young integrals in the Hilbert framework. Section 3, devoted to stochastic calculus, is the core of the paper: we introduce the concepts of χ -Dirichlet processes, ν -weak-Dirichlet processes and we study their general properties. In Section 4, the developed theory is applied to the case of convolution type processes. In Appendix A we collect some useful results on projective tensor products of Hilbert spaces.

2 Stochastic integrals

2.1 Probability and stochastic processes

In the whole paper we denote by $(\Omega, \mathcal{F}, \mathbb{P})$ a complete probability space and by $\{\mathcal{F}_t\}_{t \geq 0}$ a filtration on $(\Omega, \mathcal{F}, \mathbb{P})$ satisfying the usual conditions. Given $\tilde{\Omega} \in \mathcal{F}$ we denote by $I_{\tilde{\Omega}} : \Omega \rightarrow \{0, 1\}$ the characteristic function of the set $\tilde{\Omega}$. Conformally to the Appendix, given two (real) Banach spaces B_1, B_2 $\mathcal{L}(B_1; B_2)$ will denote, as usual the space of linear bounded maps from B_1 to B_2 . Given a Banach space B we denote by $\mathcal{B}(B)$ the Borel σ -field on B . We fix $T > 0$.

By default we assume that all the processes $\mathbb{X} : [0, T] \times \Omega \rightarrow B$ are measurable functions with respect to the product σ -algebra $\mathcal{B}([0, T]) \otimes \mathcal{F}$ with values in $(B, \mathcal{B}(B))$. The dependence of a process on the

variable $\omega \in \Omega$ is emphasized only if needed by the context. When we say that a process is continuous (respectively left continuous, right continuous, càdlàg, càglàd ...) we mean that almost all its paths are continuous (respectively left-continuous, right-continuous, càdlàg, càglàd...).

Let \mathcal{G} be a sub- σ -field of $\mathcal{B}([0, T]) \otimes \mathcal{F}$. We say that such a process $\mathbb{X} : ([0, T] \times \Omega, \mathcal{G}) \rightarrow B$ is measurable with respect to \mathcal{G} if it is measurable in the usual sense. It is said *strongly (Bochner) measurable* (with respect to \mathcal{G}) if it is the limit of \mathcal{G} -measurable countable-valued functions. We recall that if \mathbb{X} is measurable and \mathbb{X} is càdlàg, càglàd or if B is separable then \mathbb{X} is strongly measurable. The σ -field \mathcal{G} will not be mentioned when it is clearly designated. We denote by \mathcal{P} the predictable σ -field on $[0, T] \times \Omega$. The processes \mathbb{X} measurable on $(\Omega \times [0, T], \mathcal{P})$ are also called *predictable* processes. All those processes will be considered as strongly measurable, with respect to \mathcal{P} . Each time we use expressions as “adapted”, “predictable” etc... we will always mean “with respect to the filtration $\{\mathcal{F}_t\}_{t \geq 0}$ ”.

The *blackboard bold* letters $\mathbb{X}, \mathbb{Y}, \mathbb{M} \dots$ are used for Banach (or Hilbert)-space valued processes, while notations X (or $Y, M \dots$) are reserved for real valued processes. We also adopt the notations introduced in Appendix A.

Notation 2.1. *We always assume the following convention: when needed all the Banach space càdlàg processes (or functions) indexed by $[0, T]$ are extended setting $\mathbb{X}(t) = \mathbb{X}(0)$ for $t \leq 0$ and $\mathbb{X}(t) = \mathbb{X}(T)$ for $t \geq T$.*

Definition 2.2. *Let B_1 and B_2 be two real separable Banach spaces. Let $\mathbb{X} : \Omega \times [0, T] \rightarrow \mathcal{L}(B_2, B_1)$ and $\mathbb{Y} : \Omega \times [0, T] \rightarrow B_2$ be two stochastic processes. Assume that \mathbb{Y} is continuous and \mathbb{X} is \mathbb{P} Bochner integrable a.s.*

If for almost every $t \in [0, T]$ the following limit (in the norm of the space B_1) exists in probability

$$\int_0^t \mathbb{X}(r) d^- \mathbb{Y}(r) := \lim_{\epsilon \rightarrow 0^+} \int_0^t \mathbb{X}(r) \left(\frac{\mathbb{Y}(r + \epsilon) - \mathbb{Y}(r)}{\epsilon} \right) dr \quad (3)$$

and the random function $t \mapsto \int_0^t \mathbb{X}(r) d^- \mathbb{Y}(r)$ admits a continuous (in B_1) version, we say that \mathbb{X} is forward integrable with respect to \mathbb{Y} . That version of $\int_0^t \mathbb{X}(r) d^- \mathbb{Y}(r)$ is called forward integral of \mathbb{X} with respect to \mathbb{Y} . Replacing $\int_0^t \mathbb{X}(r) d^- \mathbb{Y}(r)$ with $\int_0^t \mathbb{X}(r) d^+ \mathbb{Y}(r)$ and $\frac{\mathbb{Y}(r+\epsilon) - \mathbb{Y}(r)}{\epsilon}$ with $\frac{\mathbb{Y}(r) - \mathbb{Y}(r-\epsilon)}{\epsilon}$ in (3), $\int_0^t \mathbb{X}(r) d^+ \mathbb{Y}(r)$ is called backward integral of \mathbb{X} with respect to \mathbb{Y} .

The definition above is a natural generalization of that given in [8] Definition 3.4; there the forward integral is a real valued process.

2.2 The semimartingale case

Let H and U be two separable Hilbert spaces; we adopt the notations introduced in Appendix A. An U -valued measurable process $\mathbb{M} : [0, T] \times \Omega \rightarrow U$ is called martingale if, for all $t \in [0, T]$, \mathbb{M} is \mathcal{F}_t -adapted with $\mathbb{E}[|\mathbb{M}(t)|] < +\infty$ and $\mathbb{E}[\mathbb{M}(t_2) | \mathcal{F}_{t_1}] = \mathbb{M}(t_1)$ for all $0 \leq t_1 \leq t_2 \leq T$. The concept of (conditional) expectation for B -valued processes, for a separable Banach space B , is recalled for instance in [5] Section 1.3. All the considered martingales will be continuous.

We denote by $\mathcal{M}^2(0, T; H)$ the linear space of square integrable martingales equipped with the norm $|\mathbb{M}|_{\mathcal{M}^2(0, T; U)} := \left(\mathbb{E} \sup_{t \in [0, T]} |\mathbb{M}(t)|^2 \right)^{1/2}$. It is a Banach space as stated in [5], Proposition 3.9.

An U -valued measurable process $\mathbb{M} : [0, T] \times \Omega \rightarrow U$ is called local martingale if there exists a non-decreasing sequence of stopping times $\tau_n : \Omega \rightarrow [0, T] \cup \{+\infty\}$ such that $\mathbb{M}(t \wedge \tau_n)$ for $t \in [0, T]$ is a martingale and $\mathbb{P}[\lim_{n \rightarrow \infty} \tau_n = +\infty] = 1$. All the considered local martingales will be continuous.

Given a local martingale $\mathbb{M} : [0, T] \times \Omega \rightarrow U$, the process $|\mathbb{M}|^2$ is a real local sub-martingale, see Theorem 2.11 in [24]. The increasing predictable process, vanishing at zero, appearing in the Doob-Meyer decomposition of $|\mathbb{M}|^2$ is denoted by $[\mathbb{M}]^{\text{R}, cl}(t), t \in [0, T]$. It is of course uniquely determined and continuous.

We remind some properties of the Itô stochastic integral with respect to a local martingale \mathbb{M} . Call $\mathcal{I}_{\mathbb{M}}(0, T; H)$ the set of the processes $\mathbb{X} : [0, T] \times \Omega \rightarrow \mathcal{L}(U; H)$ that are strongly measurable from $([0, T] \times \Omega, \mathcal{P})$ to $\mathcal{L}(U; H)$ and such that

$$\|\mathbb{X}\|_{\mathcal{I}_{\mathbb{M}}(0,T;H)} := \left(\mathbb{E} \int_0^T \|\mathbb{X}(r)\|_{\mathcal{L}(U;H)}^2 d[\mathbb{M}]^{\mathbb{R},cl}(r) \right)^{1/2} < +\infty.$$

$\mathcal{I}_{\mathbb{M}}(0, T; H)$ endowed with the norm $\|\cdot\|_{\mathcal{I}_{\mathbb{M}}(0,T;H)}$ is a Banach space. The linear map

$$\begin{cases} I: \mathcal{I}_{\mathbb{M}}(0, T; H) \rightarrow \mathcal{M}^2(0, T; H) \\ \mathbb{X} \mapsto \int_0^\cdot \mathbb{X}(r) d\mathbb{M}(r) \end{cases}$$

is a contraction, see e.g. [27] Section 20.4 (above Theorem 20.5). As illustrated in [24] Section 2.2 (above Theorem 2.14), the stochastic integral with respect to \mathbb{M} extends to the integrands \mathbb{X} which are measurable from $([0, T] \times \Omega, \mathcal{P})$ to $\mathcal{L}(U; H)$ and such that

$$\int_0^T \|\mathbb{X}(r)\|_{\mathcal{L}(U;H)}^2 d[\mathbb{M}]^{\mathbb{R},cl}(r) < +\infty \quad a.s. \quad (4)$$

By $\mathcal{J}^2(0, T; U, H)$ we denote the family of integrands with respect to \mathbb{M} that satisfy (4).

Theorem 2.3. *Let us consider a continuous local martingale $\mathbb{M}: [0, T] \times \Omega \rightarrow U$ and a càglàd predictable $\mathcal{L}(U, H)$ -valued process \mathbb{X} satisfying (4). Then, the forward integral $\int_0^\cdot \mathbb{X}(r) d^-\mathbb{M}(r)$, defined in Definition 2.2 exists and coincides with the Itô integral $\int_0^\cdot \mathbb{X}(r) d\mathbb{M}(r)$.*

Proof of Theorem 2.3. We follow the arguments related to the finite-dimensional case, see Theorem 2 of [37]. Suppose first that $\mathbb{X} \in \mathcal{I}_{\mathbb{M}}(0, T; H)$. In this case it satisfies the hypotheses of the stochastic Fubini theorem in the form given in [25]. We have

$$\int_0^t \mathbb{X}(r) \frac{\mathbb{M}(r + \epsilon) - \mathbb{M}(r)}{\epsilon} dr = \int_0^t \mathbb{X}(r) \frac{1}{\epsilon} \left(\int_r^{r+\epsilon} d\mathbb{M}(\theta) \right) dr;$$

applying the stochastic Fubini Theorem, the expression above is equal to

$$\int_0^t \left(\frac{1}{\epsilon} \int_{\theta-\epsilon}^{\theta} \mathbb{X}(r) dr \right) d\mathbb{M}(\theta) + R_\epsilon(t)$$

where $R_\epsilon(t)$ is a boundary term that converges to 0 in probability, for any $t \in [0, T]$, so that we can ignore it. We can apply now the maximal inequality stated in [39], Theorem 1: there exists a universal constant $C > 0$ such that, for every $f \in L^2([0, t]; \mathbb{R})$,

$$\int_0^t \left(\sup_{\epsilon \in (0,1]} \left| \frac{1}{\epsilon} \int_{(r-\epsilon)}^r f(\xi) d\xi \right| \right)^2 dr \leq C \int_0^t f^2(r) dr. \quad (5)$$

According to the vector valued version of the Lebesgue differentiation Theorem (see Theorem II.2.9 in [12]), the quantity $\frac{1}{\epsilon} \int_{(r-\epsilon)}^r \mathbb{X}(\xi) d\xi$ converges $d\mathbb{P} \otimes dr$ a.e. to $\mathbb{X}(r)$. Consequently (5) and dominated convergence theorem imply

$$\mathbb{E} \int_0^t \left\| \left(\frac{1}{\epsilon} \int_{\theta-\epsilon}^{\theta} \mathbb{X}(r) dr \right) - \mathbb{X}(\xi) \right\|_{\mathcal{L}(U,H)}^2 d[\mathbb{M}]^{\mathbb{R},cl}(\xi) \xrightarrow{\epsilon \rightarrow 0} 0, \quad (6)$$

Since the integral is a contraction from $\mathcal{I}_{\mathbb{M}}(0, T; H)$ to $\mathcal{M}^2(0, T; H)$, this show the claim for the processes in $\mathcal{I}_{\mathbb{M}}(0, T; H)$. If \mathbb{X} is a càglàd adapted process (and then a.s. bounded and therefore in $\mathcal{J}^2(0, T; U, H)$) we use the same argument after localizing the problem by using the suitable sequence of stopping times defined by $\tau_n := \inf \left\{ t \in [0, T] : \int_0^t \|\mathbb{X}(r)\|_{\mathcal{L}(U,H)}^2 d[\mathbb{M}]^{\mathbb{R},cl}(r) \geq n \right\}$ (and $+\infty$ if the set is void). \square

An easier but still important statement concerns the integration with respect bounded variation processes.

Proposition 2.4. *Let us consider a continuous bounded variation process $\mathbb{V}: [0, T] \times \Omega \rightarrow U$ and let \mathbb{X} be a càglàd measurable process $[0, T] \times \Omega \rightarrow \mathcal{L}(U, H)$. Then the forward integral $\int_0^\cdot \mathbb{X}(r) d^-\mathbb{V}(r)$, defined in Definition 2.2 exists and coincides with the Lebesgue-Bochner integral $\int_0^\cdot \mathbb{X}(r) d\mathbb{V}(r)$.*

Proof. The proof is similar to the one of Theorem 2.3. \square

As for the case of finite dimensional integrators, forward integrals also extend Young type integrals. Let $\mathbb{Y}: [0, T] \times \Omega \rightarrow U$ be a γ -Hölder continuous process and \mathbb{X} be an $\mathcal{L}(U; H)$ -valued α -Hölder continuous with $\alpha + \gamma > 1$. Then, the so called *Young integral* $\int_0^t \mathbb{X} d^y \mathbb{Y}$ is well-defined, see Proposition [22], similarly as in the one-dimensional case. Moreover the H -valued integral process has also γ -Hölder continuous paths.

An example of process Y can arise as follows. Let (e_n) be an orthonormal basis in U , (α_n) be a sequence of real numbers such that $\sum_n \alpha_n^2 < \infty$. Let β^n be sequence of real valued fractional Brownian motions of Hurst index $H > \gamma$. It is not difficult to show that the U -valued random function $\mathbb{Y}_t = \sum_{n=1} \alpha_n \beta_n e_n$ is well-defined. In particular for $0 \leq t$, and $\ell \in \mathbb{N}$ we get

$$E(|Y(t) - \mathbb{Y}(s)|^{2\ell}) \leq \sum_{n=1} \alpha_n^2 (t - s)^{2\ell H}.$$

By Kolmogorov-Centsov theorem, taking ℓ large enough, it is possible to show that \mathbb{Y} has a γ -Hölder continuous version.

Proposition 2.5. *Let $\alpha, \gamma, \mathbb{Y}, \mathbb{X}$ as above. Then the forward integral $\int_0^\cdot \mathbb{X} d^-\mathbb{Y}$ (resp. $\int_0^\cdot \mathbb{X} d^+\mathbb{Y}$) exists and equals $\int_0^\cdot \mathbb{X} d^y \mathbb{Y}$.*

Proof. The proof is very similar to the one in Section 2.2 of [37] concerning the one-dimensional case, by making use of Bochner integrals. \square

A very natural research line is to study possible extensions of the results of this section to study the equivalence of the forward integral and the stochastic integral in UMD spaces introduced in [40]. Similarly extensions of the results of Section 4 could include the study of the applicability of concepts introduced in Section 3 to convolution type processes in Banach space and, notably, to mild solution of stochastic evolution equations in UMD spaces as developed e.g. in [41]. Some relevant results in that direction have been done by [33]. Relevant applications to optimal control in the framework of UMD spaces were obtained e.g. in [2].

3 χ -quadratic variation and χ -Dirichlet processes

3.1 χ -quadratic variation processes

We denote by $\mathcal{C}([0, T])$ the space of the real continuous processes equipped with the ucp (uniform convergence in probability) topology. Consider two real separable Banach spaces B_1 and B_2 with the same notations as in Appendix A.

Following [8, 11] a *Chi-subspace* (of $(B_1 \hat{\otimes}_\pi B_2)^*$) is defined as any Banach subspace $(\chi, |\cdot|_\chi)$ which is continuously embedded into $(B_1 \hat{\otimes}_\pi B_2)^*$: in other words, there is some constant C such that $|\cdot|_{(B_1 \hat{\otimes}_\pi B_2)^*} \leq C |\cdot|_\chi$.

Concrete examples of Chi-subspaces are provided e.g. in [11], see Example 3.4. For example, for a fixed positive number τ , in the case $B_1 = B_2 = \mathcal{C}([-\tau, 0])$, the space of finite signed Borel measures on $[-\tau, 0]^2$, $\mathcal{M}([-\tau, 0]^2)$ (equipped with the total variation norm) is shown to be a Chi-subspace and it is used, together with some specific subspaces, to prove Ito-type formulas and Fukushima-type decompositions. Another concrete example will be used in Section 4 where, given $B_1 = B_2 = H$ for some

separable Hilbert H space and denoted by A the generator of a C_0 -semigroup on H , we use the space $\bar{\chi} := D(A^*) \hat{\otimes}_\pi D(A^*)$ (see (39)). Among the several possible examples (see e.g. [6]) we have for instance, in the case $H = L^2(\mathcal{O})$ for some bounded regular domain $\mathcal{O} \subseteq \mathbb{R}^m$, the heat semigroup with Dirichlet condition, in this case we have and $D(A^*) = D(A) = H_0^1(\mathcal{O}) \cap H^2(\mathcal{O})$.

Let χ be a generic Chi-subspace. We introduce the following definition.

Definition 3.1. *Given \mathbb{X} (respectively \mathbb{Y}) a B_1 -valued (respectively B_2 -valued) process, we say that (\mathbb{X}, \mathbb{Y}) admits a χ -covariation if the two following conditions are satisfied.*

H1 *For any sequence of positive real numbers $\epsilon_n \searrow 0$ there exists a subsequence ϵ_{n_k} such that*

$$\sup_k \int_0^T \frac{|(J(\mathbb{X}(r + \epsilon_{n_k}) - \mathbb{X}(r)) \otimes (\mathbb{Y}(r + \epsilon_{n_k}) - \mathbb{Y}(r)))|_{\chi^*}}{\epsilon_{n_k}} dr < \infty \text{ a.s.}, \quad (7)$$

where $J : B_1 \hat{\otimes}_\pi B_2 \rightarrow (B_1 \hat{\otimes}_\pi B_2)^{**}$ is the canonical injection between a space and its bidual.

H2 *If we denote by $[\mathbb{X}, \mathbb{Y}]_\chi^\epsilon$ the application*

$$\begin{cases} [\mathbb{X}, \mathbb{Y}]_\chi^\epsilon : \chi \rightarrow \mathcal{C}([0, T]) \\ \phi \mapsto \int_0^\cdot \left\langle \phi, \frac{J((\mathbb{X}(r + \epsilon) - \mathbb{X}(r)) \otimes (\mathbb{Y}(r + \epsilon) - \mathbb{Y}(r)))}{\epsilon} \right\rangle_{\chi^*} dr, \end{cases} \quad (8)$$

the following two properties hold.

(i) *There exists an application, denoted by $[\mathbb{X}, \mathbb{Y}]_\chi$, defined on χ with values in $\mathcal{C}([0, T])$, satisfying*

$$[\mathbb{X}, \mathbb{Y}]_\chi^\epsilon(\phi) \xrightarrow[\epsilon \rightarrow 0_+]{ucp} [\mathbb{X}, \mathbb{Y}]_\chi(\phi), \quad (9)$$

for every $\phi \in \chi \subset (B_1 \hat{\otimes}_\pi B_2)^*$.

(ii) *There exists a strongly measurable process $\widetilde{[\mathbb{X}, \mathbb{Y}]_\chi} : \Omega \times [0, T] \rightarrow \chi^*$, such that*

- *for almost all $\omega \in \Omega$, $\widetilde{[\mathbb{X}, \mathbb{Y}]_\chi}(\omega, \cdot)$ is a (càdlàg) bounded variation process,*
- *$\widetilde{[\mathbb{X}, \mathbb{Y}]_\chi}(\cdot, t)(\phi) = [\mathbb{X}, \mathbb{Y}]_\chi(\phi)(\cdot, t)$ a.s. for all $\phi \in \chi$, $t \in [0, T]$.*

Lemma 3.2. *In the setting of Definition 3.1, we set*

$$A(\epsilon) := \int_0^T \frac{|(J(\mathbb{X}(r + \epsilon) - \mathbb{X}(r)) \otimes (\mathbb{Y}(r + \epsilon) - \mathbb{Y}(r)))|_{\chi^*}}{\epsilon} dr. \quad (10)$$

(i) *If $\lim_{\epsilon \rightarrow 0} A(\epsilon)$ exists in probability then Condition **H1** of Definition 3.1 is verified.*

(ii) *If $\lim_{\epsilon \rightarrow 0} A(\epsilon) = 0$ in probability then (\mathbb{X}, \mathbb{Y}) admits a χ -covariation and $\widetilde{[\mathbb{X}, \mathbb{Y}]_\chi}$ vanishes.*

Proof. It is an easy consequence of Remark 3.10 and Lemma 3.18 in [11]. □

If (\mathbb{X}, \mathbb{Y}) admits a χ -covariation we call χ -covariation of (\mathbb{X}, \mathbb{Y}) the χ^* -valued process $\widetilde{[\mathbb{X}, \mathbb{Y}]_\chi}$ defined for every $\omega \in \Omega$ and $t \in [0, T]$ by $\phi \mapsto \widetilde{[\mathbb{X}, \mathbb{Y}]_\chi}(\omega, t)(\phi) = [\mathbb{X}, \mathbb{Y}]_\chi(\phi)(\omega, t)$. By abuse of notation, $[\mathbb{X}, \mathbb{Y}]_\chi$ will also be often called χ -covariation and it will be confused with $\widetilde{[\mathbb{X}, \mathbb{Y}]_\chi}$. We say that a process \mathbb{X} admits a χ -quadratic variation if (\mathbb{X}, \mathbb{X}) admits a χ -covariation. The process $\widetilde{[\mathbb{X}, \mathbb{X}]_\chi}$, often denoted by $\widetilde{[\mathbb{X}]_\chi}$, is also called χ -quadratic variation of \mathbb{X} .

Definition 3.3. If $\chi = (B_1 \hat{\otimes}_\pi B_2)^*$ the χ -covariation is called global covariation. In this case we omit the index $(B_1 \hat{\otimes}_\pi B_2)^*$ using the notations $[\mathbb{X}, \mathbb{Y}]$ and $\widetilde{[\mathbb{X}, \mathbb{Y}]}$.

Lemma 3.4. Let \mathbb{X} and \mathbb{Y} as in Definition 3.1. The properties below hold.

- (i) If (\mathbb{X}, \mathbb{Y}) admits a global covariation then it admits a χ -covariation for any Chi-subspace χ . Moreover $[\mathbb{X}, \mathbb{Y}]_\chi(\phi) = [\mathbb{X}, \mathbb{Y}](\phi)$ for all $\phi \in \chi$.
- (ii) Suppose that \mathbb{X} and \mathbb{Y} admit a scalar quadratic variation (Definitions 1.2) and (\mathbb{X}, \mathbb{Y}) has a tensor covariation (Definition 1.3), denoted by $[\mathbb{X}, \mathbb{Y}]^\otimes$. Then (\mathbb{X}, \mathbb{Y}) admits a global covariation $[\mathbb{X}, \mathbb{Y}]$. In particular, recalling that $B_1 \hat{\otimes}_\pi B_2$ is embedded in $(B_1 \hat{\otimes}_\pi B_2)^{**}$, we have $\widetilde{[\mathbb{X}, \mathbb{Y}]} = [\mathbb{X}, \mathbb{Y}]^\otimes$.

Proof. Part (i) follows by the definitions. For part (ii) the proof is a slight adaptation of the one of Proposition 3.14 in [11]. In particular condition **H1** holds using Cauchy-Schwarz inequality. \square

The product of a real finite quadratic variation process and a zero real quadratic variation process is again a zero quadratic variation processes. Under some conditions this can be generalized to the infinite dimensional case as shown in the following proposition.

Proposition 3.5. Let $i = 1, 2$ and ν_i be a Banach space continuously embedded in the dual B_i^* of a real separable Banach space B_i . Let consider the Chi-subspace of the type $\chi_1 = \nu_1 \hat{\otimes}_\pi B_2^*$ and $\chi_2 = B_1^* \hat{\otimes}_\pi \nu_2$, $\hat{\chi}_i = \nu_i \hat{\otimes}_\pi \nu_i$, $i = 1, 2$. Let \mathbb{X} (respectively \mathbb{Y}) be a process with values in B_1 (respectively B_2).

- (i) Suppose that \mathbb{X} admits a $\hat{\chi}_1$ -quadratic variation and \mathbb{Y} a zero scalar quadratic variation. Then $[\mathbb{X}, \mathbb{Y}]_{\chi_1} = 0$.
- (ii) Similarly suppose that \mathbb{Y} admits a $\hat{\chi}_2$ -quadratic variation and \mathbb{X} a zero scalar quadratic variation. Then $[\mathbb{X}, \mathbb{Y}]_{\chi_2} = 0$.

Proof. We remark that Lemma A.5 implies that χ_i and $\hat{\chi}_i$, $i = 1, 2$ are indeed Chi-subspaces. By Lemma 3.2(ii), it is enough to show that $A(\varepsilon)$ defined in (10) converge to zero, with $\chi = \chi_i$, $i = 1, 2$. By symmetry it is enough to show (i).

The Banach space B_i it isometrically embedded in its bidual B_i^{**} , $i = 1, 2$, so, since $\nu_1 \subseteq B_1^*$ with continuous inclusion, we have $B_1 \subseteq B_1^{**} \subseteq \nu_1^*$ where the inclusion are continuous.

Moreover, since $\chi_1 = \nu_1 \hat{\otimes}_\pi B_2^* \subseteq B_1^* \hat{\otimes}_\pi B_2^* \subseteq (B_1 \hat{\otimes}_\pi B_2)^*$, with continuous inclusions, taking into account Lemma A.5, we have $J(B_1 \hat{\otimes}_\pi B_2) \subseteq (B_1 \hat{\otimes}_\pi B_2)^{**} \subseteq \chi_1^*$. Let $a \in B_1$ and $b \in B_2$. We have

$$\begin{aligned} |J(a \otimes b)|_{\chi_1^*} &= \sup_{|\varphi|_{\nu_1} \leq 1, |\psi|_{B_2^*} \leq 1} |\chi_1^* \langle J(a \otimes b), \varphi \otimes \psi \rangle_{\chi_1}| \\ &= \sup_{|\varphi|_{\nu_1} \leq 1} |\nu_1 \langle \varphi, a \rangle_{\nu_1^*}| \sup_{|\psi|_{B_2^*} \leq 1} |B_2^* \langle \psi, b \rangle_{B_2^{**}}| = |a|_{\nu_1^*} |b|_{B_2^{**}} = |a|_{\nu_1^*} |b|_{B_2}. \end{aligned} \quad (11)$$

Consequently, with $a = \mathbb{X}(r + \varepsilon) - \mathbb{X}(r)$ and $b = \mathbb{Y}(r + \varepsilon) - \mathbb{Y}(r)$ for $r \in [0, T]$, we have

$$\begin{aligned} A(\varepsilon) &= \int_0^T \frac{|(J(\mathbb{X}(r + \varepsilon) - \mathbb{X}(r)) \otimes (\mathbb{Y}(r + \varepsilon) - \mathbb{Y}(r)))|_{\chi_1^*}}{\varepsilon} dr = \int_0^T |\mathbb{X}(r + \varepsilon) - \mathbb{X}(r)|_{\nu_1^*} |\mathbb{Y}(r + \varepsilon) - \mathbb{Y}(r)|_{B_2} \frac{dr}{\varepsilon} \\ &\leq \left(\int_0^T |\mathbb{X}(r + \varepsilon) - \mathbb{X}(r)|_{\nu_1^*}^2 \frac{dr}{\varepsilon} \int_0^T |\mathbb{Y}(r + \varepsilon) - \mathbb{Y}(r)|_{B_2}^2 \frac{dr}{\varepsilon} \right)^{1/2} \\ &= \left(\int_0^T \frac{|(J(\mathbb{X}(r + \varepsilon) - \mathbb{X}(r)) \otimes (\mathbb{X}(r + \varepsilon) - \mathbb{X}(r)))|_{\hat{\chi}_1^*}}{\varepsilon} dr \right)^{1/2} \left(\int_0^T \frac{|\mathbb{Y}(r + \varepsilon) - \mathbb{Y}(r)|_{B_2}^2}{\varepsilon} dr \right)^{1/2}. \end{aligned} \quad (12)$$

The last equality is obtained using an argument similar to (11). The condition **H1** related to the $\hat{\chi}_1$ -quadratic variation of \mathbb{X} and the zero scalar quadratic variation of \mathbb{Y} , imply that previous expression converges to zero. \square

3.2 Tensor covariation and classical tensor covariation

The notions of tensor covariation recalled in Definition 1.3, denoted by $[\cdot, \cdot]^\otimes$, concerns Banach space-valued processes. In the specific case when H_1 and H_2 are two separable Hilbert spaces and $\mathbb{M}: [0, T] \times \Omega \rightarrow H_1$, $\mathbb{N}: [0, T] \times \Omega \rightarrow H_2$ are two continuous local martingales, another (classical) notion of tensor covariation is defined, see for instance in Section 23.1 of [27]. This is denoted by $[\mathbb{M}, \mathbb{N}]^{cl}$ and (see Chapters 22 and 23 of [27]) it is an $(H_1 \hat{\otimes}_\pi H_2)$ -valued process. Recall that $(H_1 \hat{\otimes}_\pi H_2) \subseteq (H_1 \hat{\otimes}_\pi H_2)^{**}$.

Remark 3.6. *We observe the following facts.*

- (i) *Taking into account Lemma A.2 we know that, given $h \in H_1$ and $k \in H_2$, $h^* \otimes k^*$ can be considered as an element of $(H_1 \hat{\otimes}_\pi H_2)^*$. One has*

$$[\mathbb{M}, \mathbb{N}]^{cl}(t)(h^* \otimes k^*) = [\langle \mathbb{M}, h \rangle, \langle \mathbb{N}, k \rangle](t), \quad (13)$$

where h^* (respectively k^*) is associated with h (respectively k) via Riesz theorem. This property characterizes $[\mathbb{M}, \mathbb{N}]^{cl}$, see e.g. [5], Section 3.4 after Proposition 3.11.

- (iii) *If $H_2 = \mathbb{R}$ and $\mathbb{N} = N$ is a real continuous local martingale then, identifying $H_1 \hat{\otimes}_\pi H_2$ with H_1 , $[\mathbb{M}, \mathbb{N}]^{cl}$ can be considered as an H_1 -valued process. The characterization (13) can be translated into*

$$[\mathbb{M}, \mathbb{N}]^{cl}(t)(h^*) = [\langle \mathbb{M}, h \rangle, N](t), \forall h \in H_1. \quad (14)$$

By inspection, this allows us to see that the classical covariation between \mathbb{M} and N can be expressed as

$$[\mathbb{M}, N]^{cl}(t) := \mathbb{M}(t)N(t) - \mathbb{M}(0)N(0) - \int_0^t N(r) d\mathbb{M}(r) - \int_0^t \mathbb{M}(r) dN(r). \quad (15)$$

In the sequel H will denote a real separable Hilbert space.

Lemma 3.7. *Any continuous local martingale with values in H has a scalar quadratic variation and a tensor quadratic variation. If \mathbb{M}_1 and \mathbb{M}_2 are continuous local martingales with values respectively in H_1 and H_2 then $(\mathbb{M}_1, \mathbb{M}_2)$ admits a tensor covariation.*

Proof. See Proposition 1.7 and Proposition 1.6 in [11]. □

Lemma 3.8. *Let \mathbb{M} (respectively \mathbb{N}) be a continuous local martingale with values in H . Then (\mathbb{M}, \mathbb{N}) admits a tensor covariation and*

$$[\mathbb{M}, \mathbb{N}]^\otimes = [\mathbb{M}, \mathbb{N}]^{cl}. \quad (16)$$

In particular (\mathbb{M}, \mathbb{N}) admits a global covariation and

$$\widetilde{[\mathbb{M}, \mathbb{N}]} = [\mathbb{M}, \mathbb{N}]^{cl}. \quad (17)$$

Proof. Thanks to Lemma 3.7, \mathbb{M} and \mathbb{N} admit a scalar quadratic variation and (\mathbb{M}, \mathbb{N}) a tensor covariation. By Lemma 3.4(ii) they admit a global covariation. It is enough to show that they are equal as elements of $(H_1 \hat{\otimes}_\pi H_2)^{**}$, so one needs to prove that

$$[\mathbb{M}, \mathbb{N}]^\otimes(\phi) = [\mathbb{M}, \mathbb{N}]^{cl}(\phi), \quad (18)$$

for every $\phi \in (H_1 \hat{\otimes}_\pi H_2)^*$.

Given $h \in H_1$ and $k \in H_2$, we consider (via Lemma A.2) $h^* \otimes k^*$ as an element of $(H_1 \hat{\otimes}_\pi H_2)^*$. According to Lemma A.6, $H_1^* \hat{\otimes}_\pi H_2^*$ is sequentially dense in $(H_1 \hat{\otimes}_\pi H_2)^*$ in the weak-* topology. Therefore, taking into account item (i) of Remark 3.6 we only need to show that

$$[\mathbb{M}, \mathbb{N}]^\otimes(h^* \otimes k^*) = [\langle \mathbb{M}, h \rangle, \langle \mathbb{N}, k \rangle], \quad (19)$$

for every $h \in H_1, k \in H_2$. By the usual properties of Bochner integral the left-hand side of (19) is the limit of

$$\begin{aligned} & \frac{1}{\epsilon} \int_0^\cdot (M(r+\epsilon) - M(r)) \otimes (N(r+\epsilon) - N(r))(h^* \otimes k^*) dr \\ &= \frac{1}{\epsilon} \int_0^\cdot \langle (M(r+\epsilon) - M(r)), h \rangle \langle (N(r+\epsilon) - N(r)), k \rangle dr. \end{aligned} \quad (20)$$

Since $\langle \mathbb{M}, h \rangle$ and $\langle \mathbb{N}, k \rangle$ are real local martingales, the covariation $[\langle \mathbb{M}, h \rangle, \langle \mathbb{N}, k \rangle]$ exists and equals the classical covariation of local martingales because of Proposition 2.4(3) of [36]. \square

Lemma 3.9. *Let $\mathbb{M}: [0, T] \times \Omega \rightarrow H$ be a continuous local martingale and \mathbb{Z} a measurable process from $([0, T] \times \Omega, \mathcal{P})$ to H^* and such that $\int_0^T \|\mathbb{Z}(r)\|^2 d[\mathbb{M}]^{\mathbb{R}, cl}(r) < +\infty$ a.s. We define*

$$X(t) := \int_0^t \langle \mathbb{Z}(r), d\mathbb{M}(r) \rangle. \quad (21)$$

Then X is a real continuous local martingale and, for every continuous real local martingale N , the (classical, one-dimensional) covariation process $[X, N]$ is given by

$$[X, N](t) = \int_0^t \langle \mathbb{Z}(r), d[\mathbb{M}, N]^{cl}(r) \rangle; \quad (22)$$

in particular the integral in the right-side is well-defined.

Proof. The fact that X is a local martingale is part of the result of Theorem 2.14 in [24]. For the other claim we can reduce, using a sequence of suitable stopping times as in the proof of Theorem 2.3, to the case in which \mathbb{Z} , \mathbb{M} and N are square integrable martingales. Taking into account the characterization (14) and the discussion developed in [29], page 456, (22) follows. \square

When one of the processes is real the formalism of global covariation can be simplified as shown in the following proposition.

Proposition 3.10. *Let be $\mathbb{X}: [0, T] \times \Omega \rightarrow H$ a Bochner integrable process and $Y: [0, T] \times \Omega \rightarrow \mathbb{R}$ a real valued process. Suppose the following.*

- (a) *For any ϵ , $\frac{1}{\epsilon} \int_0^T |\mathbb{X}(r+\epsilon) - \mathbb{X}(r)| |Y(r+\epsilon) - Y(r)| dr$ is bounded by a r.v. $A(\epsilon)$ such that $A(\epsilon)$ converges in probability when $\epsilon \rightarrow 0$.*
- (b) *For every $h \in H$ the limit*

$$C(t)(h) := \lim_{\epsilon \rightarrow 0^+} \frac{1}{\epsilon} \int_0^t \langle h, \mathbb{X}(r+\epsilon) - \mathbb{X}(r) \rangle (Y(r+\epsilon) - Y(r)) dr$$

exists ucp and there exists a continuous process $\tilde{C}: [0, T] \times \Omega \rightarrow H$ such that $\langle \tilde{C}(t, \omega), h \rangle = C(t)(h)(\omega)$ for \mathbb{P} -a.s. $\omega \in \Omega$, for all $t \in [0, T]$ and $h \in H$.

Then \mathbb{X} and Y admit a global covariation and, after identifying H with $(H \hat{\otimes}_\pi \mathbb{R})^$, $\tilde{C} = \widetilde{[\mathbb{X}, Y]}$.*

Proof. Taking into account the identification of H with $(H \hat{\otimes}_\pi \mathbb{R})^*$ the result is a consequence of Corollary 3.26 of [11]. \square

Proposition 3.11. *If $\mathbb{M}: [0, T] \times \Omega \rightarrow H$ and $N: [0, T] \times \Omega \rightarrow \mathbb{R}$ are continuous local martingales. Then \mathbb{M} and N admit a global covariation and $\widetilde{[\mathbb{M}, N]} = [\mathbb{M}, N]^{cl}$.*

Proof. We have to check the conditions stated in Proposition 3.10 for \tilde{C} equal to the right side of (15). Concerning (a), by Cauchy-Schwarz inequality we have

$$\frac{1}{\epsilon} \int_0^T |N(r+\epsilon) - N(r)| |\mathbb{M}(r+\epsilon) - \mathbb{M}(r)| dr \leq [N, N]^{\epsilon, \mathbb{R}}(T) [\mathbb{M}, \mathbb{M}]^{\epsilon, \mathbb{R}}(T).$$

Since both N and \mathbb{M} are local martingales they admit a scalar quadratic variation (as recalled in Lemma 3.7), the result is established. Concerning (b), taking into account (14), we need to prove that for any $h \in H$

$$\lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} \int_0^T (M^h(r+\epsilon) - M^h(r))(N(r+\epsilon) - N(r)) dr = [\langle M, h \rangle, N] \quad (23)$$

ucp, where M^h is the real local martingale $\langle \mathbb{M}, h \rangle$. (23) follows by Proposition 2.4(3) of [36]. \square

3.3 χ -Dirichlet and ν -weak Dirichlet processes

We have now at our disposal all the elements we need to introduce the concepts of χ -Dirichlet process and ν -weak Dirichlet process.

Definition 3.12. Let $\chi \subseteq (H \hat{\otimes}_\pi H)^*$ be a Chi-subspace. A continuous H -valued process $\mathbb{X}: ([0, T] \times \Omega, \mathcal{P}) \rightarrow H$ is called χ -Dirichlet process if there exists a decomposition $\mathbb{X} = \mathbb{M} + \mathbb{A}$ where

- (i) \mathbb{M} is a continuous local martingale,
- (ii) \mathbb{A} is a continuous χ -zero quadratic variation process with $\mathbb{A}(0) = 0$.

Definition 3.13. Let H and H_1 be two separable Hilbert spaces. Let $\nu \subseteq (H \hat{\otimes}_\pi H_1)^*$ be a Chi-subspace. A continuous adapted H -valued process $\mathbb{A}: [0, T] \times \Omega \rightarrow H$ is said to be \mathcal{F}_t - ν -martingale-orthogonal if $[\mathbb{A}, \mathbb{N}]_\nu = 0$, for any H_1 -valued continuous local martingale \mathbb{N} .

As we have done for the expressions “stopping time”, “adapted”, “predictable”... since we always use the filtration \mathcal{F}_t , we simply write ν -martingale-orthogonal instead of \mathcal{F}_t - ν -martingale-orthogonal.

Definition 3.14. Let H and H_1 be two separable Hilbert spaces. Let $\nu \subseteq (H \hat{\otimes}_\pi H_1)^*$ be a Chi-subspace. A continuous H -valued process $\mathbb{X}: [0, T] \times \Omega \rightarrow H$ is called ν -weak-Dirichlet process if it is adapted and there exists a decomposition $\mathbb{X} = \mathbb{M} + \mathbb{A}$ where

- (i) \mathbb{M} is an H -valued continuous local martingale,
- (ii) \mathbb{A} is an ν -martingale-orthogonal process with $\mathbb{A}(0) = 0$.

The decomposition of a real weak Dirichlet process is unique, see Remark 3.5 of [21]. For the infinite dimensional case we have the following result.

Proposition 3.15. Let $\nu \subseteq (H \hat{\otimes}_\pi H)^*$ be a Chi-subspace. Suppose that ν is dense in $(H \hat{\otimes}_\pi H)^*$. Then any decomposition of a ν -weak-Dirichlet process \mathbb{X} is unique.

Proof. Assume that $\mathbb{X} = \mathbb{M}^1 + \mathbb{A}^1 = \mathbb{M}^2 + \mathbb{A}^2$ are two decompositions where \mathbb{M}^1 and \mathbb{M}^2 are continuous local martingales and $\mathbb{A}^1, \mathbb{A}^2$ are ν -martingale-orthogonal processes. If we call $\mathbb{M} := \mathbb{M}^1 - \mathbb{M}^2$ and $\mathbb{A} := \mathbb{A}^1 - \mathbb{A}^2$ we have $0 = \mathbb{M} + \mathbb{A}$.

By Lemma 3.8, \mathbb{M} has a global quadratic variation. In particular it also has a ν -quadratic variation and, thanks to the bilinearity of the ν -covariation,

$$0 = [\mathbb{M}, 0]_\nu = [\mathbb{M}, \mathbb{M} + \mathbb{A}]_\nu = [\mathbb{M}, \mathbb{M}]_\nu + [\mathbb{M}, \mathbb{A}]_\nu = [\mathbb{M}, \mathbb{M}]_\nu + 0 = [\mathbb{M}, \mathbb{M}]_\nu.$$

We prove now that \mathbb{M} has also zero global quadratic variation. We have denoted by $\mathcal{C}([0, T])$ the space of the real continuous processes defined on $[0, T]$. We introduce, for $\epsilon > 0$, the operators

$$\begin{cases} [\mathbb{M}, \mathbb{M}]^\epsilon: (H \hat{\otimes}_\pi H)^* \rightarrow \mathcal{C}([0, T]) \\ ([\mathbb{M}, \mathbb{M}]^\epsilon(\phi))(t) := \frac{1}{\epsilon} \int_0^t \langle (\mathbb{M}_{r+\epsilon} - \mathbb{M}_r)^{\otimes 2}, \phi \rangle_{(H \hat{\otimes}_\pi H)^*} dr. \end{cases} \quad (24)$$

Observe the following.

- (a) $[\mathbb{M}, \mathbb{M}]^\epsilon$ are linear and bounded operators.
- (b) For $\phi \in (H \hat{\otimes}_\pi H)^*$ the limit $[\mathbb{M}, \mathbb{M}](\phi) := \lim_{\epsilon \rightarrow 0} [\mathbb{M}, \mathbb{M}]^\epsilon(\phi)$ exists.
- (c) If $\phi \in \nu$ we have $[\mathbb{M}, \mathbb{M}](\phi) = 0$.

Thanks to (a) and (b) and Banach-Steinhaus theorem for F -spaces (see Theorem 17, Chapter II in [14]) we know that $[\mathbb{M}, \mathbb{M}]$ is linear and bounded. Thanks to (c) and the fact that the inclusion $\nu \subseteq (H \hat{\otimes}_\pi H)^*$ is dense, it follows $[\mathbb{M}, \mathbb{M}] = 0$. By Lemma 3.8 $[\mathbb{M}, \mathbb{M}]$ coincides with the classical quadratic variation $[\mathbb{M}, \mathbb{M}]^{cl}$ and it is characterized by

$$0 = [\mathbb{M}, \mathbb{M}]^{cl}(h^*, k^*) = [\langle \mathbb{M}, h \rangle, \langle \mathbb{M}, k \rangle],$$

by Remark 3.6(i). Since $\mathbb{M}(0) = 0$ and therefore $\langle \mathbb{M}, h \rangle(0) = 0$ it follows that $\langle \mathbb{M}, h \rangle \equiv 0$ for any $h \in H$. Finally $\mathbb{M} \equiv 0$, which concludes the proof. \square

Proposition 3.16. *Let H and H_1 be two separable Hilbert spaces. Let $\chi = \chi_0 \hat{\otimes}_\pi \chi_0$ for some χ_0 Banach space continuously embedded in H^* . Define $\nu = \chi_0 \hat{\otimes}_\pi H_1^*$. Then an H -valued continuous zero χ -quadratic variation process \mathbb{A} is a ν -martingale-orthogonal process.*

Proof. Taking into account Lemma A.5, χ is a Chi-subspace of $(H \hat{\otimes}_\pi H)^*$ and ν is a Chi-subspace of $(H \hat{\otimes}_\pi H_1)^*$. Let \mathbb{N} be a continuous local martingale with values in H_1 . We need to show that $[\mathbb{A}, \mathbb{N}]_\nu = 0$. We consider the random maps $T^\epsilon: \nu \times \Omega \rightarrow \mathcal{C}([0, T])$ defined by

$$T^\epsilon(\phi) := [\mathbb{A}, \mathbb{N}]_\nu^\epsilon(\phi) = \frac{1}{\epsilon} \int_0^\cdot \nu^* \langle (\mathbb{A}(r + \epsilon) - \mathbb{A}(r)) \otimes (\mathbb{N}(r + \epsilon) - \mathbb{N}(r)), \phi \rangle_\nu dr,$$

for $\phi \in \nu$.

Step 1. Suppose that $\phi = h^* \otimes k^*$ for $h^* \in \chi_0$ and $k \in H_1$. Then

$$\begin{aligned} T^\epsilon(\phi)(t) &= \frac{1}{\epsilon} \int_0^t \chi_0^* \langle (\mathbb{A}(r + \epsilon) - \mathbb{A}(r)), h^* \rangle_{\chi_0} \langle (\mathbb{N}(r + \epsilon) - \mathbb{N}(r)), k \rangle_{H_1} dr \\ &\leq \left[\frac{1}{\epsilon} \int_0^t \chi_0^* \langle (\mathbb{A}(r + \epsilon) - \mathbb{A}(r)), h^* \rangle_{\chi_0}^2 dr \frac{1}{\epsilon} \int_0^t \langle (\mathbb{N}(r + \epsilon) - \mathbb{N}(r)), k \rangle_{H_1}^2 dr \right]^{1/2} \\ &= \left[\frac{1}{\epsilon} \int_0^t \chi^* \langle (\mathbb{A}(r + \epsilon) - \mathbb{A}(r))^{\otimes 2}, h^* \otimes h^* \rangle_\chi dr \right]^{1/2} \\ &\quad \times \left[\frac{1}{\epsilon} \int_0^t \langle (\mathbb{N}(r + \epsilon) - \mathbb{N}(r)), k \rangle_{H_1}^2 dr \right]^{1/2}, \quad (25) \end{aligned}$$

that converges ucp to

$$([\mathbb{A}, \mathbb{A}](t)(h^* \otimes h^*)[\mathbb{N}, \mathbb{N}]^{cl}(t)(k^* \otimes k^*))^{1/2} = 0,$$

since the quadratic variation of a local martingale is the classical one and taking into account item (i) of Remark 3.6.

Step 2. We denote by \mathcal{D} the linear combinations of elements of the form $h^* \otimes k^*$ for $h^* \in \chi_0$ and $k \in H_1$. We remark that \mathcal{D} is dense in ν . From the convergence found in *Step 1*, it follows that, for every $\phi \in \mathcal{D}$, ucp we have $T^\epsilon(\phi) \xrightarrow{\epsilon \rightarrow 0} 0$.

Step 3. We consider a generic $\phi \in \nu$. By Lemma A.2, for $t \in [0, T]$ it follows

$$\begin{aligned}
|T^\epsilon(\phi)(t)| &\leq |\phi|_\nu \int_0^t \frac{|(\mathbb{A}(r+\epsilon) - \mathbb{A}(r)) \otimes (\mathbb{N}(r+\epsilon) - \mathbb{N}(r))|_{\nu^*}}{\epsilon} dr \\
&= |\phi|_\nu \frac{1}{\epsilon} \int_0^t |(\mathbb{N}(r+\epsilon) - \mathbb{N}(r))|_{H_1} |(\mathbb{A}(r+\epsilon) - \mathbb{A}(r))|_{\chi_0^*} dr \\
&\leq |\phi|_\nu \left(\frac{1}{\epsilon} \int_0^t |(\mathbb{N}(r+\epsilon) - \mathbb{N}(r))|_{H_1}^2 dr \frac{1}{\epsilon} \int_0^t |(\mathbb{A}(r+\epsilon) - \mathbb{A}(r))|_{\chi_0^*}^2 dr \right)^{\frac{1}{2}} \\
&= |\phi|_\nu \left(\frac{1}{\epsilon} \int_0^t |(\mathbb{N}(r+\epsilon) - \mathbb{N}(r))|_{H_1}^2 dr \times \frac{1}{\epsilon} \int_0^t |(\mathbb{A}(r+\epsilon) - \mathbb{A}(r))^{\otimes 2}|_{\chi^*} dr \right)^{\frac{1}{2}}. \quad (26)
\end{aligned}$$

To prove that $[\mathbb{A}, \mathbb{N}]_\nu = 0$ we check the corresponding conditions **H1** and **H2** of the Definition 3.1. By Lemma 3.8 we know that \mathbb{N} admits a global quadratic variation i.e. a $(H_1 \otimes H_1)^*$ -quadratic variation. By condition **H1** of the Definition 3.1 related to $(H_1 \otimes H_1)^*$ -quadratic variation for the process \mathbb{N} and the χ -quadratic variation of \mathbb{A} , for any sequence (ϵ_n) converging to zero, there is a subsequence (ϵ_{n_k}) such that the sequence $T^{\epsilon_{n_k}}(\phi)$ is bounded for any ϕ in the $\mathcal{C}[0, T]$ metric a.s. condition **H1** of the ν -covariation. By Banach-Steinhaus for F -spaces (Theorem 17, Chapter II in [14]) it follows that $T^\epsilon(\phi) \xrightarrow{\epsilon \rightarrow 0} 0$ ucp for all $\phi \in \nu$ and so condition **H2** and the final result follows. \square

Corollary 3.17. *Assume that the hypotheses of Proposition 3.16 are satisfied. If \mathbb{X} is a χ -Dirichlet process then we have the following.*

- (i) \mathbb{X} is a ν -weak-Dirichlet process.
- (ii) \mathbb{X} is a χ -weak Dirichlet process.
- (iii) \mathbb{X} is a χ -finite-quadratic-variation process.

Proof. (i) follows by Proposition 3.16. As far as (ii) is concerned, let $\mathbb{X} = \mathbb{M} + \mathbb{A}$ be a χ -Dirichlet process decomposition, where \mathbb{M} is a local martingale. Setting $H_1 = H$, then χ is included in ν , so Proposition 3.16 implies that \mathbb{A} is a χ -orthogonal process and so (ii) follows. We prove now (iii): By Lemma 3.8 and Lemma 3.4(i) \mathbb{M} admits a χ -quadratic variation. By the bilinearity of the χ -covariation, it is enough to show that $[\mathbb{M}, \mathbb{A}]_\chi = 0$. This follows from item (ii). \square

Proposition 3.18. *Let B_1 and B_2 be two real separable Banach spaces and χ a Chi-subspace of $(B_1 \hat{\otimes}_\pi B_2)^*$. Let \mathbb{X} and \mathbb{Y} be two stochastic processes with values respectively in B_1 and B_2 such that (\mathbb{X}, \mathbb{Y}) admits a χ -covariation. Let G be a continuous measurable process $G : [0, T] \times \Omega \rightarrow \mathcal{K}$ where \mathcal{K} is a closed separable subspace of χ . Then for every $t \in [0, T]$*

$$\int_0^t \chi \langle G(\cdot, r), [\mathbb{X}, \mathbb{Y}]^\epsilon(\cdot, r) \rangle_{\chi^*} dr \xrightarrow{\epsilon \rightarrow 0} \int_0^t \chi \langle G(\cdot, r), d[\widetilde{\mathbb{X}}, \widetilde{\mathbb{Y}}](\cdot, r) \rangle_{\chi^*} \quad (27)$$

in probability.

Proof. See [10] Proposition 3.7. \square

We state below the most important result related to the stochastic calculus part of the paper. It generalizes the finite dimensional result contained in [21] Theorem 4.14. Indeed it provides a (generalized) Fukushima-Dirichlet decomposition of a $C^{0,1}$ function u of a (suitably defined weak) Dirichlet process \mathbb{X} . The result below constitutes indeed a chain rule type allowing to expand a non-smooth function u of a convolution type process \mathbb{X} , in substitution of Itô formula. Similar results have been used for several purposes in stochastic control and forward-backward stochastic differential equations (FBSDEs): in the first case in finite dimension by [20] to obtain a verification theorem and by [18] to obtain identify the

solution the process Z as part of the solution of an infinite dimensional FBSDE, as a *generalized gradient* of a solution a semilinear PDE. The difficulty here is that the underlying process \mathbb{X} is far from being a semimartingale.

The definition of real weak Dirichlet process is recalled in Definition 1.1.

Theorem 3.19. *Let ν_0 be a Banach subspace continuously embedded in H^* . Define $\nu := \nu_0 \hat{\otimes}_\pi \mathbb{R}$ and $\chi := \nu_0 \hat{\otimes}_\pi \nu_0$. Let $F: [0, T] \times H \rightarrow \mathbb{R}$ be a $C^{0,1}$ -function. Denote with $\partial_x F$ the Frechet derivative of F with respect to x and assume that the mapping $(t, x) \mapsto \partial_x F(t, x)$ is continuous from $[0, T] \times H$ to ν_0 . Let $\mathbb{X}(t) = \mathbb{M}(t) + \mathbb{A}(t)$ for $t \in [0, T]$ be an ν -weak-Dirichlet process with finite χ -quadratic variation. Then $Y(t) := F(t, \mathbb{X}(t))$ is a (real) weak Dirichlet process with local martingale part*

$$R(t) = F(0, \mathbb{X}(0)) + \int_0^t \langle \partial_x F(r, \mathbb{X}(r)), d\mathbb{M}(r) \rangle.$$

Proof. By definition \mathbb{X} can be written as the sum of a continuous local martingale \mathbb{M} and a ν -martingale-orthogonal process \mathbb{A} .

Let N be a real-valued local martingale. Taking into account Lemma 3.9 and that the covariation of two real local martingales defined in (1), coincides with the classical covariation, it is enough to prove that

$$[F(\cdot, \mathbb{X}(\cdot)), N](t) = \int_0^t \langle \partial_x F(r, \mathbb{X}(r)), d[\mathbb{M}, N]^{cl}(r) \rangle, \quad \text{for all } t \in [0, T].$$

Let $t \in [0, T]$. We evaluate the ϵ -approximation of the covariation, i.e.

$$\frac{1}{\epsilon} \int_0^t (F(r + \epsilon, \mathbb{X}(r + \epsilon)) - F(r, \mathbb{X}(r))) (N(r + \epsilon) - N(r)) \, dr.$$

It equals $I_1(t, \epsilon) + I_2(t, \epsilon)$, where

$$I_1(t, \epsilon) = \int_0^t (F(r + \epsilon, \mathbb{X}(r + \epsilon)) - F(r + \epsilon, \mathbb{X}(r))) \frac{(N(r + \epsilon) - N(r))}{\epsilon} \, dr$$

and

$$I_2(t, \epsilon) = \int_0^t (F(r + \epsilon, \mathbb{X}(r)) - F(r, \mathbb{X}(r))) \frac{(N(r + \epsilon) - N(r))}{\epsilon} \, dr.$$

We prove now that

$$I_1(t, \epsilon) \xrightarrow{\epsilon \rightarrow 0} \int_0^t \langle \partial_x F(r, \mathbb{X}(r)), d[\mathbb{M}, N]^{cl}(r) \rangle \quad (28)$$

in probability; in fact $I_1(t, \epsilon) = I_{11}(t, \epsilon) + I_{12}(t, \epsilon)$ where

$$I_{11}(t, \epsilon) := \int_0^t \frac{1}{\epsilon} \langle \partial_x F(r, \mathbb{X}(r)), \mathbb{X}(r + \epsilon) - \mathbb{X}(r) \rangle (N(r + \epsilon) - N(r)) \, dr,$$

$$I_{12}(t, \epsilon) := \int_0^1 \int_0^t \frac{1}{\epsilon} \langle \partial_x F(r + \epsilon, a\mathbb{X}(r) + (1 - a)\mathbb{X}(r + \epsilon)) - \partial_x F(r, \mathbb{X}(r)), \mathbb{X}(r + \epsilon) - \mathbb{X}(r) \rangle (N(r + \epsilon) - N(r)) \, dr \, da.$$

Now we apply Proposition 3.18 with $B_1 = H$, $B_2 = \mathbb{R}$, $\mathbb{X} = \mathbb{M}$, $\mathbb{Y} = N$, $\chi = \nu$

$$I_{11}(t, \epsilon) \xrightarrow{\epsilon \rightarrow 0} \int_0^t \nu \left\langle \partial_x F(r, \mathbb{X}(r)), d[\widetilde{\mathbb{X}}, \widetilde{N}](r) \right\rangle_{\nu^*}. \quad (29)$$

Recalling that $\mathbb{X} = \mathbb{M} + \mathbb{A}$, we remark that $[\mathbb{X}, N]_\nu$ exists and the ν^* -valued process $[\widetilde{\mathbb{X}}, N]_\nu$ equals $[\widetilde{\mathbb{M}}, N]_\nu + [\widetilde{\mathbb{A}}, N]_\nu = [\widetilde{\mathbb{M}}, N]_\nu$, since \mathbb{A} is a ν -martingale orthogonal process. Taking into account the formalism of Proposition 3.10, Lemma 3.4(i) and Proposition 3.11 if $\Phi \in H \equiv H^*$, we have

$$\left\langle \Phi, [\widetilde{\mathbb{M}}, N]_\nu \right\rangle_{\nu^*} = \left\langle \Phi, [\widetilde{\mathbb{M}}, N]_\nu \right\rangle_{\nu_0^*} =_{H^*} \left\langle \Phi, [\widetilde{\mathbb{M}}, N] \right\rangle_{H^{**}} =_{H^*} \left\langle \Phi, [\mathbb{M}, N]^{cl} \right\rangle_H.$$

Consequently, it is not difficult to show that the right-hand side of (29) gives $\int_0^t \left\langle \partial_x F(r, \mathbb{X}(r)), d[\mathbb{M}, N]^{cl}(r) \right\rangle_H$.

For a fixed $\omega \in \Omega$ we consider the function $\partial_x F$ restricted to $[0, T] \times K$ where K is the (compact) subset of H obtained as convex hull of $\{a\mathbb{X}(r_1) + (1-a)\mathbb{X}(r_2) : r_1, r_2 \in [0, T]\}$. $\partial_x F$ restricted to $[0, T] \times K$ is uniformly continuous with values in ν_0 . Consequently, for ω -a.s.

$$|I_{12}(t, \epsilon)| \leq \int_0^T \delta \left(\partial_x F_{|[0, T] \times K}; \epsilon + \sup_{|r-t| \leq \epsilon} |\mathbb{X}(r) - \mathbb{X}(t)|_{\nu_0^*} \right) \times |\mathbb{X}(r + \epsilon) - \mathbb{X}(r)|_{\nu_0^*} \frac{1}{\epsilon} |N(r + \epsilon) - N(r)| dr, \quad (30)$$

where, for a uniformly continuous function $g : [0, T] \times K \rightarrow \nu_0$, $\delta(g; \epsilon)$ is the modulus of continuity $\delta(g; \epsilon) := \sup_{|x-y| \leq \epsilon} |g(x) - g(y)|_{\nu_0}$. In previous formula we have identified H with H^{**} so that $|x|_H \leq |x|_{\nu_0^*}, \forall x \in H$. So (30) is lower than

$$\begin{aligned} & \delta \left(\partial_x F_{|[0, T] \times K}; \epsilon + \sup_{|s-t| \leq \epsilon} |\mathbb{X}(s) - \mathbb{X}(t)|_{\nu_0^*} \right) \times \left(\int_0^T \frac{1}{\epsilon} |N(r + \epsilon) - N(r)|^2 dr \int_0^T \frac{1}{\epsilon} |(\mathbb{X}(r + \epsilon) - \mathbb{X}(r))|_{\nu_0^*}^2 dr \right)^{1/2} \\ &= \delta \left(\partial_x F_{|[0, T] \times K}; \epsilon + \sup_{|s-t| \leq \epsilon} |\mathbb{X}(s) - \mathbb{X}(t)|_{\nu_0^*} \right) \times \left(\int_0^T \frac{1}{\epsilon} |N(r + \epsilon) - N(r)|^2 dr \int_0^T \frac{1}{\epsilon} |(\mathbb{X}(r + \epsilon) - \mathbb{X}(r))^{\otimes 2}|_{\chi^*} dr \right)^{1/2}, \end{aligned} \quad (31)$$

when $\epsilon \rightarrow 0$, where we have used Lemma A.2, for $\alpha = \pi$ with the usual identification. The right-hand side of (31), of course converges to zero, since \mathbb{X} (respectively N) is a χ -finite quadratic variation process (respectively a real finite quadratic variation process) and X is also continuous as a ν_0^* -valued process.

To conclude the proof of the proposition we only need to show that $I_2(t, \epsilon) \xrightarrow{\epsilon \rightarrow 0} 0$. This is relatively simple since

$$I_2(t, \epsilon) = \frac{1}{\epsilon} \int_0^t \Gamma(u, \epsilon) dN(u) + R(t, \epsilon),$$

where $R(t, \epsilon)$ is a boundary term such that $R(t, \epsilon) \xrightarrow{\epsilon \rightarrow 0} 0$ in probability and

$$\Gamma(u, \epsilon) = \frac{1}{\epsilon} \int_{(u-\epsilon)_+}^u (F(r + \epsilon, \mathbb{X}(r)) - F(r, \mathbb{X}(r))) dr.$$

Since $\int_0^T (\Gamma(u, \epsilon))^2 d[N](u) \rightarrow 0$ in probability, Problem 2.27, chapter 3 of [23] implies that $I_2(\cdot, \epsilon) \rightarrow 0$ ucp. The result finally follows. \square

4 The case of convolution type processes

This section concerns applications of the stochastic calculus via regularization to convolution type processes. As a particular case the results apply to mild solutions of infinite dimensional stochastic evolution equations.

Assume that H and U are real separable Hilbert spaces, $Q \in \mathcal{L}(U)$, $U_0 := Q^{1/2}(U)$. Assume that $\mathbb{W}_Q = \{\mathbb{W}_Q(t) : 0 \leq t \leq T\}$ is an U -valued \mathcal{F}_t - Q -Wiener process with $\mathbb{W}_Q(0) = 0$, \mathbb{P} a.s. See Sections

2.1 and 2.2 of [19] for the definition and properties of Q -Wiener processes and the definition of stochastic integral with respect to \mathbb{W}_Q . Denote by $\mathcal{L}_2(U_0, H)$ the Hilbert space of the Hilbert-Schmidt operators from U_0 to H^4 .

In the whole section, as usual in the literature we refer (see e.g. [5, 19]) we will always identify, via Riesz Theorem, the spaces H and H^* . In this way the duality between H^* and H reduces simply to the scalar product in H .

Lemma 4.1. *Consider an $\mathcal{L}(U, H)$ -valued predictable process \mathbb{A} such that $\int_0^T \text{Tr}[\mathbb{A}(r)Q^{1/2}(\mathbb{A}(r)Q^{1/2})^*] dr < \infty$ a.s. and define*

$$\mathbb{M}_t = \int_0^t \mathbb{A}(r) d\mathbb{W}_Q(r), t \in [0, T]. \quad (32)$$

If \mathbb{X} is an H -valued predictable process such that

$$\int_0^T \langle \mathbb{X}(r), \mathbb{A}(r)Q^{1/2}(\mathbb{A}Q^{1/2})^*\mathbb{X}(r) \rangle dr < \infty, \text{ a.s.}, \quad (33)$$

then

$$N(t) = \int_0^t \langle \mathbb{X}(r), d\mathbb{M}(r) \rangle, t \in [0, T], \quad (34)$$

is well-defined and it equals $N(t) = \int_0^t \langle \mathbb{X}(r), \mathbb{A}(r) d\mathbb{W}_Q(r) \rangle$ for $t \in [0, T]$.

Proof. See Section 4.7 of [5]. □

We recall in the following proposition some significant properties of the stochastic integral with respect to local martingales.

Proposition 4.2. *Let \mathbb{A} , \mathbb{M} and N as in Lemma 4.1.*

- (i) $\mathbb{N}(t) := \int_0^t \mathbb{X}(r)d\mathbb{M}(r)$ is a well defined (\mathcal{F}_t) -local martingale.
- (ii) Let \mathbb{K} be an (\mathcal{F}_t) -predictable process such that $\mathbb{K}\mathbb{X}$ fulfills (4). Then the Itô-type stochastic integral $\int_0^t \mathbb{K}d\mathbb{N}$ for $t \in [0, T]$ is well-defined and it equals $\int_0^t \mathbb{K}\mathbb{X}d\mathbb{M}$.
- (iii) If \mathbb{M} is a Q -Wiener process W_Q , then, whenever \mathbb{X} is such that

$$\int_0^T \text{Tr} \left[\left(\mathbb{X}(r)Q^{1/2} \right) \left(\mathbb{X}(r)Q^{1/2} \right)^* \right] dr < +\infty \text{ a.s.}, \quad (35)$$

then $\mathbb{N}(t) = \int_0^t \mathbb{X}(r)d\mathbb{W}_Q(r)$ is a local martingale and $[N]^{\mathbb{R},cl}(t) = \int_0^t \left(\mathbb{X}(r)Q^{1/2} \right) \left(\mathbb{X}(r)Q^{1/2} \right)^* dr$.

- (iv) If in (iii), the expectation of the quantity (35) is finite, then $\mathbb{N}(t) = \int_0^t \mathbb{X}(r)d\mathbb{W}_Q(r)$ is a square integrable continuous martingale.
- (v) If \mathbb{M} is defined as in (32) and \mathbb{X} fulfills (33), then \mathbb{M} is a real local martingale. If moreover, the expectation of (33) is finite, then N , defined in (34), is a square integrable martingale.

Proof. For (i) see [24] Theorem 2.14 page 14-15. For (ii) see [28], proof of Proposition 2.2 Section 2.4. (iii) and (iv) are contained in [5] Theorem 4.12 Section 4.4. (v) is a consequence of (iii) and (iv) and of Lemma 4.1. □

We recall the following fact that concerns the classical tensor covariation.

⁴The definition and the first properties of Hilbert-Schmidt operators can be found in Appendix A.2 of [32] and for more details in [38].

Lemma 4.3. Let $\Psi : ([0, T] \times \Omega, \mathcal{P}) \rightarrow \mathcal{L}_2(U_0, H)$ be a strongly measurable process satisfying

$$\int_0^T \|\Psi(r)\|_{\mathcal{L}_2(U_0, H)}^2 dr < +\infty \quad a.s.$$

Consider the local martingale $\mathbb{M}(t) := \int_0^t \Psi(r) d\mathbb{W}_Q(r)$. Then $[\mathbb{M}, \mathbb{M}]^{cl}(t) = \int_0^t g(r) dr$, where $g(r)$ is the element of $H \hat{\otimes}_\pi H$ associated with the nuclear operator $G_g(r) := (\Psi(r)Q^{1/2})(\Psi(r)Q^{1/2})^*$.

Proof. See [5] Section 4.7. □

We introduce now the class of convolution type processes that includes mild solutions of (2). We denote by $A : D(A) \subseteq H \rightarrow H$ the generator of the C_0 -semigroup e^{tA} (for $t \geq 0$) on H . The reader may consult for instance [1] Part II, Chapter 1 for basic properties of C_0 -semigroups. A^* denotes the adjoint of A . $D(A)$ and $D(A^*)$ are Hilbert spaces when endowed with the graph norm: for any $x \in D(A^*)$ (respectively $x \in D(A)$), $|x|_{D(A^*)}^2 := |x|^2 + |A^*x|^2$ (respectively $|x|_{D(A)}^2 := |x|^2 + |Ax|^2$).

Observe that, as a consequence of the identification of H and H^* , A^* is the generator of a C_0 -semigroup on H and $D(A^*)$ is continuously embedded in H .

Let b be a predictable process with values in H and σ be a predictable process with values in $\mathcal{L}_2(U_0, H)$ such that

$$\mathbb{P} \left[\int_0^T |b(t)| + \|\sigma(t)\|_{\mathcal{L}_2(U_0, H)}^2 dt < +\infty \right] = 1. \quad (36)$$

We are interested in the process

$$\mathbb{X}(t) = e^{tA}x + \int_0^t e^{(t-r)A}b(r) dr + \int_0^t e^{(t-r)A}\sigma(r) d\mathbb{W}_Q(r). \quad (37)$$

We call the processes of this form *convolution type processes*. A mild solution to an equation of type (2) is a particular case of (37). Indeed, once existence and uniqueness of the solution $\mathbb{X}(\cdot)$ of (37) is proved, we can simply take $b(r) = b(r, \mathbb{X}(r))$ and $\sigma(r) = \sigma(r, \mathbb{X}(r))$. Examples of SPDEs that can be rewritten in the form (2) (see e.g. [6] Part III) are for example stochastic heat (and more general parabolic) equations with Dirichlet or Neumann boundary conditions, wave equations, delay equations, reaction-diffusion equations. We define

$$\mathbb{Y}(t) := \mathbb{X}(t) - \int_0^t b(r) dr - \int_0^t \sigma(r) d\mathbb{W}_Q(r) - x. \quad (38)$$

Lemma 4.4. Let b (respectively σ) be a predictable process with values in H (respectively with values in $\mathcal{L}_2(U_0, H)$) such that (36) is satisfied. Let $\mathbb{X}(t)$ be defined by (37) and \mathbb{Y} defined by (38). If $z \in D(A^*)$ we have $\langle \mathbb{Y}(t), z \rangle = \int_0^t \langle \mathbb{X}(r), A^*z \rangle dr$.

Proof. See [30] Theorem 12. □

We want now to prove that \mathbb{Y} has zero- $\bar{\chi}$ -quadratic variation for a suitable space $\bar{\chi}$. We will see that the space

$$\bar{\chi} := D(A^*) \hat{\otimes}_\pi D(A^*) \quad (39)$$

does the job. We set $\bar{v}_0 := D(A^*)$ which is clearly continuously embedded into H . By Lemma A.5, $\bar{\chi}$ is a Chi-subspace of $(H \hat{\otimes}_\pi H)^*$.

Proposition 4.5. The process \mathbb{Y} has zero $\bar{\chi}$ -quadratic variation.

Proof. Observe that, thanks to Lemma 3.18 in [11] it is enough to show that

$$I(\epsilon) := \frac{1}{\epsilon} \int_0^T |(\mathbb{Y}(r+\epsilon) - \mathbb{Y}(r))^{\otimes 2}|_{\bar{\chi}^*} dr \xrightarrow{\epsilon \rightarrow 0} 0, \quad \text{in probability.}$$

In fact, identifying $\bar{\chi}^*$ with $\mathcal{Bi}(\bar{\nu}_0, \bar{\nu}_0; \mathbb{R})$ (see after Notation A.3) we get

$$\begin{aligned} I(\epsilon) &= \frac{1}{\epsilon} \int_0^T \sup_{|\phi|_{\bar{\nu}_0}, |\psi|_{\bar{\nu}_0} \leq 1} |\langle (\mathbb{Y}(r+\epsilon) - \mathbb{Y}(r)), \phi \rangle \langle (\mathbb{Y}(r+\epsilon) - \mathbb{Y}(r)), \psi \rangle| dr \\ &\leq \frac{1}{\epsilon} \int_0^T \sup_{|\phi|_{\bar{\nu}_0}, |\psi|_{\bar{\nu}_0} \leq 1} \left\{ \left| \int_r^{r+\epsilon} \langle \mathbb{X}(\xi), A^* \phi \rangle d\xi \right| \left| \int_r^{r+\epsilon} \langle \mathbb{X}(\xi), A^* \psi \rangle d\xi \right| \right\} dr, \quad (40) \end{aligned}$$

where we have used Lemma 4.4. This is smaller than $\frac{1}{\epsilon} \int_0^T \left| \int_r^{r+\epsilon} |\mathbb{X}(\xi)| d\xi \right|^2 dr \leq \epsilon \sup_{\xi \in [0, T]} |\mathbb{X}(\xi)|^2$, which converges to zero almost surely. \square

Corollary 4.6. *The process \mathbb{X} is a $\bar{\chi}$ -Dirichlet process. Moreover it is also a $\bar{\chi}$ finite quadratic variation process and a $\bar{\nu}_0 \hat{\otimes}_\pi \mathbb{R}$ -weak-Dirichlet process.*

Proof. For $t \in [0, T]$, we have $\mathbb{X}(t) = \mathbb{M}(t) + \mathbb{A}(t)$, where $\mathbb{M}(t) := x + \int_0^t \sigma(r) d\mathbb{W}_Q(r)$, $\mathbb{A}(t) := \mathbb{V}(t) + \mathbb{Y}(t)$ and $\mathbb{V}(t) = \int_0^t b(r) dr$. \mathbb{M} is a local martingale by Proposition 4.2 (iii) and \mathbb{V} is a bounded variation process. By Proposition 3.5 and Lemma 3.4(i), we get $[\mathbb{V}, \mathbb{V}]_{\bar{\chi}} = [\mathbb{V}, \mathbb{Y}]_{\bar{\chi}} = [\mathbb{Y}, \mathbb{V}]_{\bar{\chi}} = 0$. By Proposition 4.5 and the bilinearity of the $\bar{\chi}$ -covariation, it yields that \mathbb{A} has a zero $\bar{\chi}$ -quadratic variation and so \mathbb{X} is a $\bar{\chi}$ -Dirichlet process. The second part of the statement is a consequence of Corollary 3.17. \square

Remark 4.7. *Corollary 4.6 allows to apply the Fukushima type Theorem 3.19 to expand a $F(t, \mathbb{X}(t))$, where F is a $C^{0,1}$ -function and \mathbb{X} is a convolution type process, so for instance the solution of a stochastic differential equation of evolution type. This looks as a Itô generalized chain rule for non-smooth function F and it was applied in [16] to obtain the stochastic control verification Theorem 6.11.*

In the sequel we will denote by $UC([0, T] \times H; D(A^*))$ ($D(A^*)$ is equipped with the graph norm) the F -space of the functions $G : [0, T] \times H \rightarrow D(A^*)$ which are uniformly continuous on sets $[0, T] \times D$ for any closed ball D of H , equipped with the topology of the uniform convergence on closed balls.

The theorem below generalizes for some aspects the Itô formula of [11], i.e. their Theorem 5.2, to the case when the second derivatives do not necessarily belong to the Chi-subspace χ .

Theorem 4.8. *Let $F : [0, T] \times H \rightarrow \mathbb{R}$ of class $C^{1,2}$. Suppose that $(t, x) \mapsto \partial_x F(t, x)$ belongs to $UC([0, T] \times H; D(A^*))$. Let \mathbb{X} be an H -valued process admitting a $\bar{\chi}$ -quadratic variation. We suppose the following.*

(i) *There exists a (càdlàg) bounded variation process $C : [0, T] \times \Omega \rightarrow (H \hat{\otimes}_\pi H)$ such that, for all t in $[0, T]$ and $\phi \in \bar{\chi}$,*

$$C(t, \cdot)(\phi) = [\mathbb{X}, \mathbb{X}]_{\bar{\chi}}(\phi)(t, \cdot) \quad \text{a.s.}$$

(ii) *For every continuous function $\Gamma : [0, T] \times H \rightarrow D(A^*)$ the following integral exists:*

$$\int_0^t \langle \Gamma(r, \mathbb{X}(r)), d^- \mathbb{X}(r) \rangle. \quad (41)$$

Then, for any $t \in [0, T]$,

$$\begin{aligned} F(t, \mathbb{X}(t)) &= F(0, \mathbb{X}(0)) + \int_0^t \langle \partial_r F(r, \mathbb{X}(r)), d^- \mathbb{X}(r) \rangle \\ &\quad + \frac{1}{2} \int_0^t \langle \partial_{xx}^2 F(r, \mathbb{X}(r)), dC(r) \rangle_{H \hat{\otimes}_\pi H} + \int_0^t \partial_r F(r, \mathbb{X}(r)) dr, \quad \text{a.s.} \quad (42) \end{aligned}$$

Proof. Step 1. Let $\{e_i^*\}_{i \in \mathbb{N}}$ be an orthonormal basis of H made of elements of $D(A^*) \subseteq H$. This is always possible since $D(A^*) \subseteq H$ densely embedded, via a Gram-Schmidt orthogonalization procedure. For $N \geq 1$ we denote by $P_N: H \rightarrow H$ the orthogonal projection on the span of the vectors $\{e_1^*, \dots, e_N^*\}$. $P_\infty: H \rightarrow H$ simply denotes the identity.

Let us for a moment omit the time dependence on F , which is supposed to be of class C^2 from H to \mathbb{R} . We define $F_N: H \rightarrow \mathbb{R}$ as $F_N(x) := F(P_N X(x))$. We have

$$\partial_x F_N(x) = P_N \partial_x F(P_N X(x)) \quad (43)$$

and

$$\partial_{xx}^2 F_N(x) = (P_N \otimes P_N) \partial_{xx}^2 F(P_N X(x)),$$

where the latter equality has to be understood as

$${}_{(H \hat{\otimes}_\pi H)^*} \langle \partial_{xx}^2 F_N(x), h_1 \otimes h_2 \rangle_{(H \hat{\otimes}_\pi H)} = {}_{(H \hat{\otimes}_\pi H)^*} \langle \partial_{xx}^2 F(P_N(x)), (P_N(h_1)) \otimes (P_N(h_2)) \rangle_{H \hat{\otimes}_\pi H}, \quad (44)$$

for all $h_1, h_2 \in H$. $\partial_{xx}^2 F_N(x)$ is an element of $(H \hat{\otimes}_\pi H)^*$ but it belongs to $(D(A^*) \hat{\otimes}_\pi D(A^*))$ as well; indeed it can be written as

$$\sum_{i,j=1}^N {}_{(H \hat{\otimes}_\pi H)^*} \langle \partial_{xx}^2 F(P_N(x)), e_i^* \otimes e_j^* \rangle_{H \hat{\otimes}_\pi H} (e_i^* \otimes e_j^*)$$

and $e_i^* \otimes e_j^*$ are in fact elements of $D(A^*) \hat{\otimes}_\pi D(A^*)$.

We come back now again to the time dependence notation $F(t, x)$. We can apply the Itô formula proved in [11], Theorem 5.2, and with the help of Assumption (i), we find

$$\begin{aligned} F_N(t, \mathbb{X}(t)) &= F_N(0, \mathbb{X}(0)) + \int_0^t \langle \partial_x F_N(r, \mathbb{X}(r)), d^- \mathbb{X}(r) \rangle \\ &\quad + \frac{1}{2} \int_0^t {}_{(H \hat{\otimes}_\pi H)^*} \langle \partial_{xx}^2 F_N(r, \mathbb{X}(r)), dC(r) \rangle_{H \hat{\otimes}_\pi H} + \int_0^t \partial_r F_N(r, \mathbb{X}(r)) dr. \end{aligned} \quad (45)$$

Step 2. We consider, for fixed $\epsilon > 0$, the map

$$\begin{cases} T_\epsilon: UC([0, T] \times H; D(A^*)) \rightarrow L^0(\Omega) \\ T_\epsilon: G \mapsto \int_0^t \left\langle G(r, \mathbb{X}(r)), \frac{\mathbb{X}(r+\epsilon) - \mathbb{X}(r)}{\epsilon} \right\rangle dr, \end{cases}$$

where the set $L^0(\Omega)$ of all real random variables is equipped with the topology of the convergence in probability. Assumption (ii) implies that $\lim_{\epsilon \rightarrow 0} T_\epsilon G$ exists for every G . By Banach-Steinhaus for F -spaces (see Theorem 17, Chapter II in [14]) it follows that the map

$$\begin{cases} UC([0, T] \times H; D(A^*)) \rightarrow L^0(\Omega) \\ G \mapsto \int_0^t \langle G(r, \mathbb{X}(r)), d^- \mathbb{X}(r) \rangle \end{cases}$$

is linear and continuous.

Step 3. If $K \subseteq H$ is a compact set then the set $P(K) := \{P_N(y) : y \in K, N \in \mathbb{N} \cup +\infty\}$ is compact as well. Indeed, consider $\{P_{N_l}(y_l)\}_{l \geq 1}$ be a sequence in $P(K)$. We look for a subsequence convergence to an element of $P(K)$. Since K is compact we can assume, without loss of generality, that y_l converges, for $l \rightarrow +\infty$, to some $y \in K$. If $\{N_l\}$ assumes only a finite number of values then (passing if necessary to a subsequence) $N_l \equiv \bar{N}$ for some $\bar{N} \in \mathbb{N} \cup +\infty$ and then $P_{N_l}(y_l) \xrightarrow{l \rightarrow +\infty} P_{\bar{N}}(y)$. Otherwise we can

assume (passing if necessary to a subsequence) that $N_l \xrightarrow{l \rightarrow +\infty} +\infty$ and then it is not difficult to prove that $P_{N_l}(y_l) \xrightarrow{l \rightarrow +\infty} y$, which belongs to $P(K)$ since $y = P_\infty y$. In particular, being $\partial_x F$ continuous,

$$\mathcal{D} := \{\partial_x F(P_N(x)) : x \in K, N \in \mathbb{N} \cup \{+\infty\}\}$$

is compact in $D(A^*)$. Since the sequence of maps $\{P_N\}$ is uniformly continuous, it follows that

$$\sup_{x \in \mathcal{D}} |(P_N - I)(x)| \xrightarrow{N \rightarrow \infty} 0. \quad (46)$$

Step 4. We show now that

$$\lim_{N \rightarrow \infty} \int_0^t \langle \partial_x F_N(r, \mathbb{X}(r)), d^- \mathbb{X}(r) \rangle = \int_0^t \langle \partial_x F(r, \mathbb{X}(r)), d^- \mathbb{X}(r) \rangle \quad (47)$$

holds in probability for every $t \in [0, T]$. Let K be a compact subset of H . In fact

$$\sup_{t \in [0, T], x \in K} |\partial_x F(t, P_N(x)) - \partial_x F(t, x)| \xrightarrow{N \rightarrow \infty} 0,$$

since $\partial_x F$ is continuous. On the other hand

$$\sup_{t \in [0, T], x \in K} |(P_N - I)(\partial_x F(t, P_N x))| \xrightarrow{N \rightarrow \infty} 0,$$

because of (46). Consequently, by (43),

$$\partial_x F_N \rightarrow \partial_x F, \quad (48)$$

uniformly on each compact, with values in H . This yields that ω -a.s.

$$\partial_x F_N(r, \mathbb{X}(r)) \rightarrow \partial_x F(r, \mathbb{X}(r)),$$

uniformly on each compact. By Step 2, then (47) follows.

Step 5. Finally, we prove that

$$\lim_{N \rightarrow \infty} \frac{1}{2} \int_0^t \langle \partial_{xx}^2 F_N(r, \mathbb{X}(r)), dC(r) \rangle_{H \hat{\otimes}_\pi H} = \frac{1}{2} \int_0^t \langle \partial_{xx}^2 F(r, \mathbb{X}(r)), dC(r) \rangle_{H \hat{\otimes}_\pi H}. \quad (49)$$

For a fixed $\omega \in \Omega$ we define $K(\omega)$ the compact set as $K(\omega) := \{\mathbb{X}(t)(\omega) : t \in [0, T]\}$. We write

$$\begin{aligned} & \left| \int_0^t \langle \partial_{xx}^2 F_N(r, \mathbb{X}(x)) - \partial_{xx}^2 F(r, \mathbb{X}(x)), dC(r) \rangle_{H \hat{\otimes}_\pi H} \right|(\omega) \\ & \leq \sup_{\substack{y \in K(\omega) \\ t \in [0, T]}} |\partial_{xx}^2 F_N(t, y) - \partial_{xx}^2 F(t, y)|_{(H \hat{\otimes}_\pi H)^*} \int_0^T d|C|_{\text{var}}(r)(\omega), \end{aligned} \quad (50)$$

where $r \mapsto |C(r)|_{\text{var}}$ is to total variation function of C . Using arguments similar to those used in proving (48) one can see that $\partial_{xx}^2 F_N \xrightarrow{N \rightarrow \infty} \partial_{xx}^2 F$ uniformly on each compact. Consequently

$$\sup_{r \in [0, T]} |(\partial_{xx}^2 F_N - \partial_{xx}^2 F)(r, \mathbb{X}(r))|_{(H \hat{\otimes}_\pi H)^*} \xrightarrow{N \rightarrow \infty} 0.$$

Since C has bounded variation, finally (49) holds.

Step 6. Since F_N (respectively $\partial_r F_N$) converges uniformly on each compact to F (respectively $\partial_r F$), when $N \rightarrow \infty$, then

$$\int_0^t \partial_r F_N(r, \mathbb{X}(r)) dr \xrightarrow{N \rightarrow \infty} \int_0^t \partial_r F(r, \mathbb{X}(r)) dr.$$

Taking the limit when $N \rightarrow \infty$ in (45), finally provides (42). \square

Lemma 4.9. *The conditions (i) and (ii) of Theorem 4.8 are verified if $\mathbb{X} = \mathbb{M} + \mathbb{V} + \mathbb{S}$, where \mathbb{M} is a local martingale, \mathbb{V} is an H -valued bounded variation process, and \mathbb{S} is a process verifying*

$$\langle \mathbb{S}, h \rangle (t) = \int_0^t \langle \mathbb{Z}(r), A^* h \rangle dr, \quad \text{for all } h \in D(A^*),$$

for some measurable process \mathbb{Z} with $\int_0^T |\mathbb{Z}(r)|^2 dr < +\infty$ a.s.

Proof. By Lemma 3.8, \mathbb{M} admits a global quadratic variation which can be identified with $[\mathbb{M}, \mathbb{M}]^{cl}$. On the other hand $\mathbb{A} = \mathbb{V} + \mathbb{S}$ has a zero $\bar{\chi}$ -quadratic variation, by Proposition 3.5 and the bilinearity character of the $\bar{\chi}$ -covariation. \mathbb{X} is therefore a $\bar{\chi}$ -Dirichlet process. By Corollary 3.17 and again the bilinearity of the $\bar{\chi}$ -covariation, we obtain that \mathbb{X} has a finite $\bar{\chi}$ -quadratic variation. Taking also into account Lemma 3.8 and Lemma 3.4(i), we get $\widetilde{[\mathbb{X}, \mathbb{X}]_{\bar{\chi}}}(\Phi)(\cdot) = \langle [\mathbb{M}, \mathbb{M}]^{cl}, \Phi \rangle$ if $\Phi \in \bar{\chi}$. Consequently, we can set $C = [\mathbb{M}, \mathbb{M}]^{cl}$ and condition (i) is verified.

To prove (ii) consider a continuous function $\Gamma: [0, T] \times H \rightarrow D(A^*)$. The integral of $(\Gamma(r), \mathbb{X}(r))$ with respect to the semimartingale $\mathbb{M} + \mathbb{V}$ where $\mathbb{M}(t) = x + \int_0^t \sigma(r) dW_Q(r)$ exists and equals the classical Itô integral, by Proposition 2.3 and Proposition 2.4. Therefore, we only have to prove that

$$\int_0^t \langle \Gamma(r), \mathbb{X}(r) \rangle, d^- \mathbb{S}(r) \rangle, \quad t \in [0, T],$$

exists. For every $t \in [0, T]$ the ϵ -approximation of such an integral gives, up to a remainder boundary term $C(\epsilon, t)$ which converges in probability to zero,

$$\begin{aligned} \frac{1}{\epsilon} \int_0^t \langle \Gamma(r), \mathbb{X}(r) \rangle, \mathbb{S}(r + \epsilon) - \mathbb{S}(r) \rangle dr &= \frac{1}{\epsilon} \int_0^t \int_r^{r+\epsilon} \langle \mathbb{Z}(u), A^* \Gamma(r, \mathbb{X}(r)) \rangle du dr \\ &= \frac{1}{\epsilon} \int_0^t \int_{u-\epsilon}^u \langle \mathbb{Z}(u), A^* \Gamma(r, \mathbb{X}(r)) \rangle dr du \xrightarrow{\epsilon \rightarrow 0} \int_0^t \langle \mathbb{Z}(u), A^* \Gamma(u, \mathbb{X}(u)) \rangle du, \end{aligned} \quad (51)$$

in probability by classical Lebesgue integration theory. The right-hand side of (51) has obviously a continuous modification so (41) exists by definition and condition (ii) is fulfilled. \square

Next result can be considered a Itô formula for mild type processes. An interesting contribution in this direction, but in a different spirit appears in [4].

Corollary 4.10. *Assume that b is a predictable process with values in H and σ is a predictable process with values in $\mathcal{L}_2(U_0, H)$ satisfying (36). Define \mathbb{X} as in (37). Let x be an element of H . Assume that $f \in C^{1,2}([0, T] \times H)$ with $\partial_x f \in UC([0, T] \times H, D(A^*))$. Then, \mathbb{P} -a.s.,*

$$\begin{aligned} f(t, \mathbb{X}(t)) &= f(0, x) + \int_0^t \partial_s f(r, \mathbb{X}(r)) dr + \int_0^t \langle A^* \partial_x f(r, \mathbb{X}(r)), \mathbb{X}(r) \rangle dr + \int_0^t \langle \partial_x f(r, \mathbb{X}(r)), b(r) \rangle dr \\ &\quad + \frac{1}{2} \int_0^t \text{Tr} \left[\left(\sigma(r) Q^{1/2} \right) \left(\sigma(r) Q^{1/2} \right)^* \partial_{xx}^2 f(r, \mathbb{X}(r)) \right] dr + \int_0^t \langle \partial_x f(r, \mathbb{X}(r)), \sigma(r) dW_Q(r) \rangle, \end{aligned} \quad (52)$$

where the partial derivative $\partial_{xx}^2 f(r, x)$ for any $r \in [0, T]$ and $x \in H$ stands in fact for its associated linear bounded operator in the sense of (60).

Proof. It is a consequence of Theorem 4.8 taking into account Lemma 4.9: we have $\mathbb{M}(t) = x + \int_0^t \sigma(r) dW_Q(r)$, $t \in [0, T]$, $\mathbb{V}(t) = \int_0^t b(r) dr$, $\mathbb{S} = \mathbb{Y}$ with $\mathbb{Z}(r) = \mathbb{X}(r)$. According to that lemma, in Theorem 4.8 we set $C = [\mathbb{M}, \mathbb{M}]^{cl}$. We also use the chain rule for Itô's integrals in Hilbert spaces, see the considerations before Proposition 4.2, together with Lemma 4.3. The fourth integral in the right-hand side of (52) appears from the second integral in (42) using Proposition A.4 and Lemma 4.3. \square

4.1 Extension to fractional processes

The class of convolution processes on which the concepts of Section 3 can be applied can be extended to include fractional processes. We sketch this fact in a simplified framework. We consider again H a separable Hilbert space and A the generator of a C_0 -semigroup on H .

Let $\mathbb{B}: [0, T] \times \Omega \rightarrow H$ be a continuous progressively measurable H -valued process with a finite number of modes of the form

$$\mathbb{B}(t) := \sum_{i=1}^N h_i \beta_i(t),$$

where, for any i , h_i is a fixed elements of H and β_i is a γ -Hölder continuous real-valued process, $\gamma > \frac{1}{2}$. We suppose that, for any $i = 1, \dots, N$, the H -valued function $t \mapsto e^{tA} h_i$ (defined on $[0, T]$), is α -Hölder continuous with values in H . By Proposition 2.5

$$\int_0^t e^{(t-s)A} d^- \mathbb{B}(s) \quad \text{and} \quad \int_0^t e^{(t-s)A} d^+ \mathbb{B}(s)$$

are well defined and equal

$$\sum_{i=1}^N \int_0^t e^{(t-s)A} h_i d^y \beta_i(s).$$

Example 4.11. *Suppose that A is the generator of an analytic semigroup on H and that 0 is in the resolvent of A . It is the case for instance of the heat semigroup with Dirichlet boundary condition acting on bounded and regular domains in \mathbb{R}^m . If we choose $h_i \in D((-A)^\alpha)$ for $\alpha \in (0, 1]$, we have (see e.g. [31], Theorem 6.13, page 74), for any $s, t \in [0, T]$ with $s < t$, $|(e^{tA} - e^{sA})h_i| \leq C_T(t-s)^\alpha |h_i|_{D((-A)^\alpha)}$ and then $t \mapsto e^{tA} h_i$ is α -Hölder continuous.*

In the setting described above consider the process

$$\mathbb{X}_1(t) := \mathbb{X}(t) + \int_0^t e^{(t-s)A} d^- \mathbb{B}(s),$$

where \mathbb{X} is of the form (37). We show now that Corollaries 4.6 and 4.10 also hold in the present extended framework.

Proposition 4.12. *The process \mathbb{X}_1 is a $\bar{\chi}$ -Dirichlet process. Moreover it is also a $\bar{\chi}$ finite quadratic variation process and a $\bar{\nu}_0 \hat{\otimes}_\pi \mathbb{R}$ -weak-Dirichlet process.*

Sketch of the proof. Using the properties of the Young integral one can show that the process $\mathbb{A}_1(t) := \int_0^t e^{(t-s)A} d^- \mathbb{B}(s)$ is Hölder continuous with parameter $\gamma > 1/2$. So \mathbb{A}_1 has a zero scalar quadratic variation and then, by additivity and Corollary 4.6, \mathbb{X}_1 has a $\bar{\chi}$ quadratic variation and $[\mathbb{X}_1, \mathbb{X}_1]_{\bar{\chi}} = [\mathbb{X}, \mathbb{X}]_{\bar{\chi}}$. A similar argument shows that \mathbb{X}_1 is a $\bar{\nu}_0 \hat{\otimes}_\pi \mathbb{R}$ -weak-Dirichlet process. \square

Proposition 4.13. *Suppose that hypotheses of Corollary 4.10 hold. Suppose that the trajectories of the process b belong a.s. to $L^p(0, T; H)$ for any $p < 2$. Then the formula (52) still holds with the addition of the term $\int_0^t \langle \partial_x f(s, \mathbb{X}_1(s)), d^- \mathbb{B}(s) \rangle$*

Remark 4.14. *Observe that $\int_0^t \langle \partial_x f(s, \mathbb{X}_1(s)), d^- \mathbb{B}(s) \rangle$ equals $\sum_{i=1}^N \int_0^t \partial_x f(s, \mathbb{X}_1(s)) h_i d^y \beta_i(s)$. Indeed $\int_0^\cdot \sigma(s) d\mathbb{W}(s)$ and \mathbb{V} are α -Hölder continuous for any $\alpha < 1/2$. So \mathbb{X} is α -Hölder continuous for any $\alpha < 1/2$ and the same holds for $\partial_x f(\cdot, \mathbb{X}_1(\cdot))$ since $\partial_x f$ is locally Lipschitz-continuous.*

Sketch of the proof of Proposition 4.13. We define $\mathbb{Y}_1(t) := \mathbb{Y}(t) + \mathbb{B}(t)$ where $\mathbb{Y}(t)$ is defined in (38). Applying Lemma 4.4 to $\mathbb{Y}^\varepsilon(t) := \mathbb{Y}(t) - \int_0^t \frac{\mathbb{B}(s) - \mathbb{B}(s-\varepsilon)}{\varepsilon} ds$ we obtain, for any $z \in D(A^*)$, $\langle \mathbb{Y}^\varepsilon(t), z \rangle = \int_0^t \langle \mathbb{X}^\varepsilon(s), A^* z \rangle ds$ where $\mathbb{X}^\varepsilon(t) := \mathbb{X}(t) + \int_0^t e^{(t-s)A} \frac{\mathbb{B}(s) - \mathbb{B}(s-\varepsilon)}{\varepsilon} ds$. Using the fact that $\int_0^t e^{(t-s)A} d^+ \mathbb{B}(s) = \lim_{\varepsilon \rightarrow 0} \int_0^t e^{(t-s)A} \frac{\mathbb{B}(s) - \mathbb{B}(s-\varepsilon)}{\varepsilon} ds$ the statement of Lemma 4.4 extends to \mathbb{X}_1 , \mathbb{Y}_1 replacing, respectively, \mathbb{X} and \mathbb{Y} . Then, by Proposition 4.5, \mathbb{Y}_1 has a zero $\bar{\chi}$ quadratic variation. The proof can be completed by following the lines of that of Corollary 4.10. \square

References

- [1] A. Bensoussan, G. Da Prato, M. C. Delfour, and S. K. Mitter. *Representation and control of infinite dimensional systems*. Birkhäuser, Boston, 2007.
- [2] Z. Brzezniak and R. Serrano. Optimal relaxed control of dissipative stochastic partial differential equations in banach spaces. *SIAM J. Control Optim.*, 51(3):2664–2703, 2013.
- [3] F. Coquet, A. Jakubowski, J. Mémin, and L. Słomiński. Natural decomposition of processes and weak Dirichlet processes. In *In memoriam Paul-André Meyer: Séminaire de Probabilités XXXIX*, volume 1874 of *Lecture Notes in Math.*, pages 81–116. Springer, Berlin, 2006.
- [4] G. Da Prato, A. Jentzen, and M. Röckner. A mild Itô formula for SPDEs. *Preprint Arxiv*, <http://arxiv.org/pdf/1009.3526.pdf>, 2011.
- [5] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*. Cambridge University Press, 1992.
- [6] G. Da Prato and J. Zabczyk. *Ergodicity for infinite-dimensional systems*. Cambridge University Press, 1996.
- [7] L. Denis. Solutions of stochastic partial differential equations considered as Dirichlet processes. *Bernoulli*, 10(5):783–827, 2004.
- [8] C. Di Girolami and F. Russo. Infinite dimensional stochastic calculus via regularization. *Preprint HAL-INRIA*, <http://hal.archives-ouvertes.fr/inria-00473947/fr/>, 2010.
- [9] C. Di Girolami and F. Russo. Clark-Ocone type formula for non-semimartingales with finite quadratic variation. *C. R. Acad. Sci. Paris Sér. I Math.*, 349(3-4):209 – 214, 2011.
- [10] C. Di Girolami and F. Russo. Generalized covariation and extended Fukushima decomposition for Banach space-valued processes. Applications to windows of Dirichlet processes. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, 15(2), 2012.
- [11] C. Di Girolami and F. Russo. Generalized covariation for Banach space valued processes and Itô formula. *Osaka journal of mathematics.*, 51(3), 2014.
- [12] J. Diestel and J. J. Uhl. *Vector measures*. AMS, Providence, 1977.
- [13] N. Dinculeanu. *Vector integration and stochastic integration in Banach spaces*. Wiley, New York, 2000.
- [14] N. Dunford and J. T. Schwartz. *Linear operators. Part I*. Wiley, New York, 1988.
- [15] M. Errami and F. Russo. n -covariation, generalized Dirichlet processes and calculus with respect to finite cubic variation processes. *Stochastic Process. Appl.*, 104(2):259–299, 2003.
- [16] G. Fabbri and F. Russo. Infinite dimensional weak dirichlet processes, stochastic pdes and optimal control. arXiv preprint arXiv:1207.5710, Unpublished., 2012.
- [17] H. Föllmer. Dirichlet processes. In *Stochastic integrals (Proc. Sympos., Univ. Durham, Durham, 1980)*, volume 851 of *Lecture Notes in Math.*, pages 476–478. Springer, Berlin, 1981.
- [18] M. Fuhrman and G. Tessitore. Generalized directional gradients, backward stochastic differential equations and mild solutions of semilinear parabolic equations. *Appl. Math. Optim.*, 51(3):279–332, 2005.
- [19] L. Gawarecki and V Mandrekar. *Stochastic differential equations in infinite dimensions with applications to stochastic partial differential equations*. Springer, Heidelberg, 2011.
- [20] F. Gozzi and F. Russo. Verification theorems for stochastic optimal control problems via a time dependent Fukushima-Dirichlet decomposition. *Stochastic Process. Appl.*, 116(11):1530–1562, 2006.
- [21] F. Gozzi and F. Russo. Weak Dirichlet processes with a stochastic control perspective. *Stochastic Process. Appl.*, 116(11):1563–1583, 2006.
- [22] M. Gubinelli, A. Lejay, and S. Tindel. Young integrals and SPDEs. *Potential Anal.*, 25(4):307–326, 2006.
- [23] I. Karatzas and S. E. Shreve. *Brownian motion and stochastic calculus*. Springer, New York, 1988.
- [24] N. V. Krylov and B. L. Rozovskii. Stochastic evolution equations. In P. H. Baxendale and S. V. Lototsky, editors, *Stochastic differential equations: theory and applications*, volume 2 of *Interdisciplinary Mathematical Sciences*, pages 1–70. World Scientific, 2007. Translated from Itogi Naukii Tekhniki, Seriya Sovremennye Problemy Matematiki.

- [25] J. A. Leon. Stochastic Fubini theorem for semimartingales in Hilbert space. *Can. J. Math.*, 42(5):890–901, 1990.
- [26] Z.M. Ma and M. Röckner. *Introduction to the theory of (nonsymmetric) Dirichlet forms*. Springer, Berlin, 1992.
- [27] M. Métivier. *Semimartingales: a course on stochastic processes*. Gruyter, Berlin, 1982.
- [28] M. Métivier and J. Pellaumail. *Stochastic integration*. Academic Press, New York, 1980.
- [29] P.A. Meyer. Notes sur les intégrales stochastiques. I Intégrales hilbertiennes. In C. Dellacherie, P. Meyer, and M. Weil, editors, *Séminaire de Probabilités XI*, volume 581 of *Lecture Notes in Mathematics*, pages 446–481. 1977.
- [30] M. Ondreját. Uniqueness for stochastic evolution equations in Banach spaces. *Dissertationes Math. (Rozprawy Mat.)*, 426:63 p., 2004.
- [31] A. Pazy. *Semigroups of linear operators and applications to partial differential equations*, volume 44 of *Applied Mathematical Sciences*. Springer, New York, 1983.
- [32] S. Peszat and J. Zabczyk. *Stochastic partial differential equations with Lévy noise*. Cambridge University Press, 2007.
- [33] M. Pronk and M. Veraar. Forward integration, convergence and non-adapted pointwise multipliers. *Infin. Dimens. Anal. Quantum Probab. Relat. Top.*, 18(1):1550005, 23, 2015.
- [34] F. Russo and P. Vallois. Intégrales progressive, rétrograde et symétrique de processus non adaptés. *C. R. Acad. Sci. Paris Sér. I Math.*, 312(8):615–618, 1991.
- [35] F. Russo and P. Vallois. Forward, backward and symmetric stochastic integration. *Probab. Theory Related Fields*, 97(3):403–421, 1993.
- [36] F. Russo and P. Vallois. Noncausal stochastic integration for l ad l ag processes. In *Stochastic analysis and related topics (Oslo, 1992)*, volume 8 of *Stochastics Monogr.*, pages 227–263. Gordon and Breach, Montreux, 1993.
- [37] F. Russo and P. Vallois. Elements of stochastic calculus via regularization. In *Séminaire de Probabilités XL*, volume 1899 of *Lecture Notes in Math.*, pages 147–185. Springer, Berlin, 2007.
- [38] R. A. Ryan. *Introduction to tensor products of Banach spaces*. Springer, London, 2002.
- [39] E. M. Stein. *Singular integrals and differentiability properties of functions*. Princeton University Press, 1970.
- [40] J. M. A. M. van Neerven, M. C. Veraar, and L. Weis. Stochastic integration in umd Banach spaces. *Ann. Probab.*, 35(4):1438–1478, 2007.
- [41] J. M. A. M. Van Neerven, M. C. Veraar, and L. Weis. Stochastic evolution equations in umd Banach spaces. *J. Appl. Funct. Anal.*, 255(4):940–993, 2008.

A Some functional analysis results

A.1 Notation and basic facts

Let us consider two real separable Banach spaces B_1 and B_2 . We denote by $C(B_1; B_2)$ the set of the continuous functions from B_1 to B_2 . It is a topological vector space if equipped with topology of the uniform convergence on compact sets. If $B_2 = \mathbb{R}$ we often simply use the notation $C(B_1)$ instead of $C(B_1; \mathbb{R})$. Similarly, given a real interval I , continuous B_2 -valued functions defined on $I \times B_1$ while we use the lighter notation $C(I \times B_1)$ when $B_2 = \mathbb{R}$. $C^1(I \times B_1)$ denotes the space of Fréchet continuous differentiable functions $u : I \times B_1 \rightarrow \mathbb{R}$. For a function $u : I \times B_1 \rightarrow \mathbb{R}$, we denote by $\partial_x u(t, x)$ (respectively $\partial_{xx}^2 u(t, x)$), if it exists, the first (respectively second) Fréchet derivative with respect to the variable $x \in B_1$). A function $u \in C(I \times B_1)$ (respectively $u \in C^1(I \times B_1)$) will be said to belong to $C^{0,1}(I \times B_1)$ (respectively $C^{1,2}(I \times B_1)$) if $\partial_x u$ exists and it is continuous, i.e. it belongs to $C(I \times B_1; B_1^*)$ (respectively $\partial_{xx}^2 u(t, x)$ exists for any $(t, x) \in [0, T] \times B_1$ and it is continuous, i.e. it belongs to $C(I \times B_1; \mathcal{B}i(B_1, B_1))$, where $\mathcal{B}i(B_1, B_2)$ is the linear topological space of bilinear bounded forms on $B_1 \times B_2$.

We denote by $\mathcal{L}(B_1; B_2)$ the space of linear bounded maps from B_1 to B_2 and by $\|\cdot\|_{\mathcal{L}(B_1; B_2)}$ the corresponding norm. We indicate by a double bar, i.e. $\|\cdot\|$, the norm of an operator. Often we will consider the case of two separable Hilbert spaces U and H . We denote $|\cdot|$ and $\langle \cdot, \cdot \rangle$ (respectively $|\cdot|_U$ and $\langle \cdot, \cdot \rangle_U$) the norm and the inner product on H (respectively U).

Notation A.1. If H is a Hilbert space, in order to argue more transparently, we often distinguish between H and its dual H^* and with every element $h \in H$ we associate $h^* \in H^*$ through Riesz Theorem.

If $U = H$, we set $\mathcal{L}(U) := \mathcal{L}(U; U)$. $\mathcal{L}_2(U; H)$ will be the set of Hilbert-Schmidt operators from U to H and $\mathcal{L}_1(H)$ (respectively $\mathcal{L}_1^+(H)$) will be the space of (respectively non-negative) nuclear operators on H . For details about the notions of Hilbert-Schmidt and nuclear operator, the reader may consult [38], Section 2.6 and [32] Appendix A.2. If $T \in \mathcal{L}_2(U; H)$ and $T^* : H \rightarrow U$ is the adjoint operator, then $TT^* \in \mathcal{L}_1(H)$ and the Hilbert-Schmidt norm of T is $\|T\|_{\mathcal{L}_2(U; H)}^2 = \|TT^*\|_{\mathcal{L}_1(H)}$. We recall that, for a generic element $T \in \mathcal{L}_1(H)$ and given a basis $\{e_n\}$ of H , the sum $\sum_{n=1}^{\infty} \langle Te_n, e_n \rangle$ is absolutely convergent and independent of the chosen basis $\{e_n\}$. It is called *trace* of T and denoted by $\text{Tr}(T)$. $\mathcal{L}_1(H)$ is a Banach space and we denote by $\|\cdot\|_{\mathcal{L}_1(H)}$ the corresponding norm. If T is non-negative then $\text{Tr}(T) = \|T\|_{\mathcal{L}_1(H)}$ and in general we have the inequalities

$$|\text{Tr}(T)| \leq \|T\|_{\mathcal{L}_1(H)}, \quad \sum_{n=1}^{\infty} |\langle Te_n, e_n \rangle| \leq \|T\|_{\mathcal{L}_1(H)}, \quad (53)$$

see Proposition C.1, [5]. As a consequence, if T is a non-negative operator, the relation below holds:

$$\|T\|_{\mathcal{L}_2(U; H)}^2 = \text{Tr}(TT^*). \quad (54)$$

A.2 Reasonable norms on tensor products

Consider two real separable Banach spaces B_1 and B_2 . Denote, for $i = 1, 2$, with $|\cdot|_i$ the norm on B_i . $B_1 \otimes B_2$ stands for the *algebraic tensor product* i.e. the set of the elements of the form $\sum_{i=1}^n x_i \otimes y_i$ where x_i and y_i are respectively elements of B_1 and B_2 . On $B_1 \otimes B_2$ we identify all the expressions we need in order to ensure that the product $\otimes : B_1 \times B_2 \rightarrow B_1 \otimes B_2$ is bilinear.

On $B_1 \otimes B_2$ we introduce the projective norm π defined, for all $u \in B_1 \otimes B_2$, as

$$\pi(u) := \inf \left\{ \sum_{i=1}^n |x_i|_{B_1} |y_i|_{B_2} : u = \sum_{i=1}^n x_i \otimes y_i \right\}.$$

The *projective tensor product* of B_1 and B_2 , $B_1 \hat{\otimes}_{\pi} B_2$, is the Banach space obtained as completion of $B_1 \otimes B_2$ for the norm π , see [38] Section 2.1, or [8] for further details. For $u \in B_1 \otimes B_2$ of the form $u = \sum_{i=1}^n x_i \otimes y_i$ we define

$$\varepsilon(u) := \sup \left\{ \left| \sum_{i=1}^n \Phi(x_i) \Psi(y_i) \right| : \Phi \in B_1^*, \Psi \in B_2^*, |\Phi|_{B_1^*} = |\Psi|_{B_2^*} = 1 \right\}$$

and denote by $B_1 \hat{\otimes}_{\varepsilon} B_2$ the completion of $B_1 \otimes B_2$ for such a norm: it is the *injective tensor product* of B_1 and B_2 . We remind that $\varepsilon(u)$ does not depend on the representation of u and that, for any $u \in B_1 \otimes B_2$, $\varepsilon(u) \leq \pi(u)$.

A norm α on $B_1 \otimes B_2$ is said to be *reasonable* if for any $u \in B_1 \otimes B_2$,

$$\varepsilon(u) \leq \alpha(u) \leq \pi(u). \quad (55)$$

We denote by $B_1 \hat{\otimes}_{\alpha} B_2$ the completion of $B_1 \otimes B_2$ with respect to the norm α . For any reasonable norm α on $B_1 \otimes B_2$, for any $x \in B_1$ and $y \in B_2$ one has $\alpha(x \otimes y) = |x|_{B_1} |y|_{B_2}$. See [38] Chapter 6.1 for details.

Lemma A.2. Let B_1 and B_2 be two real separable Banach spaces and α a reasonable norm on $B_1 \otimes B_2$. We denote $B := B_1 \hat{\otimes}_{\alpha} B_2$. Choose $a^* \in B_1^*$ and $b^* \in B_2^*$. One can associate to $a^* \otimes b^*$ the elements $i(a^* \otimes b^*)$ of B^* acting as follows on a generic element $u = \sum_{i=1}^n x_i \otimes y_i \in B_1 \otimes B_2$:

$$\langle i(a^* \otimes b^*), u \rangle = \sum_{i=1}^n \langle a^*, x_i \rangle \langle b^*, y_i \rangle.$$

Then $i(a^* \otimes b^*)$ extends by continuity to the whole B and

$$|i(a^* \otimes b^*)|_{B^*} = |a^*|_{B_1^*} |b^*|_{B_2^*}. \quad (56)$$

Proof. We first prove the \leq inequality in (56). Setting $\tilde{a}^* = \frac{a^*}{|a^*|}, \tilde{b}^* = \frac{b^*}{|b^*|}$ we write

$$\langle i(a^* \otimes b^*), u \rangle = \langle i(\tilde{a}^* \otimes \tilde{b}^*), u \rangle |a^*| |b^*| \leq \varepsilon(u) |a^*|_{B_1^*} |b^*|_{B_2^*}.$$

The latter inequality comes from the definition of injective tensor norm ε , considering first $\Phi = a^*, \Psi = B^*$. By (55) $\langle i(a^* \otimes b^*), u \rangle \leq \alpha(u) |a^*|_{B_1^*} |b^*|_{B_2^*}$ and the \leq inequality of (56) is proved.

Concerning the converse inequality, we have $|a^*|_{B_1^*} = \sup_{|\phi|_{B_1}=1} |B_1^* \langle a^*, \phi \rangle|_{B_1}$ and similarly for b^* . So, chosen $\delta > 0$, there exist $\phi_1 \in B_1$ and $\phi_2 \in B_2$ with $|\phi_1|_{B_1} = |\phi_2|_{B_2} = 1$ and

$$|a^*|_{B_1^*} \leq \delta + |B_1^* \langle a^*, \phi_1 \rangle|_{B_1}, \quad |b^*|_{B_2^*} \leq \delta + |B_2^* \langle b^*, \phi_2 \rangle|_{B_2}.$$

We set $u := \phi_1 \otimes \phi_2$. We obtain

$$\begin{aligned} |i(a^* \otimes b^*)|_{B^*} &\geq \frac{|B^* \langle i(a^* \otimes b^*), u \rangle|_B}{|u|_B} = \frac{|B^* \langle i(a^* \otimes b^*), u \rangle|_B}{|\phi_1|_{B_1} |\phi_2|_{B_2}} \\ &= |B_1^* \langle a^*, \phi_1 \rangle|_{B_1} |B_2^* \langle b^*, \phi_2 \rangle|_{B_2} \geq (|a^*|_{B_1^*} - \delta)(|b^*|_{B_2^*} - \delta). \end{aligned} \quad (57)$$

Since $\delta > 0$ is arbitrarily small we finally obtain $|i(a^* \otimes b^*)|_{B^*} \geq |a^*|_{B_1^*} |b^*|_{B_2^*}$. This gives the second inequality and concludes the proof. \square

Notation A.3. When $B_1 = B_2$ and $x \in B_1$ we denote by $x^{\otimes 2}$ the element $x \otimes x \in B_1 \otimes B_1$.

The dual of the projective tensor product $B_1 \hat{\otimes}_\pi B_2$, denoted by $(B_1 \hat{\otimes}_\pi B_2)^*$, can be identified isomorphically with the linear space of bounded bilinear forms on $B_1 \times B_2$ denoted by $\mathcal{B}i(B_1, B_2)$. If $u \in (B_1 \hat{\otimes}_\pi B_2)^*$ and ψ_u is the associated form in $\mathcal{B}i(B_1, B_2)$, we have

$$|u|_{(B_1 \hat{\otimes}_\pi B_2)^*} = \sup_{|a|_{B_1} \leq 1, |b|_{B_2} \leq 1} |\psi_u(a, b)|.$$

See for this [38] Theorem 2.9 Section 2.2, page 22 and also the discussion after the proof of the theorem, page 23. Every element $u \in H \hat{\otimes}_\pi H$ is isometrically associated with an element T_u in the space of nuclear operators $\mathcal{L}_1(H, H)$, defined, for u of the form $\sum_{i=1}^{\infty} a_n \otimes b_n$, as follows:

$$T_u(x) := \sum_{i=1}^{\infty} \langle x, a_n \rangle b_n,$$

see for instance [38] Corollary 4.8 Section 4.1 page 76.

T_u is self-adjoint if and only if there exists a sequence of real numbers (λ_n) and an orthonormal basis (h_n) of H such that

$$u = \sum_{n=1}^{+\infty} \lambda_n h_n \otimes h_n. \quad (58)$$

The ‘‘only is’’ part is obvious. For the ‘‘if’’ part observe that, since T_u is nuclear, it is compact (Proposition A.6 of [32]); if is also self-adjoint there exists (thanks to the spectral theorem) a sequence of real numbers (λ_n) and an orthonormal basis (h_n) of H such that T_u can be written as

$$T_u(x) = \sum_{n=1}^{+\infty} \lambda_n \langle h_n, x \rangle h_n, \quad \text{for all } x \in H; \quad (59)$$

in particular $T_u(h_n) = \lambda_n h_n$ for each n and u can be written as in (58).

To each element ψ of $(H \hat{\otimes}_\pi H)^*$ we associate a bilinear continuous operator B_ψ and a linear continuous operator $L_\psi : H \rightarrow H$ (see [38] page 24, the discussion before Proposition 2.11 Section 2.2) such that

$$\langle L_\psi(x), y \rangle = B_\psi(x, y) = \psi(x \otimes y) \quad \text{for all } x, y \in H. \quad (60)$$

Proposition A.4. Let $u \in H \hat{\otimes}_\pi H$ and $\psi \in (H \hat{\otimes}_\pi H)^*$ with associated maps $T_u \in \mathcal{L}_1(H), L_\psi \in \mathcal{L}(H)$. If T_u is self-adjoint

$${}_{(H \hat{\otimes}_\pi H)^*} \langle \psi, u \rangle_{H \hat{\otimes}_\pi H} = \text{Tr}(L_\psi T_u).$$

Proof. The claim follows from what we have recalled above. Indeed, using (58) and (60) we have

$${}_{(H \hat{\otimes}_\pi H)^*} \langle \psi, u \rangle_{H \hat{\otimes}_\pi H} = \psi(u) = \psi \left(\sum_{i=1}^{+\infty} \lambda_n h_n \otimes h_n \right) = \sum_{n=1}^{+\infty} \langle L_\psi(\lambda_n h_n), h_n \rangle$$

and the last expression is exactly $\text{Tr}(L_\psi T_u)$ when we compute it using the basis h_n . \square

Lemma A.5. *Let us consider a Banach space ν_1 (respectively ν_2) continuously embedded in B_1^* (respectively B_2^*). Then $\bar{\chi} := \nu_1 \hat{\otimes}_\pi \nu_2$ can be continuously embedded in $(B_1 \hat{\otimes}_\pi B_2)^*$. In particular there exists a constant $C > 0$ such that, for all $u \in \bar{\chi}$,*

$$|u|_{(B_1 \hat{\otimes}_\pi B_2)^*} \leq C |u|_{\bar{\chi}}, \quad (61)$$

after having identified an element of $\bar{\chi}$ with an element of $(B_1 \hat{\otimes}_\pi B_2)^$ (see Lemma A.2). In other words $\bar{\chi}$ is a Chi-subspace of $(B_1 \hat{\otimes}_\pi B_2)^*$. In particular $B_1^* \hat{\otimes}_\pi B_2^*$ is a Chi-subspace of $(B_1 \hat{\otimes}_\pi B_2)^*$.*

Proof. To simplify the notations assume the norm of the injections $\nu_1 \hookrightarrow B_1^*$ and $\nu_2 \hookrightarrow B_2^*$ to be less or equal than 1. We remind that $(B_1 \hat{\otimes}_\pi B_2)^*$ is isometrically identified with the Banach space of the bilinear bounded forms from $B_1 \times B_2$ to \mathbb{R} , denoted by $\mathcal{B}i(B_1, B_2)$.

Consider first an element $u \in \bar{\chi}$ of the form $u = \sum_{i=1}^n a_i^* \otimes b_i^*$ for some $a_i^* \in \nu_1$ and $b_i^* \in \nu_2$. u can be identified with an element of $\mathcal{B}i(B_1, B_2)$ acting as $u(\phi, \psi) := \sum_{i=1}^n \langle a_i^*, \phi \rangle \langle b_i^*, \psi \rangle$. We can choose $a_i^* \in \nu_1$ and $b_i^* \in \nu_2$ such that $u = \sum_{i=1}^n a_i^* \otimes b_i^*$ and

$$|u|_{\bar{\chi}} = \inf \left\{ \sum_{i=1}^n |x_i|_{\nu_1} |y_i|_{\nu_2} : u = \sum_{i=1}^n x_i \otimes y_i, \quad x_i \in \nu_1, y_i \in \nu_2 \right\} > -\epsilon + \sum_{i=1}^n |a_i^*|_{\nu_1} |b_i^*|_{\nu_2}.$$

Using such an expression for u we have

$$\|u\|_{\mathcal{B}i(B_1, B_2)} = \sup_{|\phi|_{B_1}, |\psi|_{B_2} \leq 1} \left| \sum_{i=1}^n \langle a_i^*, \phi \rangle \langle b_i^*, \psi \rangle \right| \leq \sum_{i=1}^n |a_i^*|_{B_1^*} |b_i^*|_{B_2^*} \leq \sum_{i=1}^n |a_i^*|_{\nu_1} |b_i^*|_{\nu_2} \leq \epsilon + |u|_{\bar{\chi}}.$$

Since ϵ is arbitrary, we conclude that $\|u\|_{\mathcal{B}i(B_1, B_2)} \leq |u|_{\bar{\chi}}$.

Since this proves that the mapping that associates to $u \in \nu_1 \hat{\otimes}_\pi \nu_2$ its corresponding element in $\mathcal{B}i(B_1, B_2)$, has norm 1 on the dense subset $\nu_1 \hat{\otimes}_\pi \nu_2$, then the claim is proved. \square

Lemma A.6. *Let H_1, H_2 be two separable Hilbert spaces. Then $H_1^* \hat{\otimes}_\pi H_2^*$ is sequentially dense in $(H_1 \hat{\otimes}_\pi H_2)^*$ in the weak-* topology.*

Proof. Let (e_i^*) and (f_i) be respectively two orthonormal bases of H_1 and H_2 . We denote by \mathcal{D} the linear span of finite linear combinations of $e_i^* \otimes f_i$. Let $T \in (H_1 \hat{\otimes}_\pi H_2)^*$, which is a linear continuous functional on $H_1 \hat{\otimes}_\pi H_2$. Using the identification of $(H_1 \hat{\otimes}_\pi H_2)^*$ with $\mathcal{B}i(H_1, H_2)$, for each $n \in \mathbb{N}$, we define the bilinear form

$$T_n(a, b) := \sum_{i=1}^n \langle a, e_i \rangle_{H_1} \langle b, f_i \rangle_{H_2} T(e_i, f_i).$$

It defines an element of $H_1^* \hat{\otimes}_\pi H_2^* \subset (H_1 \hat{\otimes}_\pi H_2)^*$. It remains to show that

$${}_{(H_1 \hat{\otimes}_\pi H_2)^*} \langle T_n, l \rangle_{H_1 \hat{\otimes}_\pi H_2} \xrightarrow{n \rightarrow \infty} {}_{(H_1 \hat{\otimes}_\pi H_2)^*} \langle T, l \rangle_{H_1 \hat{\otimes}_\pi H_2}, \quad \text{for all } l \in H_1 \hat{\otimes}_\pi H_2.$$

We show now the following.

- (i) $T_n(a, b) \xrightarrow{n \rightarrow \infty} T(a, b)$ for all $a \in H_1, b \in H_2$.
- (ii) For a fixed $l \in H_1 \hat{\otimes}_\pi H_2$, the sequence $T_n(l)$ is bounded.

Let us prove first (i). Let $a \in H_1$ and $b \in H_2$. We write

$$T_n(a, b) = T \left(\sum_{i=1}^n \langle a, e_i \rangle_{H_1} e_i, \sum_{i=1}^n \langle b, f_i \rangle_{H_2} f_i \right). \quad (62)$$

Since $\sum_{i=1}^n \langle a, e_i \rangle e_i \xrightarrow{n \rightarrow +\infty} a$ in H_1 , $\sum_{i=1}^n \langle b, f_i \rangle f_i \xrightarrow{n \rightarrow +\infty} b$ in H_2 , and T is a bounded bilinear form, the point (i) follows.

Let us prove now (ii). Let $\epsilon > 0$ fixed and $l_0 \in \mathcal{D}$ such that $|l - l_0|_{H_1 \hat{\otimes}_\pi H_2} \leq \epsilon$. Then

$$|T_n(l)| \leq |T_n(l - l_0)| + |T_n(l_0)| \leq |T_n|_{(H_1 \hat{\otimes}_\pi H_2)^*} |l - l_0|_{H_1 \hat{\otimes}_\pi H_2} + |T_n(l_0)|. \quad (63)$$

So (63) is bounded by

$$\sup_{|a|_{H_1}, |b|_{H_2} \leq 1} \sum_{i=1}^n |\langle a, e_i \rangle e_i|_{H_1} |\langle b, f_i \rangle f_i|_{H_2} |T|_{(H_1 \hat{\otimes}_\pi H_2)^*} \epsilon + \sup_n |T_n(l_0)| \leq |T|_{(H_1 \hat{\otimes}_\pi H_2)^*} \epsilon + \sup_n |T_n(l_0)|,$$

recalling that the sequence $(T_n(l_0))$ is bounded, since it is convergent. Finally (ii) is also proved. At this point (i) implies that

$${}_{(H_1 \hat{\otimes}_\pi H_2)^*} \langle T_n, l \rangle_{H_1 \hat{\otimes}_\pi H_2} \xrightarrow{n \rightarrow \infty} {}_{(H_1 \hat{\otimes}_\pi H_2)^*} \langle T, l \rangle_{H_1 \hat{\otimes}_\pi H_2}, \quad \text{for all } l \in \mathcal{D}.$$

Since \mathcal{D} is dense in $H_1 \hat{\otimes}_\pi H_2$, the conclusion follows by Banach-Steinhaus theorem, see Theorem 18, Chapter II in [14]. \square

ACKNOWLEDGEMENTS. The authors are grateful to the Editor, Associated Editor and to both Referees for their valuable comments which helped them to considerably improve the first version of the paper. The research was partially supported by the ANR Project MASTERIE 2010 BLAN-0121-01 The research of the second named author also benefited partially from the support of the “FMJH Program Gaspard Monge in optimization and operation research” (Project 2014-1607H). It was partially written during the stay of the second named author in the Bernoulli Center (EPFL Lausanne) and at Bielefeld University, SFB 701 (Mathematik). The work of the first named author has been developed in the framework of the center of excellence LABEX MME-DII (ANR-11-LABX-0023-01).