

Plants of embalming in Modern Times: archaeobotanical results from the Jacobins' Convent at Rennes (Brittany, France)

Delphine Barbier-Pain, Marie-Pierre Ruas, Rémi Corbineau, Sylvie Duchesne,

Norbert Telmon, Rozenn Colleter

▶ To cite this version:

Delphine Barbier-Pain, Marie-Pierre Ruas, Rémi Corbineau, Sylvie Duchesne, Norbert Telmon, et al.. Plants of embalming in Modern Times: archaeobotanical results from the Jacobins' Convent at Rennes (Brittany, France). 17th Conference of the International workgroup for palaeoethnobotany, Jul 2016, Paris, France. , 2016. halshs-01740336

HAL Id: halshs-01740336 https://shs.hal.science/halshs-01740336

Submitted on 21 Mar 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PLANTS OF EMBALMING IN MODERN TIMES Archaeobotanical results from the Jacobins' Convent at Rennes (Brittany, France)

Delphine Barbier-Pain^{1, 2}, Marie-Pierre Ruas^{3, 4}, Rémi Corbineau², Sylvie Duchesne^{1, 5}, Norbert Telmon^{5, 6}, Rozenn Colleter^{1, 5}

A rescue excavation was conducted by the INRAP (2011-2013) in Les Jacobins' convent at Rennes (Brittany, France). About 800 burials including lead coffins dated to the 17th century have been uncovered together with 5 lead reliquaries (cardiotaphs); 4 of them had an inscription revealing the identity of the deceased.





al. 2012).





Lead coffin #1001: 3 lead cardiotaphs (CA) are visible at the head of the coffin, including heart #3

The anthropological analyses revealed marks of post-mortem manipulations of some of the skeletons: skulls were sawed and sterna were cut. In addition, compacted stuff were filling the cavities of any bodies, as well as the hearts in the cardiotaphs. These evidences testified embalming practices which were restricted to the social elites (Mafart *et al.* 2004; Georges 2006; Devriendt *et al.* 2004; Georges

MATERIAL AND METHODS

The burials which presented embalming evidence were sampled for several organic remains analyses at different locations in the lead coffin, in the different body regions and inside and outside the hearts in the cardiotaphs, as well. To date, 2 cardiotaphs (#3 & 5) and 1 coffin (1001) provided positive results for plant remains.

The sporo-pollinic material was extracted according to the current physic-chemical techniques, and analysed with a transmission photonic microscope. Subfossil plant macro-remains, dry, charred or mineralized, have been sampled without sieving and directly sorted in water under the stereomicroscope.

MAIN PLANT INGREDIENTS

The different preserved plant remains (pollen, flowers, stems, twigs, leaves, fruits or seeds) which are the embalming ingredients revealed a high taxonomic diversity (70 taxa). Hearts #3 & 5 provided 59 more or less precisely identified taxa.

Some of these plants were used because of the aromatic and antiseptic properties of their organs. They are frequently mentioned in the 17th century recipe treatises (Ruas 1992; Corbineau 2014): **blooming plant** of **mugwort, camomile, oregano, lesser calamint, lavender, thyme** and **rosemary leaves, fennel** and **juniper fruits**.



nent of twig and leaf: Th

- Rosmarinus leaf fragment - Lamiaceae calyx HEART #3 DRIED STUFF



Fragments of cloves, a common spice also used in the recipes, are the only exotic plant identified at Les Jacobins (Heart #3). *Myrtus-type* pollen may also come from flowers imported from the Mediterranean region. Flax tow was filling the cavity of the cardiotaph and the hearts. Charcoal (Angiospermae) testified the use of ashes.

Two different recipes were used for preparing the dough filling the two hearts. With 6 different morphotypes, the Lamiaceae largely predominated the pollen spectrum of heart #5 (80%); the latter mostly provided fennel seeds and lavender flowers. Heart #3 mainly yielded Asteraceae pollen (*Anthemis*-type), together with lesser proportions of Myrtaceae (*Myrtus*-type) and Lamiaceae (unident.); macro remains analyses evidenced camomile flowers and rosemary leaves.

Much noticeable is the over-representation (>80%) of **Beta-type** pollen in burial #1001. Conversely, there is not any mention of Amaranthaceae among ca. hundred plant ingredients which are mentioned in the 17 pieces of texts that we analysed (surgery and pharmacological encyclopaedias, and 14-19th c. embalming reports (Corbineau 2014). This burial opens new reflection avenues about the funerary practices during the Modern times.

Fragments of floral bud

MAIN PLANTS RECORDED BY MACRO-REMAINS & POLLEN HEARTS 17 th C. AD	PART OF EMBALMING PLANTS				
	VEGETATIVE		FLOWER		FRUIT
	LEAF	TWIG STEM FIBRE	BUD	BLOOMING	RIPE
Rosmarinus officinalis					
Thymus vulgaris					
Origanum vulgare					
cf. Linum usitatissimum					
Syzygium aromaticum					
cf. Lavandula					
cf. Myrtus					
Clinopodium nepeta					
Apiaceae, Asteraceae & Lamiaceae					
Matricaria chamomilla					
Artemisia vulgaris					
_ Mentha longifolia					
Foeniculum vulgare					
Juniperus communis/oxycedrus					

1- National Institute of Preventive Archaeological Research (INRAP)

2- UMR 6566, CReAAH, Rennes-Nantes, France

3- National Center of Scientific Research (CNRS) 4 - UMR 7209 AASPE, Muséum nationald'Histoire naturelle, Sorbonne-Universités, Paris, France

UMR 5288, Laboratory for Molecular Anthropology and Computer-generated Imagery (AMIS), Toulouse, France

6- Medico legal department, CHU Toulouse-Rangueil, Toulouse, France



<u>2.5 mm</u> <u>0.5 mm</u> Fioral organs & achene: Matricana chamomilia Plant macro-remains sampled from the infilling of two hearts. The size of some fragments suggests that the dried plants were crushed and coarsely cut (size 1-2 mm). Oils and resins mixed with the plant powder still strongly smelt during our analysis.

REFERENCES

Corbineau R. 2014. Pour une archéobotanique funéraire enquêtes interdisciplinaires et analyses polliniques autour de la tombe et du corps mort (ére chrétienne, France - Italie), Ithèse doctorat, Université du Maine, 595 p. Devriendt W., Venet S., Defnée A., Garnier N., Gillet B., Rossetti L. 2012. Découverte d'un cœur reliquaire à Douai (ancienne église Saint-Jacques, place Carnot) : approche pluridisciplinaire de l'embaumement à l'époque moderne, Archéologie Médlévale 42 : 23-42.

Sacques, pade Carlon). - approche punctuscipinatare de reinidadmiententi à repoque moderne, Anciencoigne Medicavale 42, 25-42. Georges P. 2006. L'embaumement et le préférement du coer au Moyen Age. Le sociage du stermum, in: CHARLER (P) (dir.) – Actes du 1er colloque international de Pathographie, Loches, 22-24 avril 2005, Paris, éditions de Boccard, 2006, pp. 99-112 Mafart B. Pelletier J.-P. Erron M. 2004. Destamorte ambient de la méticaval funcerary tradice. A case observed at the cometery of

Mafart B., Pelletier J.-P., Fixot M. 2004. Post-mortem ablation of the heart: a medieval funerary practice. A case observed at the cemetery of Ganagobie Priory in the French Department of Alpes-de-Haute-Provence, International Journal of Osteoarchaeology 14: 67-73... Ruas M.-P. 1932. Matières d'embaumement dans la sépulture du Château de Laval : analyse des graines, La Mayenne: Archéologie, Histoire, supplément 2, 1992 : 87-91.