



**HAL**  
open science

## Small headwater stream evolution in response to Lateglacial and Early Holocene climatic changes and geomorphological features in the Saint-Gond marshes (Paris Basin, France)

Bruno Depreux, Amélie Quiquerez, Carole Bégeot, Christian Camerlynck,  
Anne-Véronique Walter-Simonnet, Pascale Ruffaldi, Rémi Martineau

### ► To cite this version:

Bruno Depreux, Amélie Quiquerez, Carole Bégeot, Christian Camerlynck, Anne-Véronique Walter-Simonnet, et al.. Small headwater stream evolution in response to Lateglacial and Early Holocene climatic changes and geomorphological features in the Saint-Gond marshes (Paris Basin, France). *Geomorphology*, 2019, 345, pp.106830. 10.1016/j.geomorph.2019.07.017 . halshs-02294136

**HAL Id: halshs-02294136**

**<https://shs.hal.science/halshs-02294136>**

Submitted on 5 Nov 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NoDerivatives 4.0 International License

# Small headwater stream evolution in response to Lateglacial and Early Holocene climatic changes and geomorphological features in the Saint-Gond marshes (Paris Basin, France)

Bruno Depreux, Amélie Quiquerez, Carole Bégeot, Christian Camerlynck, Anne-Véronique Walter-Simonnet, Pascale Ruffaldi, Rémi Martineau

**This is an author generated version of the article:**

Bruno Depreux, Amélie Quiquerez, Carole Bégeot, Christian Camerlynck, Anne-Véronique Walter-Simonnet, Pascale Ruffaldi, Rémi Martineau (2019). Small headwater stream evolution in response to Lateglacial and Early Holocene climatic changes and geomorphological features in the Saint-Gond marshes (Paris Basin, France), *Geomorphology*, Volume 345, 106830, ISSN 0169-555X.

Copyright: Elsevier (<https://www.elsevier.com>)

The final publication is available on <https://doi.org/10.1016/j.geomorph.2019.07.017>

1 **Small headwater stream evolution in response to Lateglacial and Early Holocene climatic changes and**  
2 **geomorphological features in the Saint-Gond marshes (Paris Basin, France).**

3 Bruno Depreux <sup>a,\*</sup>, Amélie Quiquerez <sup>b</sup>, Carole Bégeot <sup>c</sup>, Christian Camerlynck <sup>d</sup>, Anne-Véronique Walter-Simonnet <sup>c</sup>,  
4 Pascale Ruffaldi <sup>c</sup>, Rémi Martineau <sup>b</sup>

5 <sup>a</sup> Université Paul Valéry Montpellier 3, UMR 5140 ASM, CNRS, MCC, 34199 Montpellier, France

6 <sup>b</sup> Université de Bourgogne Franche-Comté, UMR 6298 ARTEHIS, CNRS, 21000 Dijon, France

7 <sup>c</sup> Université de Bourgogne Franche-Comté, UMR 6249 Chrono-Environnement, CNRS, 25030 Besançon, France

8 <sup>d</sup> Université Pierre et Marie Curie Paris 6, UMR 7619 METIS, CNRS, 75252 Paris, France

9  
10 \* Corresponding author:

11 E-mail address: [bruno.depreux@univ-montp3.fr](mailto:bruno.depreux@univ-montp3.fr)

12 Full postal address: Bruno Depreux, Université Paul Valéry Montpellier 3, Route de Mende, UMR 5140 –  
13 ASM, 34199 Montpellier cedex 05, France

14  
15 **Abstract**

16 The study focuses on river dynamics and vegetation changes during the last Glacial-Interglacial transition from a  
17 headwater stream located in the eastern part of the Paris Basin. We adopt a spatial multiproxy approach combining  
18 image analyses, geophysical surveys, sedimentary and pollen record analyses to document the impact of Lateglacial  
19 climate changes and geomorphological features on vegetation and fluvial dynamics in the small Boitet catchment (ca.  
20 20 km<sup>2</sup>). Our results show that the sedimentary record is organized into six successive alluvial sequences reflecting  
21 changes in discharge and channel morphology in response to short phases of climate oscillations during the Lateglacial  
22 and Early Holocene periods. The Boitet alluvial sequences present some similarities with other NW European rivers  
23 that are interpreted as large-scale fluvial system evolution to climate changes. However, some local differences have  
24 also been highlighted partly related to the upstream position of the catchment. Among them, two distinctive features  
25 of the Boitet catchment are 1) the preservation of the Oldest Dryas deposits, which have been rarely described in this  
26 area, and 2) the recognition of multichannel river dynamics during the Oldest Dryas and Younger Dryas. We  
27 demonstrate that the fluvial evolution is firstly triggered by climate changes and that land surface features may also  
28 influence specifically upstream areas revealing contrasting responses of the river system.

29 **Keywords:** Lateglacial; anastomosed system; headwater catchment; climate changes; Saint-Gond marshes; Paris Basin

## 31 1 Introduction

32 In NW Europe, the Last Glacial Maximum (LGM), reached at ca.  $22.1 \pm 4.3$  ka, was followed by a Lateglacial interstadial  
33 during which glaciers retreated and temperatures increased (Shakun and Carlson, 2010; Clark et al., 2012). Over the  
34 last three decades, numerous paleoclimate records based on marine and continental sedimentary archives from the  
35 North Atlantic realm, indicate a non-linear warming punctuated by prominent abrupt climate changes (Björck et al.,  
36 1996; NGRIP members, 2004). The failure of the North Atlantic thermohaline circulation caused by freshwater inputs  
37 is supposedly partly responsible for these cooling events (Barber et al., 1999; Teller et al., 2002; Alley and Agustsdottir,  
38 2005; Fleitmann et al., 2008), and these have been well documented in the Greenland ice cores. At the transition from  
39 the LGM to the Lateglacial, analyses of oxygen isotope ratios in ice cores have allowed the identification of two stadials  
40 associated with cold conditions (GS-2 and GS-1), interspersed with one interstadial displaying milder conditions (GI-1)  
41 (Björck et al., 1998). In terrestrial records from northern Europe, these abrupt cooling events were firstly recognised  
42 from pollen analyses by Jessen (1935) and Iversen (1954). An inferred Lateglacial biostratigraphy zonation was defined  
43 by a conventional sequence (Oldest Dryas, Bølling, Allerød and Younger Dryas), which is often used in a  
44 chronostratigraphic and/or climatostratigraphic sense (Litt et al., 2001; Rasmussen et al., 2014). Sedimentary records  
45 from European rivers have been widely studied in order to document these periods, as in northern France (Antoine et  
46 al., 2003; Pastre et al., 2003; Deschodt et al., 2004), but also in the Netherlands (Vandenberghe et al., 1987), in the  
47 United Kingdom (Brown, 2001; Bridgland, 2010; Macklin et al., 2010; Walker et al., 2012), in Germany (Kasse et al.,  
48 2005), or in Poland (Vandenberghe et al., 1994). They all highlight that NW European river morphodynamics evolve in  
49 response to the short-term climate changes and follow broadly similar fluvial evolution patterns. During the colder  
50 period, e.g. during the Upper Pleniglacial or Younger Dryas period (respectively GS-2 and GS-1), fluvial dynamics are  
51 generally characterized by the development of braided channels in a steppe environment (Vandenberghe, 2003).  
52 Conversely, a warmer climate, e.g. the Bølling-Allerød (GI-1), induces a water discharge decrease and a temperature  
53 increase and favours conditions for organic deposition in channels and floodplains of NW European rivers (Antoine,  
54 1997; Pastre et al., 2000). However, if isotopic analyses on ice cores provide direct records of climatic oscillations,  
55 sedimentary archives are rather controlled by local hydrological events, which means that climate changes are  
56 indirectly recorded in the sedimentary archives, with possibly some delays and under various preservation potential  
57 conditions. This also implies that the chronology defined by the reference conventional sequence may not be entirely

58 preserved, and if preserved may be temporally shifted. For instance, among all of the studies performed in NW  
59 European rivers, only a very few of them have been able to describe Oldest Dryas deposits even though it is a well-  
60 known cooling climate period in ice cores. Therefore, there is a need to study environmental and fluvial  
61 morphodynamics changes in various catchments throughout Europe to better conceive the responses of rivers to  
62 climate changes and the spatial variability of the conventional chronostratigraphic sequence.

63 In this paper, we document the morphosedimentary dynamics and the vegetation cover evolution during the last  
64 Glacial-Interglacial transition of a small headwater basin located in eastern part of the Paris Basin (Saint-Gond marshes,  
65 NE France) where no previous studies have been carried out. We first present our results obtained from a  
66 multidisciplinary approach combining both spatial and temporal data: sedimentological and pollen analyses  
67 supplemented by aerial image processes and sub-surface geophysical surveys. Then, considering the  
68 geomorphological context and climate reconstructions from other alluvial sequences in NW Europe, we discuss the  
69 preservation potential of the sedimentary archives and the local geomorphological features as key factors for  
70 recording Lateglacial alluvial deposits.


## 72 2 Study area

73 The Boitet catchment (48°846 N, 3°92 E) belongs to a large peatland called “Marais de Saint-Gond”, located in the  
74 eastern part of the Paris Basin (Marne, France), at the foothills of the Brie plateau (Fig. 1). These peatlands have  
75 developed over a Late Cretaceous chalky substrate in the alluvial valley of the Petit-Morin River which is the trunk  
76 drainage of the catchment. The Boitet River is a small low-gradient tributary river draining a 20 km<sup>2</sup> catchment in the  
77 upstream part of the Saint-Gond marshes. The present-day mean slope is less than 1 ‰ and the mean annual water  
78 discharge is estimated at 2 m<sup>3</sup>.s<sup>-1</sup> in the Montmirail hydrological station, which is located 20 km downstream.

79 The Saint-Gond marshes are bordered to the west by an upland plateau characterized by a cuesta morphology where  
80 chalky slopes are covered by Mesozoic and Cenozoic deposits, composed of clays, silts and sands. The present-day  
81 morphology of the investigated catchment has been shaped during the Quaternary period. Periglacial processes  
82 favoured the weathering of the chalky substrate, delivering sediments consisting of chalky cryoclasts and flints to the  
83 valley floor. On the hillslopes, Cenozoic formations are overlain by colluvial deposits containing chalky cryoclasts  
84 packed into a silty matrix. Close to the peatlands, the core drillings performed by the French geological survey (BRGM)  
85 indicate that the chalky substrate is covered by Weichselian colluvial-alluvial deposits, whose thickness may reach up

86 to 10-15 m, and that are related to a terrace system associated with the Petit-Morin drainage network (Hatrival et al.,  
87 1988) (Fig. 1). During the Pleistocene, the drainage basin of the Saint-Gond marshes was modified to its eastern and  
88 upstream part by the hydrographic captures of the Somme-Soude and the Vaure rivers. These captures have induced  
89 a major disruption causing possibly a water discharge decrease and the alluvial aggradation of the Petit-Morin valley  
90 (Tricart, 1949).

91 Since the Pleniglacial, the Petit-Morin catchment did not experience any other significant landscape transformation  
92 apart from those let by anthropogenic activities. The surface of these peatbogs were broadly exploited from the 17<sup>th</sup>  
93 century as combustible fuel or fertilizer, as evidenced by the numerous traces of extraction pits of several meter thick  
94 (Salaün and Marre, 2005). Large drainage operations were carried out. The hydrographic network has been modified  
95 and regularly cleaned. As a result, in some areas, these anthropogenic activities have possibly provoked the  
96 destruction of the superficial layers of sedimentary sequence. Therefore, it is expected that Petit-Morin catchment  
97 hosts well-preserved sedimentary records for the Late Pleniglacial-Holocene transition period.



98  
99 Fig. 1. Study area location maps. A: Area of interest in France. B: Topographical map of the Saint-Gond marshes region and location of the study  
100 area at the mouth of the Boitet River. C: Location map of the coring transects (yellow circles) and the geophysical profiles (blue lines).

### 3 Material and Methods

The morphosedimentary dynamics of the valley has been investigated by combining geophysics, geomorphology, sedimentology and palynology in the central part of the Boitet valley.

#### 3.1 Planimetric analysis of the floodplain from image processing

A geomorphological study combining field surveys (core drilling and geophysical investigations) with image analysis techniques have been performed in order to obtain a synoptic view of the morphodynamics of the valley. We have used historical aerial images supplied by the French Geographic National Institute (IGN) (1946, 1969 and 1974) and historical maps, i.e., the cadastral plan called "Napoleonic cadastre" (1830) and the map of "Etat-Major" produced by the French army (1832). Images and maps have been georeferenced and also processed with ENVI/IDL (2018) remote sensing software to better reveal spectral anomalies linked to alluvial formations. We have reconstructed a diachronic image in which each spectral band is composed respectively of the aerial images of 1946, 1969 and 1974. Then, a Principal Component Analyses (PCA) have been performed to highlight spectral anomalies linked to channel systems, and then digitize the detected traces (Fig. 3).

#### 3.2 Geophysical investigations

We used a shallow geophysical method to reconstruct the geometry and thickness of the fluvial depositional system and to define the drilling location. To this end, four electrical resistivity tomography (ERT) profiles have been undertaken using a multimode resistivity imaging system (Syscal Pro Switch, Iris Instruments©). This method is widely employed in alluvial system (Vandenberghe and Desmedt, 1979; Vandenberghe, 1981; Van Huissteden et al., 1986; Chambers et al., 2012; Laigre et al., 2012; Hausmann et al., 2013; Rey et al., 2013; Matys Grygar et al., 2016; Bábek et al., 2018). The survey positioning was chosen in such a way that the paleochannels identified by aerial imagery were intersected. The profiles 1-3 were acquired perpendicularly to the river flow direction in order to image the sedimentary filling of alluvial valley where the valley was the widest (Fig. 4). We have positioned a longitudinal profile, parallel to the valley axis, to investigate spatial lateral variation of the sedimentary records. The 192 m long profiles were acquired surveys with 2 m-electrode spacing using a Wenner-Schlumberger array to maintain both a sufficiently high horizontal resolution and restrict sensitivity to vertical variations. Investigation depths, that reach 20 m for the profiles 1-3, respectively 8 m for the profile 4, are deep enough to attain the horizontal calcareous substrate and reconstruct the geometry of the alluvial record of the valley.

### 3.3 Core sedimentology

A total of eight cores, named MDV-1 to MDV-5, MDV-9, MDV-10 and MDV-12, were collected as close as possible to the geophysical profiles. However, for reasons of accessibility in the field, the geophysical profile and the coring locations were sometimes slightly shifted (Fig. 1). The cross-section consists of six cores, spaced 20 m apart in the central part, and spaced 40 m apart near the edges (Fig. 6).

Sediment cores were described on the basis of lithology and sediment grain-size characteristics. Lithology is dominated by the presence of clayey, silty chalky layers in which some cherts were identified and peaty sediments. According to the Udden-Wentworth Scale, sediment grain size ranges from clays, to silt to silty sand to pebbles. The sediment color, on the basis on the Munsell Colour Chart, can reflect either the lithology and/or the degradation of the organic content, and was used as a complementary parameter to define sedimentary facies. Using these criteria, ten major sedimentary facies (F1 to F10) were identified with their characteristics summarized in Table 1.

Table 1. Synthetic classification of the sedimentary facies and units of the Boitet River.

Sedimentary Unit (SU)	Sedimentary facies	Texture	Munsell color
SU 1	F1	clayey sand	2.5 Y 8/4 pale yellow
SU 2	F2	silty clay	10 YR 7/1 light gray
	F3	silty sand	10 YR 7/1 light gray
	F4	clayey-sandy silt	2.5 Y 8/1 white
	F5	silty-sandy clay	2.5 Y 6/4 light yellowish brown
	F10	sandy silt	10 YR 6/3 pale brown
SU 3	F6	clayey peat	10 YR 2/1 black
	F2	silty clay	10 YR 7/1 light gray
	F8	silty clay	7.5 YR 3/3 dark brown
SU 4	F2	silty clay	10 YR 7/1 light gray
	F3	silty sand	10 YR 7/1 light gray
	F7	sandy silt	10 YR 5/2 grayish brown
	F8	silty clay	7.5 YR 3/3 dark brown
SU 5	F6	clayey peat	10 YR 2/1 black
	F8	silty clay	7.5 YR 3/3 dark brown
SU 6	F9	silty clay	2.5 Y 2.5/1 black

#### 3.3.1 Laboratory sedimentary analyses

A maximum thickness of 4 m was obtained on MDV-4 in the central part of the Boitet valley. Therefore, we have chosen two reference cores (MDV-4 and MDV-1 near the geophysical profile 1), to be representative of the hydro-



144 sedimentary dynamics of the river and on which complementary laboratory analyses (magnetic susceptibility, loss on  
145 ignition, pollinic spectra) were performed.

146 Magnetic susceptibility was measured in SI unit with Bartington MS2E sensor, to detect any presence of detrital  
147 colluvial materials (Vanni re et al., 2004). Organic matter or calcite will lead to very low negative values, oxides and  
148 clays will have slightly positive values and ferrimagnetic materials, such as iron oxides will display high values (Dearing,  
149 1999).


150 Loss on ignition (LOI) method was used in order to determine organic matter (OM) and carbonate contents. The values  
151 of these contents are expressed in percentage of the mass of bulk. Samples were dried 12h at 105 C then burned in a  
152 muffle furnace in two steps. First, OM content was measured under oxidising conditions at 550 C during 5h. Then,  
153 carbonate content was obtained under pyrolysis conditions at 950 C during 2h (Heiri et al., 2001).

### 154 3.3.2 Chronostratigraphic framework

155 Seven <sup>14</sup>C ages were carried out on three samples in the MDV-4 core and four samples in the MDV-1 core, using AMS  
156 method in organic levels (Table 2) to propose chronostratigraphic framework and constrain the temporal evolution of  
157 the depositional environment. The age-depth model was established on organic material using the R package 'clam'  
158 (Fig. 2). No material was available to date the bottom of the sequence. Calibration was performed by using the CALIB  
159 programme version 7.1 (Stuiver and Reimer, 1993) with the IntCal13 calibration curve (Reimer et al., 2013). In the  
160 MDV-1, <sup>14</sup>C ages extend from 12460 ± 70 BP (15024-14220 cal. BP) at a depth of 191 cm to 10220 ± 50 BP (12131-  
161 11752 cal. BP) at the depth of 70 cm. In the MDV-4, the older <sup>14</sup>C dating is of 12140 ± 70 BP (14184-13779 cal. BP) at  
162 about 127 cm, and the younger <sup>14</sup>C dating is of 10110 ± 60 BP (11993-11400 cal. BP).

163 Thus, these <sup>14</sup>C ages span the Lateglacial period, from the Oldest Dryas – B lling transition to the Younger Dryas –  
164 Preboreal transition. In this paper, we employ the term 'Oldest Dryas', which encompass ca. 18 – 14.7 cal. BP, as a  
165 chronostratigraphic period, such as the associated conventional nomenclature (B lling, Aller d and Younger Dryas)  
166 (Alley and Clark, 1999; Ivy-Ochs et al., 2005; Shakun and Carlson, 2010; Clark et al., 2012).

167



168

169

Fig. 2. Age-depth model of the MDV1 core. Age spectra (grey shading) correspond to the 2σ ranges.

170

171

Table 2. AMS radiocarbon dates from the Boitet River. Calibration with IntCal13 (Reimer et al., 2013).

Lab. code	Core	Depth (cm)	<sup>14</sup> C yr BP	Cal. BP (2σ)	Biozones
Poz-70902	MDV-4	68-69	10110 ± 60	11993-11400	Younger Dryas
Poz-70901	MDV-4	91.5-92.5	10890 ± 60	12905-12688	Allerød
Poz-70903	MDV-4	126.5-127.5	12140 ± 70	14184-13779	Bølling
Poz-86926	MDV-1	70	10220 ± 50	12131-11752	Younger Dryas
Poz-86927	MDV-1	127	10560 ± 50	12661-12410	Younger Dryas
Poz-86928	MDV-1	159	10970 ± 60	12993-12718	Allerød
Poz-70904	MDV-1	191-192	12460 ± 70	15024-14220	Bølling

172

173

### 3.4 Palynology

174

Pollen analysis was carried out on 60 samples collected at four centimetres intervals from the MDV1 core. Around 1 g

175

of fresh matter was prepared using the standard method established by Faegri and Iversen (1989). Material was

176

filtered to 200 μm, then treated with HCl 10%, HF 40%, NaOH 10% and acetolysis to colour the pollen grains. A

177

minimum of 300 pollen grains were counted from each sample. Pollen grains were identified using a light microscope

178

(magnification 250x and 400x) with comparison to modern material and the standard identification keys of the Central

179 European pollen flora (Beug, 1961; Punt, 1976; Punt and Clarke, 1980, 1981, 1984; Punt et al., 1988; Faegri and Iversen,  
180 1989; Moore et al., 1991; Reille, 1992, 1995).

181 The construction of the pollen diagram was performed using the program TILIA 1.5.12 (Grimm, 1991). The main pollen  
182 sum includes all pollen types excluding spores. Six pollen assemblage zones (PAZ) were identified according to the  
183 changes on trends in the percentages of the main taxa.

## 184 4 Results

### 185 4.1 Mapping the traces of paleochannels from aerial images and geophysical investigation

186 Aerial image processes reveal the presence of numerous linear and sinuous spectral anomalies in the floodplain,  
187 displaying straight, to sinuous, to meandering geometries and that may intersect themselves. These detected  
188 anomalies could be indicative of an alluvial history of the valley before the development of the marshes (Fig. 3) and  
189 are interpreted as remnant palaeochannels. To test this hypothesis, the positioning of geophysical profiles 1 to 4 were  
190 chosen in such a way that some of the possible palaeochannels identified by aerial image processing were crossed,  
191 and this, where field condition accessibility in the marsh was possible (dense shrub vegetation, open water patches).

192 The low resistivity values, ranging from 15 to 107  $\Omega$ .m, suggest a mixture of clay, silt, and sand deposits within a water-  
193 saturated context (Fig. 4). In all ERT profiles, it is possible to distinguish a basal homogeneous unit defined by a tabular  
194 geometry and the relatively highest resistivity values ( $> 50 \Omega$ . m). This basal unit is overlain by a 1-4 m thick  
195 heterogeneous unit, displaying the lowest resistivity values ( $< 20$  to  $50 \Omega$ .m). This unit exhibits some gully  
196 morphologies, symmetrical or asymmetrical, that can be recognized by their low resistivity ( $< 35 \Omega$ .m).





Fig. 3. Network of multiple palaeochannels revealed by the PCA of the historical aerial images performed in 1946, 1969 and 1974.

In this alluvial context, the geometries and the resistivity values of the basal unit are consistent with deposits, composed of gravel, sand and silt. The low-resistivity gully morphologies identified in the upper geophysical unit are located just below the traces of palaeochannels that were detected by the aerial images. The geometries and low resistivity values are consistent with paleochannels filled by organic and silty-clayey deposits.

The results of photo-interpretation and geophysics highlight a fluvial system dominated by multiple channels exhibiting various shapes, ranging from meander to slightly sinuous to straight channel. Morphology of the channels has been estimated by the width/depth ratio that is indicative of the energy of the river regime. In our case, we have estimated these values on the conductive sedimentary bodies, identified in the geophysical profiles for depth of less than 2 m. The Wenner-Schlumberger array, as it sounds the sub-surface horizontally and vertically, makes it possible to assess the channel morphology, and give some estimation of the depth by the contrast between high and low

209 resistivity values. Smith and Smith (1980) have shown that values ranging from 7 to 20 are indicative of an  
210 anastomosed fluvial system.



211

212 Fig. 4. Reconstruction of the Lateglacial multichannel river network from geophysical and coring surveys.

#### 213 4.2 Sedimentology, palynology and chronostratigraphic framework

214 Here, we describe the sedimentary and palynological record subdivided respectively into six sedimentary units (SU)  
215 and five pollinic units (PAZ) on the basis of (i) the ten sedimentary facies defined previously, (ii) the sedimentary  
216 analyses (grain size, magnetic susceptibility, carbonate and organic contents), (iii) the palynological analyses and (iv)  
217 the age constraints. The chronostratigraphic framework was constrained by  $^{14}\text{C}$  ages, by relative chronology between  
218 the different units, and/or by comparison to facies defined by the BRGM (Hatrival et al., 1988). The depositional  
219 geometry of the cross-section has been reconstructed from the correlation of six SU.


#### 220 Unit 1 (SU 1 / PAZ 1)

221 *Sedimentology:* The base of the sedimentary record begins with a yellow oxidised clayey sand (F1) overlaying the  
222 Campanian chinks. This bed contains angular chalky gravel layer and abundant flints, coming from the erosion of  
223 Campanian chalky bedrock. The top of SU is delimited by an undulating erosional surface, forming a topographic  
224 depression of more than 1 m between the MDV-3 and MDV-2 cores (Fig. 66).

225 *Palynology*: This zone corresponds to a very low pollen concentration with much of the pollen being eroded and not  
226 identifiable. Levels at 246 and 250 cm depth do not contain any pollen. The spectra is dominated by Poaceae,  
227 Cyperaceae, *Betula* and *Juniperus* (Fig. 77).

228 *Chronostratigraphic constraints*: A lack of organic matter does not permit any  $^{14}\text{C}$  dating of Unit 1. However, since this  
229 unit displays similar characteristics to the Weichselian facies described by the BRGM (Hatrival et al., 1988), we propose  
230 to attribute Unit 1 to the Late Pleniglacial period.

231 *Interpretation*: The coarse texture of the deposits and the presence of abundant flint slivers suggest a periglacial  
232 environment dominated by cryoclastic processes and a sediment transport governed by high-energy braided rivers.  
233 The incision phase results from a change in the river discharge occurring at the end of the Pleniglacial period.




234  
235 Fig. 5. Magnetic susceptibility, organic matter and carbonates contents from MDV4 core.

## 236 Unit 2 (SU 2 / PAZ 2)

237 *Sedimentology*: The thickness and the facies of the SU 2, overlaying the erosional surface, differ from one core to the  
238 other (Fig. 66). The unit is the thickest in the MDV-4 core (> 3 m thick since the SU 1 has not been reached) and the  
239 thinnest in the MDV-1 core (only about 50 cm). In all of the cores where the unit base was attained, the basal deposits  
240 consist of yellowish-grey silty clay beds (F2) that have weakly positive magnetic susceptibility values ( $1$  to  $5 \times 10^{-5}$  SI).  
241 In the lowermost part of this unit, some greyish silty sand beds (F3) displaying a null to weakly negative signal ( $-3$  to  
242  $50 \times 10^{-5}$  SI) have been encountered in the MDV-3 and MDV-4 cores (Fig. 5). In the upper deposits of the unit, several



243 facies evolutions have been observed. In the MDV-2, MDV-1 and MDV-12 cores, some colour changes, from yellow-  
 244 brown to grey-blue towards the top, suggest an evolution towards a reducing environment. At the uppermost part of  
 245 the cores, some thin organic-rich beds have been encountered. In these cores, the transition towards Unit 3 is gradual.  
 246 Conversely, the MDV-5 core consists of a whitish clayey-sandy silt facies (F4). In the MDV-4 and MDV-5 cores, no  
 247 increase in the organic content has been observed and the uppermost part of the unit is marked by a second incision  
 248 surface that also was identified in the MDV-3 core. Sedimentological analyses show that SU 2 displays a very low  
 249 organic matter content (about 4 %), and a high content of silicates and oxides (63 %). The carbonate content represents  
 250 about a third of the total composition (Fig. 5).



251

252

Fig. 6. Cross-section of the Lateglacial and Early Holocene sequence from the Boitet valley with the sedimentary units and the radiocarbon dates.

253 *Palynology*: The pollen composition of this zone indicates a large open herbaceous landscape with dominance of sedge  
254 communities, probably linked to swamp soils where important shrub vegetation with willow and birch can grow (Fig.  
255 77). Standing water is assumed because of consequent pollen percentages of aquatic plants at the end of the zone,  
256 between 222 cm and 198 cm. The extension of local dry habitats was limited as it was reflected by low representation  
257 of steppe vegetation but nevertheless more significant at the beginning of the zone with the development of *Ephedra*  
258 and *Hippophae*.

259 *Chronostratigraphic constraints*: The radiocarbon dates performed at the base of the Unit 3 corresponds to the onset  
260 of the Bølling. It is likely that Unit 2 records the Oldest Dryas period and partly an early phase of the Bølling.

261 *Interpretation*: This assumption seems to be confirmed by the pollen assemblages of treeless vegetation which can be  
262 compared with the Oldest Dryas of the Lateglacial stratigraphy of higher altitude areas although the percentages of  
263 *Artemisia* were relatively low. The silty-clayey sequence suggests a gradual filling of the channels by a fine alluvial load.  
264 A significant river and sediment discharge change has possibly occurred between the Pleniglacial period and the Oldest  
265 Dryas to explain the absence of a gradual evolution of the sediment supply from coarse to fine load. The high silt/clay  
266 percentage, the low width/depth ratio and the aggradation geometry of the deposits are all consistent with a low-  
267 energy system of anastomosed channels. We note a peculiarity of channel III: where the whitish clayey-sandy silt (F4)  
268 of the MDV-5 core, which is also observed in MDV-9 further upstream, could correspond either to lateral bank deposits  
269 or to the infilling of another channel. This singular deposit may correspond to a distinctive channel at the east of the  
270 alluvial plain, as these cores are exactly located on a wide meandering channel observed on aerial images and  
271 geophysical surveys (Fig. 3). Nevertheless, we are unable to link this channel to the chronostratigraphic framework to  
272 fully confirm this hypothesis.

273 In the MDV-1 and MDV-2, the colour, evolving from yellow-brown to grey-blue towards the top, is indicative of a  
274 transformation from oxidised towards reducing conditions. The reducing environment, the preservation of muddy  
275 organic deposits and the gradual facies transition towards the Units 3 suggest that these muddy organic-rich deposits  
276 accumulated in a channel that was abandoned. In contrast, MDV-4 and MDV-5 document the functioning of active  
277 channels. On the whole, our data suggest that the landscape is a marshy expanse with a low-energy multichannel  
278 system running through it with some abandoned and active channels. These hypotheses are consistent with  
279 geophysical and image data that highlight a multichannel river system, although it cannot be totally excluded that  
280 some of the channels in the floodplain are more recent.



282 *Sedimentology:* The base of the unit presents some differences of facies and thicknesses between the abandoned  
283 channel (MDV-1 and MDV-2) and the active channels (MDV-3, MDV-4 and MDV-5). The peat thickness varies between  
284 the cores: the thickness is highest in the MDV-1 and MDV-2 cores, and thinnest in the MDV-5 and MDV-3 cores (Fig.  
285 66). In the MDV-1 and MDV-2, the unit starts with thin organic centimetre thick layers that are interbedded with silty  
286 beds. These deposits evolve towards a highly-decomposed black clayey peat (F6) that fills the space available left by a  
287 channelized topography. In the MDV-3, MDV-4 and MDV-5, no silty beds have been observed in the lowermost part  
288 of the unit, leaving only the black peaty layer (F6) that overlays directly the incision surface. This facies is composed of  
289 75 % of organic matter, 2-3 % of carbonate content and 20 % of silicates and oxides (Fig. 5). The susceptibility values  
290 are weakly negative to null, which is consistent with the presence of a high organic matter content. The top of the unit  
291 is truncated by a sharp undulating erosional surface that is observed in all cores.


292 *Palynology:* This zone corresponds to the pioneer shrub colonization (PAZ3a) which continues with the installation of  
293 pine (PAZ3b) in or near the investigated area (Fig. 77). In PAZ3a, the first stage of dynamic is characterized by an  
294 increase of *Juniperus* rapidly followed by a rise in *Betula*. *Salix* communities remain locally in the surrounding wet  
295 areas. The depletion of pollen from aquatic plants with simultaneous and continuous increase of hygrophilous taxa  
296 like *Filipendula* and *Equisetum* indicates the paludification, i.e., the peat initialisation, of the area probably also linked  
297 with the augmentation of *Poaceae* and *Cyperaceae*. In PAZ3b, the continuous representation of *Pinus* percentages of  
298 around 30% signals the forest development of the surrounding area. The intensification of tree cover has led to a  
299 decline of heliophilous shrubs like *Betula* and *Juniperus* and herbaceous taxa like *Artemisia* and *Helianthemum*.

300 *Chronostratigraphic constraints:* Four radiocarbon dates have been performed in this unit:  $12460 \pm 70$  (15024-14220  
301 cal. BP),  $12140 \pm 70$  (14184-13779 cal. BP),  $10970 \pm 60$  (12993-12718 cal. BP) and  $10890 \pm 60$  (12905-12688 cal. BP).  
302 The oldest dates at the base of the peaty Unit 3 coincide with the Oldest Dryas-Bølling transition (Fig. 2). Since the  
303 younger dates at the top of the unit are concomitant with the Allerød-Younger Dryas transition, we suggest that Unit  
304 3 was deposited during the Bølling-Allerød period.

305 *Interpretation:* The increase in shrub representation is consistent with the vegetation changes recorded in Western  
306 Europe as "The Bølling biozone" corresponding to the GI-1e of the event stratigraphy (Lowe et al., 2008). The  
307 consistent increase in *Pinus* percentages is probably linked to the development of mixed pine forest and is assigned to

308 the Allerød biozone in NW Europe. The deposits are defined by the presence of a homogeneous peat layer and a low  
 309 abundance of a clay fraction. These characteristics are indicative of a phase of decreasing river dynamics that evolves  
 310 towards a swamp environment. The peaty F6 probably developed under water-saturated and anaerobic conditions.  
 311 The evolution from silt to organic-rich sediment could record a decrease in sediment supply due to soil stabilization  
 312 linked to a densification of the vegetation cover induced by the temperature increase and wetter conditions during  
 313 the Bølling-Allerød period.

314 The difference of facies observed in the lowermost part of Unit 3 in MDV-1 and MDV-2 (absence of erosion surface,  
 315 differential preservation of silty beds, thick peat layer) and MDV-3, MDV-4 and MDV-5 (erosion surface, thinner peat  
 316 layer) could reflect a differential response during the climate transition depending of the channel activity. In the  
 317 abandoned channel, no erosion has occurred and the silt to organic-rich deposits evolution may reflect some delay,  
 318 *i.e* the time required for the vegetation to develop. In the active channel, the decrease in sediment supply and the  
 319 increase of the water table would have favoured a brief erosion phase in the channel. The only preserved deposits are  
 320 the peaty deposits. We suggest that the incision phase in the peaty deposits of the Unit 3 could be attributed to the  
 321 climate change from the Allerød to the Younger Dryas period.



322

323 Fig. 7. Pollen diagram from MDV1 core. AP: arboreal and shrub pollen; NAP: non-arboreal pollen (herbaceous pollen); PAZ: pollen assemblage  
 324 zone. Values are presented as percentages (black curves) and with a 10-times exaggeration (grey curves).

#### Unit 4 (SU 4 / PAZ 4)

*Sedimentology:* The base of SU 4 starts with a homogeneous greyish clay (F2) intercalated with few organic thin layers that fill the undulating topography formed during the previous incision phase. The unit thickness varies laterally from a few centimetres (MDV-1) to a few ten centimetres (MDV-4). The organic matter content remains very low (only a few %), as observed in Unit 2.

*Palynology:* The pollen changes indicate the return to open herbaceous communities with the extension of steppe taxa, especially *Artemisia* (Fig. 77). The presence of the low competitive taxa *Ephedra* indicates the emergence of coversand and full sun exposure areas. The openness of the forested landscape explains the increase in other heliophilous shrubs of *Juniperus* and *Hippophae* and even *Betula*. The occurrence of pollen of *Potamogeton sp.* can be explained only by the presence of standing water near the place of the core. We can detect a partition of this zone with a first part characterized by a greater pollen percentage of steppe taxa-Poaceae and aquatic plants and a second part with greater pollen percentages of Cyperaceae and *Betula* and a strong drop in Poaceae. These two biozones are separated by a sterile level.

*Chronostratigraphic constraints:* One radiocarbon date from the base of the unit returns an age of around  $10560 \pm 50$  (12661-12410 cal. BP) and confirms that U4, which follows stratigraphically the Bølling-Allerød period, is attributed to the onset of the Younger Dryas.

*Interpretation:* The preservation of silty deposits (F2) reveals an increase in sediment supply, a fluvial aggradation, and thus the reactivation of a low-energy river system. We suggest that a rarefaction of the stabilizing vegetation cover and a lowering of the water table have occurred to explain the record of detrital silty sediments that feed the channels. Therefore, it is supposed that Unit 4 records the functioning of low-energy channels in colder conditions during the Younger Dryas period.

#### Unit 5 (SU 5 / PAZ 5)

*Sedimentology:* This unit consists of a clay layer interbedded with frequent organic-matter rich beds (F8), displaying between 15 to 60% of organic matter content. Some peaty layers, containing up to 75 %, can be identified in MDV-1, MDV-2, MDV-4 and MDV-12 cores (Fig. 5). The top of the unit is truncated by an incision surface observed in all cores (Fig. 66).

351 *Palynology*: The sharp increase in *Pinus* pollen and spore monolet curves indicates a marked environmental change  
352 where the development of large forested areas caused the quasi disappearance of herbaceous and shrubs heliophilous  
353 taxa (Fig. 77). The strong decrease of Cyperaceae percentages is notable. The stable presence of taxa linked to wet  
354 soils like *Salix* and *Filipendula* shows however the persistence of swampy areas.

355 *Chronostratigraphic constraints*: Two radiocarbon dates have been performed in the Unit 5:  $10220 \pm 50$  (12131-11752  
356 cal. BP) and  $10110 \pm 60$  (11993-11400 cal. BP). These confirm that the U5 is dated to the end of the Younger Dryas.

357 *Interpretation*: The succession of detrital layers intercalated between the organic layers may reflect some fluctuations  
358 of the water table and the return to a marshy landscape. Clay deposits mark flood events in a low-energy river system  
359 whereas peaty layers reflect a water table increase. The sharp erosion observed at the top of the unit could record the  
360 transition between the Younger Dryas and the Holocene. The disappearance of cold-adapted species and the forest  
361 recovery correspond to the Younger Dryas-Holocene transitional phase.

## 362 **Unit 6 (SU 6)**

363 *Sedimentology*: It corresponds to a gyttja deposit that can be observed up to the ground surface. This non-cohesive  
364 facies (F9) is composed of clay and organic matter which can reach up to 75 % (Fig. 5). The heterogeneous and non-  
365 cohesive sediments appear fairly reworked. Magnetic susceptibility presents slightly negative values.

366 *Chronostratigraphic constraints*: No  $^{14}\text{C}$  dating have been performed. However, the gyttja deposits and the  $^{14}\text{C}$  dating  
367 in the Unit 5 are consistent with an Early Holocene record.

368 *Interpretation*: The gyttja deposit is indicative of a wet environment similar to the current fen, which signifies that the  
369 environmental conditions have not significantly evolved since the beginning of the Holocene. The hiatus highlighted  
370 between Units 5 and 6 could be either attributed to Holocene climate change or to more recent anthropogenic activity.  
371 Indeed, we cannot exclude that the industrial exploitation of the peat has contributed to the erosion and reworking  
372 of the gyttja (Salaün and Marre, 2005).

## 373 4.3 Morphodynamic evolution and climatic cycles

374 The combination of data coming from the image analyses, the geophysical acquisitions and the sedimentary and  
375 palynological records have highlighted changes in discharge and channel morphology in response to short phase  
376 climate oscillations. We observed (1) a multi-channel network that has developed in the alluvial plain, (2) the record

377 of successive cut-and-fill phases in the sedimentary record, (3) a sedimentary filling mainly composed of clay, silts and  
378 peats. All these data allow us to define periodic fluctuations of the water level, with six successive phases of river  
379 dynamics in the Boitet catchment during the Lateglacial and Early Holocene periods.

### 380 **Pleniglacial (Unit 1)**

381 During the Pleniglacial period, the landscape is dominated by high-energy braided rivers transporting coarse  
382 sediments. Pollinic data indicate both wet conditions (Cyperaceae) and steppe environment with sparse dwarf-birchs  
383 and grass vegetation. The transition from the Pleniglacial to the Oldest Dryas appears to be marked by an incision  
384 phase potentially attributed to a climate change, but it cannot be totally excluded that the channel morphology filled  
385 by the Oldest Dryas deposits was inherited from the Pleniglacial.

### 386 **Oldest Dryas (Unit 2)**

387 The hydro-sedimentary dynamic of the river evolves during the Oldest Dryas. The image analyses and geophysical  
388 prospection reveal a river system consisting of low-energy multiple channels. In this river system, the main active  
389 anastomosed channels display a silty-clayey sedimentation, which may have occurred partly during an early phase of  
390 the Bølling period as evidenced by Pastre et al. (2003), while abandoned channels are filled by organic deposits. Indeed,  
391 several gullies were visible on the orthophotographs and/or on the coring transect that indicated a complex network  
392 of palaeochannels. Besides, one larger meandering channel with a coarser sedimentation seems to flow along the  
393 easternmost part of the valley. Palynological data suggests a landscape dominated by steppe meadows (*Poaceae*,  
394 *Helianthemum*) in which some anastomosed channels bordered by shrubs (dwarf-birchs and willows) develop. The  
395 deposition of a new sedimentary unit (SU2) without any gradual transition with the former Pleniglacial coarse deposits  
396 (SU1) indicates a significant change of the sediment discharge and of the Boitet River morphodynamics, which  
397 occurred between the Last Glacial Maximum and the Oldest Dryas. The timing of this change is not clear; it could be  
398 marked and occurred in a short time period, but also, the lack of constraint on the chronology of this phase does not  
399 allow us to exclude the possibility of a long time gap between the incision of SU1 and the aggradation of SU2.

### 400 **Bølling-Allerød (Unit 3)**

401 After the cold episode of the Oldest Dryas, the detrital sedimentation disappears at the onset of the Bølling, suggesting  
402 a decline of the fluvial activity. The sedimentation, dominated by organic matter rich deposits, is indicative of a  
403 significant change in terms of fluvial processes, sedimentary supply and vegetation cover at the catchment scale. This

404 is reflected by warmer conditions favouring vegetation colonisation by shrubs and willow trees and soil stability. The  
405 presence of more aquatic vegetation suggests a mean water table rise coeval with the climatic amelioration of the  
406 Bølling period. In the second part of the Allerød, the amount of pines increases to replace an open forest vegetation  
407 cover.

### 408 **Younger Dryas (Units 4/5)**

409 A new incision phase is recorded at the end of the Allerød or at the onset of the Younger Dryas. The Younger Dryas  
410 period is marked by detrital sedimentation composed of silty-clayey load and the reactivation of a multi-channel  
411 system. The fluvial activity and the disappearance of the vegetation cover indicate both a lowering of the water table,  
412 that could be induced by climate changes, tectonics and any eustatic processes (Vandenbergh, 2015). The vertical  
413 deformation rate of the Paris Basin (0.01 m/1kyr) are too low to explain the observed base-level fall (Briais et al. 2016),  
414 and we suspect that this incision phase results from climate change, evolving towards cooler and drier conditions that  
415 characterized the Younger Dryas period. The vegetation is dominated by a steppe environment and a decrease of the  
416 amount of the pines. In the second period of the Younger Dryas, the increase of the organic matter beds in the  
417 sediment record suggests a recolonization of the vegetation. The landscape evolves towards an open birch-pine  
418 woodland in which sinuous rivers flow.

### 419 **Holocene (Unit 6)**

420 The general incision observed in all cores suggests a climate change occurring at the transition between the Younger  
421 Dryas and the Holocene. Abrupt changes both in vegetation and in the fluvial dynamics take place. Detrital  
422 sedimentation disappears to be replaced by gyttja deposits. A new phase of water table rise is recorded. The landscape  
423 is dominated by a marsh and the densification of a pine forest.

## 424 **5 Discussion**

425 Hereafter, we will compare successively the six phases of river dynamics observed in the Boitet catchment to those  
426 described in NW Europe during the Lateglacial and Early Holocene periods. Ensuingly, the preservation potential and  
427 geomorphological features as key factors for recording Lateglacial alluvial deposits will be discussed.

### 428 **5.1 Climate changes and river adjustments: an interregional analysis**

429 In NW European rivers, many studies have shown that the Oldest Dryas cooling period has been poorly preserved in  
430 fluvial deposits. In most channels located in lowland areas of northern France, Lateglacial sedimentary records have

431 been incised at the beginning of the Bølling (Antoine, 1997; Pastre et al., 2000). Just prior to this major incision phase,  
432 a few relics of fluvial deposits reworking the former Pleniglacial loess have been noticed in the Oise and Marne valleys  
433 of the Paris Basin (Antoine et al., 2003; Pastre et al., 2003), in which Upper Palaeolithic Magdalenian sites of Etiolles,  
434 Verbrerie and Pincevent are interstratified and dated from 13000 to 12000 BP (Valladas, 1994). These deposits are  
435 similar to the silty fluvial sequence (SU2) described in this paper, although the latter is well developed. Therefore, the  
436 Boitet River provides a rare case of Lateglacial sedimentary archives in northern France marked by a marked incision  
437 phase followed by an Oldest Dryas – early Bølling alluvial deposit (Fig. 88).

438 In the Boitet catchment, the onset of Bølling warming period is closely correlated to the storage of organic infilling.  
439 Ages obtained at the very bottom layer of the channels around 14612 cal. BP (15024-14220 cal. BP) and 14009 cal. BP  
440 (14184-13779 cal. BP) are consistent with the oldest Lateglacial peaty deposits of the Selle and the Thérain rivers of  
441 the Paris Basin, respectively dated from 14472 cal. BP (14827-14150 cal. BP) and 14748-14081 cal. BP (Antoine et al.,  
442 2003; Pastre et al., 2003), making the Boitet sequence the oldest alluvial record from Lateglacial in northern France.  
443 Our dates also indicate that the shift from cold to warmer climate at the very early beginning of the Bølling may have  
444 provoked the re-incision of some of the inherited channels to explain the presence of gyttja deposits around 14.6 cal.  
445 BP. Similar incision surfaces have been broadly illustrated elsewhere in NW and Central Europe (Vandenberghe, 1995;  
446 Huisink, 1999) and interpreted as a response to climate warming at the Oldest Dryas-Bølling transition. However, in  
447 our case, the wide standard deviation does not allow us to date precisely the onset of the incision. Moreover, the  
448 preservation of organo-clastic deposits prior to this peat accumulation in some channels of the Boitet River may  
449 indicate a minor incision phase which predates the main temperature increase ca. 14.7 cal. BP. Such an assumption  
450 has already been suggested in the Somme valley of the Paris Basin (Antoine et al., 2003), in the Jeetzel River in northern  
451 Germany (Turner et al., 2013) or in the Dordogne River in SW France (Bertran et al., 2013), where river incision  
452 occurred before the main climate warming in response to seasonality changes, e.g. a subtle increase in vegetation  
453 cover (Wagner-Cremer and Lotter, 2011). Vegetation cover is indeed perceived as a direct and key control mechanism  
454 on river morphodynamics (Vandenberghe, 2003).

455 During the Bølling-Allerød, no evidence of changes in the river morphodynamics have been detected. However, pollen  
456 analysis highlights a clear division of the peat accumulation which is defined firstly by a prevailing of pioneer shrub  
457 vegetation marked by an increase of *Juniperus* and *Betula*, secondly, after a transitional layer, by a sharp rise of *Pinus*  
458 leading to the high amount of AP grains (80%). This abrupt change in vegetation cover, usually characterised by a  
459 crossing-over of the *Betula* and *Pinus* curves, is commonly recognised in NW Europe and occurred within the Allerød

460 biozone (Hoek, 1997; De Klerk, 2008). Indeed, in the Paris Basin, the vegetation was largely dominated by *Betula* before  
461 being replaced by *Pinus* (Leroyer, 1997; Pastre et al., 2000), and the same trend has been observed in lowland areas  
462 of northern regions, e.g. in Belgium (Mullenders et al., 1972; Munaut and Paulissen, 1973; Crombé et al., 2013) or in  
463 the Netherlands (Bohncke, 1993; Hoek, 1997; Bos, 2001). Overall, the biozones defined from our palynological data  
464 are in accordance with regional synthetic studies from east (Turner et al., 2013) to west (Pastre et al. 2003): the Oldest  
465 Dryas open and steppe environment is respectively consistent with the OV-I (“Open Vegetation I”) and PAZ 1/2 as well  
466 as the Bølling shrub formation with the Hippophae-phase and PAZ 3. From 14 to 13.5 cal. BP, the transitional layer  
467 which is characterized by low values of *Betula* and *Pinus* and a rise of *Poaceae* may correspond to the first part of the  
468 Allerød, described as the OV-II and Allerød a-b (Turner et al., 2013) and PAZ 4/5 (Pastre et al., 2003). Finally, the *Pinus*  
469 rise occurred simultaneously around 13.5 – 13.3 cal. BP in these different areas, which confirmed the relevance of our  
470 biostratigraphical divisions and the correlation with the ice-core chronology already noticed in these previous studies.

471 The onset of the Younger Dryas is marked by the incision of the Allerød unit and then by the accumulation of a mixture  
472 of chalky and silty sediments in channels (Fig. 88). These deposits are supposedly produced by the weathering of chalky  
473 bedrock on the slopes during freeze-thaw cycles and then transported from the hillslopes by intensive runoff (Antoine  
474 et al., 2003). These interpretations imply both a climatic deterioration and a decrease of vegetation cover that lead to  
475 an increase of the fluvial activity. These hypotheses are consistent with other studies performed in the Paris Basin and  
476 in NW European rivers (Vandenberghe et al., 1994; Pastre et al., 2003). The rise in NAP percentages of steppe taxa and  
477 heliophilous shrubs and the decrease of *Pinus* pollen are representative of the beginning of the Younger Dryas biozone  
478 of NW Europe around 10950 BP (Isarin, 1997). Therefore, the sedimentological and palynological features of the Boitet  
479 sequence are archetypal of the onset of Younger Dryas (Limondin-Lozouet and Antoine 2001; Pastre et al., 2003). Our  
480 data reveal a twofold division for this period in the pollen record that we interpret as reflecting the shift from a cold  
481 and wet phase toward a warmer and drier climate (Fig. 88). A similar subdivision of malacological and palynological  
482 sequences has been identified in the Oise and the Somme valleys (Paris Basin), (Limondin-Lozouet and Antoine, 2001;  
483 Ponel et al., 2005), but also has been broadly recognised throughout Western Europe (Bohncke et al., 1993; Walker,  
484 1995) as far as the Mediterranean area (Burjachs et al., 2016), and eastward up to the Jeetzel valley in Germany (Turner  
485 et al., 2013). This subdivision is attributed to changes in the North Atlantic thermohaline circulation (Isarin et al., 1998).






Fig. 8. Comparison of the main climatic, fluvial and vegetation changes in the Boitet valley during Lateglacial.

## 5.2 Preservation potential and geomorphological features as key factors for recording Lateglacial alluvial deposits

Sedimentary sequences in the Marais de Saint Gond catchment reveal global similarities with NW European rivers that are interpreted as fluvial system evolution to climate changes. However, some local differences have also been highlighted. Among them, two distinctive features of the Boitet sequence are the preservation of Oldest Dryas deposits, and the development of low-energy multichannel systems during the Oldest Dryas and Younger Dryas. It is likely that these features can be partly explained by some local geomorphological context (Mol et al., 2000), illustrated by the investigation of a headwater drainage area of a tributary river, with a small-sized catchment.

Our study reveals a low-energy anastomosed system of numerous narrow channels of 10-20 m width prior to Bølling organic rich sedimentation. Some narrow multichannel systems occurring after the Pleniglacial and displaying a low sinuosity have been described in NW European rivers, such as those highlighted in the Selle valley (Antoine et al., 2003; Ponel et al., 2005) or in the Maas and Warta rivers in Poland (Vandenberghé et al., 1994). These multichannel systems

500 have been interpreted as 'transitional systems' to explain the fluvial changes from Pleniglacial braided to Bølling  
501 meandering rivers (Vandenberghe, 1995; Huisink, 1997; Kasse et al., 2005; Kasse et al., 2010; Erkens et al., 2011;  
502 Turner et al., 2013). Several have demonstrated that this transitional phase could take place before 14.7 cal. BP  
503 (Huisink, 1997).

504 Transitional patterns in alluvial system was initially perceived as an intrinsic change of the river morphodynamics in  
505 response to climate change. For instance, such patterns have been commonly described in cold-to-warm transition,  
506 such as Pleniglacial to Lateglacial (Huisink, 1997). In the case of anastomosed rivers, they are characterised by  
507 aggradational processes and developed under different conditions; favoured by a low-gradient plain where the  
508 amount of available sediment is higher than the capacity of transport or by an upstream area with high sediment  
509 supply. Aggradation arises in downstream accumulation zones where channels are not able to transport all sediment  
510 coming from upstream, as it is the case in alluvial fans (Smith and Smith, 1980; Smith, 1986; Mather et al., 2017) or  
511 deltas (Makaske, 2001; Makaske et al., 2017). Alternatively, such patterns are found in tributaries or headwater  
512 drainage areas due to a high-discharge/excess of sediment supply, e.g. the Morava River (Bábek et al., 2018), the  
513 Jeetzel valley (Turner et al., 2013) or the Moervaart palaeolake (Crombé et al., 2013). The latter reveals an evolution  
514 of the drainage system, from an anastomosed network to a single meandering channel, caused by lacustrine level  
515 decrease. In the Dordogne valley, Bertran et al. (2013) demonstrate the development of an anabranching river around  
516 18000 – 17000 cal. BP which lasted throughout the Lateglacial. Therefore, the identification of an anastomosed river  
517 system in the Boitet catchment is consistent with these observations. Moreover, this anastomosed system is a  
518 tributary of a large and marshy accumulation area (Saint-Gond marshes) resulted from the two hydrographic captures.  
519 The capture of the Somme-Soude River at the end of Pleniglacial, which has taken place upstream of the Saint-Gond  
520 marshes and the Boitet catchment, probably induced a major disruption in the drainage system (Tricart, 1949). As  
521 demonstrated by Maher et al. (2007), such capture lead to a decrease in water and sediment discharges and to a valley  
522 infilling with fines in the beheaded river system. This suggests that it was partly responsible for an alluvial aggradation  
523 of the Saint-Gond marshes area and of its tributaries. In addition to this geomorphological context, the headwater  
524 position of the Boitet catchment characterised by a connectivity of the hillslope to the valley floor, and the chalky  
525 bedrock weathered by freeze-thaw cycles, may explain the development of this anastomosed pattern characterised  
526 by fine alluvial aggradation during the Oldest Dryas cold period. Such a fluvial evolution as a response to both climatic  
527 change and particular catchment features have been already demonstrated in Belgium (Crombé et al., 2013) and may  
528 correspond to allocyclic-autocyclic coupling factors which could have triggered the moment when a threshold was

529 exceeded (Schumm, 1977). Indeed, the latter developed the idea that fluvial records may not only reflect external  
530 forcing factors like climatic changes but also intrinsic factors. Intrinsic features, e.g. topography, geology, vegetation  
531 cover, may be responsible for delayed responses or autonomous changes, and these factors and processes could differ  
532 from one basin to another (Prosser et al., 1994).

533 Analyses of depositional sequences in river systems show that alluvial records are formed by successive cut-and-fill  
534 cycles. As a consequence, the preservation potential of sedimentary sequences and river morphodynamic evolution is  
535 ultimately dependent of the intensity of the incision/sedimentation phases. For instance, most studies performed in  
536 NW Europe are focused on large catchments (>2500 km<sup>2</sup>) in downstream areas of main rivers and have stressed a  
537 broad incision at the onset of Bølling warming, hence the Oldest Dryas aggradation may have been eroded. Conversely,  
538 we suspect that the preservation potential of sedimentary sequences in those small-scale catchments is higher than  
539 for larger rivers, favoured by an upstream position, as suggested by Houben (2003). Indeed, in a headstream area, the  
540 river dynamics are more dependent on internal factors such as topography or subsoil lithology than that of a wider  
541 lowland river which gathers many tributaries (Vandenberghe, 1995; Vandenberghe, 2003). The relative importance of  
542 local factors in the alluvial preservation may explain why some headwater catchments present some peculiarities, e.g.  
543 the Rochy-Condé in the Thérain valley (Paris Basin). Conversely, sedimentary deposits in downstream areas reflect the  
544 global evolution of the entire catchment as the sum of the tributary responses. These will erase their specific local  
545 features leading to a climate-driven record, such as it has been shown in the Somme and Seine rivers. In the Paris  
546 Basin, Pastre et al. (2003) had already noticed that the geomorphological context contributes to the river evolution in  
547 spite of the major influence of climatic shifts. They demonstrate the contrast between main rivers active channels and  
548 small catchments. During Bølling, channel infilling deposits of the main valleys (e.g. Oise valley at Houdancourt and  
549 Lacroix-Saint-Ouen sites or Seine valley at Bazoches site) are characterised by a sandy-silty load. On the contrary, like  
550 the Boitet valley, organic deposits and peat formation are much more common for this period in small valleys and  
551 abandoned channels e.g. the Selle and Thérain valleys described above. The Thérain River is also a good point of  
552 comparison with our study; a pre-Bølling organic-rich deposit, subsequent to an undated clayey-silty deposition, has  
553 been dated to around 13,250 ± 84 BP (16126-15475 cal. BP; Pastre et al., 2003). Despite the lack of sedimentological  
554 details, the presence of a few organic silty layers prior to Bølling warming, which is rarely observed, may be related to  
555 the first organic sedimentation of the channel II (Fig. 6). These features further highlight the importance of the local

geomorphological conditions, such as the catchment size or the channel activity, and the resulting preservation potential, on the river evolution.

Besides temporal dynamics, spatial dynamics also have significant implications for river evolution. In addition to the narrow multichannel system, one larger paleomeander of 40 m width characterized by coarser sedimentation is observed in the easternmost part of the valley (Fig. 3) which hypothetically may have remained partly active during Bølling-Allerød stadial. Such variability of the fluvial dynamics in the alluvial plain is frequently observed: numerous NW European studies have noticed a complex Lateglacial multichannel transition from braided to meandering patterns during which these dynamics differed depending on channel activity, e.g. the abandonment of some of the channels whereas some of them remained active at the same time (Vandenberghe et al., 1994; Crombé et al., 2013; Turner et al. 2013).

## 6 Conclusion

We demonstrate that the fluvial evolution is triggered by climate changes and by geomorphological controls that can be strongly coupled, particularly in headwater areas where preservation potential reveals contrasting responses with different timings. Therefore, the choice of the geomorphological area (headwater area; wetland or lakes; alluvial plain of a wide drainage basin) seems decisive and determines the preservation potential of alluvial river systems. Upstream sections are important areas to investigate hydro-sedimentary delivery from slopes to valley floors. As such, they record more directly hydrosystem responses to forcing factors. It appears that focus on the headwater streams provides new data about hydrosystem adjustment to the last Glacial-Interglacial transition. The coupled 'spatial-multiproxy' analyses and comparisons with more regional environmental data and climate reconstructions demonstrate the importance of climatic mechanisms on the whole hydrosystem and particularly the strong potential of preservation of alluvial archives in headwater drainage basins.

## 7 Acknowledgments

This work has benefited from the support of the research program "Programme sur le Néolithique des Marais de Saint-Gond et de la vallée du Petit Morin (Marne, France)" (R. Martineau dir.) and the PRES Bourgogne Franche-Comté for funding of the fieldwork, the laboratory analyses and the radiocarbon dating. The laboratory measurements were undertaken at the PES and CPEP platforms of the UMR 6249 Chrono-environnement - UBFC University. We thank Julien Didier, Maxime Mermet, Léa Bidaud and Vincent Favreau for their help with the coring and the geophysical

583 investigations and Thomas Keep for assistance in translation. We are also grateful to Mr. and Mrs. Mathieu for their  
584 logistical support on the fieldwork. Suggestions from Jean-François Pastre, Jef Vandenberghe, Laurent Deschodt and  
585 Martin Stokes improved this manuscript.

## 586 8 References

- 587 Alley, R., Agustsdottir, A., 2005. The 8k event: cause and consequences of a major Holocene abrupt climate change. *Quaternary Science Reviews*  
588 24, 1123–1149. <https://doi.org/10.1016/j.quascirev.2004.12.004>
- 589 Antoine, P., 1997. Modifications des systèmes fluviaux à la transition Pléni-glaciaire-Tardiglaciaire et à l'Holocène : l'exemple du bassin de la  
590 Somme (Nord de la France). *Géographie physique et Quaternaire* 51, 93. <https://doi.org/10.7202/004763ar>
- 591 Antoine, P., Munaut, A.-V., Limondin-Lozouet, N., Ponel, P., Dupéron, J., Dupéron, M., 2003. Response of the Selle River to climatic modifications  
592 during the Lateglacial and Early Holocene (Somme Basin-Northern France). *Quaternary Science Reviews* 22, 2061–2076.  
593 [https://doi.org/10.1016/S0277-3791\(03\)00180-X](https://doi.org/10.1016/S0277-3791(03)00180-X)
- 594 Bábek, O., Sedláček, J., Novák, A., Létal, A., 2018. Electrical resistivity imaging of anastomosing river subsurface stratigraphy and possible controls  
595 of fluvial style change in a graben-like basin, Czech Republic. *Geomorphology* 317, 139–156.  
596 <https://doi.org/10.1016/j.geomorph.2018.05.012>
- 597 Barber, D.C., Dyke, A., Hillaire-Marcel, C., Jennings, A.E., Andrews, J.T., Kerwin, M.W., Bilodeau, G., McNeely, R., Southon, J., Morehead, M.D.,  
598 Gagnon, J.-M., 1999. Forcing of the cold event of 8,200 years ago by catastrophic drainage of Laurentide lakes. *Nature* 400, 344–348.  
599 <https://doi.org/10.1038/22504>
- 600 Beug, H.-J., 1961. Leitfaden der Pollenbestimmung für Mitteleuropa und angrenzende Gebiete. G. Fischer, Stuttgart.
- 601 Bertran, P., Frouin, M., Mercier, N., Naessens, F., Prodeo, F., Queffelec, A., Sirieix, C., Sitzia, L., 2013. Architecture of the lower terraces and  
602 evolution of the Dordogne River at Bergerac (south-west France) during the last glacial-interglacial cycle. *Journal of Quaternary Science*  
603 19, 605–616. <https://doi.org/10.1002/jqs.2656>
- 604 Björck, S., Kromer, B., Johnsen, S., Bennike, O., Hammarlund, D., Lemdahl, G., Possnert, G., Rasmussen, T.L., Wohlfarth, B., Hammer, C.U., Spurk,  
605 M., 1996. Synchronized Terrestrial-Atmospheric Deglacial Records Around the North Atlantic. *Science* 274, 1155–1160.
- 606 Björck, S., Walker, M.J.C., Cwynar, L.C., Johnsen, S., Knudsen, K.-L., Lowe, J.J., Wohlfarth, B., INTIMATE Members, 1998. An event stratigraphy  
607 for the Last Termination in the North Atlantic region based on the Greenland ice-core record: a proposal by the INTIMATE group.  
608 *Journal of Quaternary Science* 13, 283–292. [https://doi.org/10.1002/\(SICI\)1099-1417\(199807/08\)13:4<283::AID-JQS386>3.0.CO;2-A](https://doi.org/10.1002/(SICI)1099-1417(199807/08)13:4<283::AID-JQS386>3.0.CO;2-A)
- 609 Bos, J.A.A., 2001. Lateglacial and Early Holocene vegetation history of the northern Wetterau and the Amöneburger Basin (Hessen), central-west  
610 Germany. *Review of Palaeobotany and Palynology* 115, 177–212.
- 611 Bohncke, S.P.J., 1993. Lateglacial environmental changes in The Netherlands: spatial and temporal patterns. *Quaternary Science Reviews* 12,  
612 707–717.
- 613 Bohncke, S.P.J., Vandenberghe, J., Huijzer, A.S., 1993. Periglacial environments during the Weichselian Lateglacial in the Maas valley, the  
614 Netherlands. *Geologie en Mijnbouw* 72, 193–210.
- 615 Briais, J., Guillocheau, F., Lasseur, E., Robin, C., Châteauneuf, J.J., Serrano, O., 2016. Response of a low-subsiding intracratonic basin to long  
616 wavelength deformations: the Palaeocene–early Eocene period in the Paris Basin. *Solid Earth* 7, 205–228. <https://doi.org/10.5194/se-7-205-2016>
- 617
- 618 Bridgland, D.R., 2010. The record from British Quaternary river systems within the context of global fluvial archives. *Journal of Quaternary Science*  
619 25, 433–446. <https://doi.org/10.1002/jqs.1383>
- 620 Brown, A.G., 2001. *Alluvial geoarchaeology: floodplain archaeology and environmental change*, second. ed. Cambridge University Press, New  
621 York.
- 622 Burjachs, F., Jones, S.E., Giralt, S., Fernández-López de Pablo, J., 2016. Lateglacial to Early Holocene recursive aridity events in the SE  
623 Mediterranean Iberian Peninsula: The Salines playa lake case study. *Quaternary International* 403, 187–200.  
624 <https://doi.org/10.1016/j.quaint.2015.10.117>
- 625 Chambers, J.E., Wilkinson, P.B., Wardrop, D., Hameed, A., Hill, I., Jeffrey, C., Loke, M.H., Meldrum, P.I., Kuras, O., Cave, M., Gunn, D.A., 2012.  
626 Bedrock detection beneath river terrace deposits using three-dimensional electrical resistivity tomography. *Geomorphology* 177–178,  
627 17–25. <https://doi.org/10.1016/j.geomorph.2012.03.034>
- 628 Crombé, P., De Smedt, P., Davies, N.S., Gelorini, V., Zwertvaegher, A., Langohr, R., Van Damme, D., Demiddele, H., Van Strydonck, M., Antrop,  
629 M., Bourgeois, J., De Maeyer, P., De Reu, J., Finke, P.A., Van Meirvenne, M., Verniers, J., 2013. Hunter-gatherer responses to the  
630 changing environment of the Moervaart palaeolake (Nw Belgium) during the Late Glacial and Early Holocene. *Quaternary International*  
631 308–309, 162–177. <https://doi.org/10.1016/j.quaint.2013.05.035>
- 632 de Klerk, P., 2008. Patterns in vegetation and sedimentation during the Weichselian Late-glacial in north-eastern Germany. *Journal of*  
633 *Biogeography* 35, 1308–1322. <https://doi.org/10.1111/j.1365-2699.2007.01866.x>
- 634 Dearing, J., 1999. *Environmental Magnetic Susceptibility: Using the Bartington MS2 System*, second. ed. Chi Publishing, Kenilworth.
- 635 Deschodt, L., Salvador, P.G., Boulou, M., 2004. Formations sédimentaires et évolution de la vallée de la Deûle depuis le Pléni-glaciaire supérieur  
636 à Houplin-Ancoisne (Nord de la France). *Quaternaire* 15, 269–284. <https://doi.org/10.3406/quate.2004.1774>
- 637 ENVI/IDL (2018). <https://www.harrisgeospatial.com/Software-Technology/ENVI> (accessed 20/06/2019).
- 638 Erkens, G., Hoffmann, T., Gerlach, R., Klostermann, J., 2011. Complex fluvial response to Lateglacial and Holocene allogenic forcing in the Lower  
639 Rhine Valley (Germany). *Quaternary Science Reviews* 30, 611–627. <https://doi.org/10.1016/j.quascirev.2010.11.019>
- 640 Faegri, K., Iversen, J., 1989. *Textbook of Pollen Analysis (IV Ed.)*. J. Wiley and sons Ltd, 328 p.
- 641 Fleitmann, D., Mudelsee, M., Burns, S.J., Bradley, R.S., Kramers, J., Matter, A., 2008. Evidence for a widespread climatic anomaly at around 9.2  
642 ka before present: CLIMATIC ANOMALY AT AROUND 9.2 ka B.P. *Paleoceanography* 23, n/a-n/a.  
643 <https://doi.org/10.1029/2007PA001519>
- 644 Hausmann, J., Steinel, H., Kreck, M., Werban, U., Vienken, T., Dietrich, P., 2013. Two-dimensional geomorphological characterization of a filled

- 645 abandoned meander using geophysical methods and soil sampling. *Geomorphology* 201, 335–343.  
646 <https://doi.org/10.1016/j.geomorph.2013.07.009>
- 647 Hatrival, J.N., Chertier, B., Morfaux, P., 1988. Notice explicative de la feuille Montmort à 1/50 000. Bureau de Recherches Géologiques et Minières  
648 (BRGM), Orléans, 37 p. <http://infoterre.brgm.fr/> (accessed 17/06/2019).
- 649 Heiri, O., Lotter, A.F., Lemcke, G., 2001. Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility  
650 and comparability of results. *Journal of Paleolimnology* 25, 101–110. <https://doi.org/10.1023/A:1008119611481>
- 651 Hoek, W.Z., 1997. Atlas to Paleogeography of Lateglacial Vegetations. Maps of Lateglacial and Early Holocene landscape and vegetation in the  
652 Netherlands, with an extensive review of available palynological data. *Nederlandse Geografische studies* 231, Utrecht/Amsterdam,  
653 165 p.
- 654 Houben, P., 2003. Spatio-temporally variable response of fluvial systems to Late Pleistocene climate change: a case study from central Germany.  
655 *Quaternary Science Reviews* 22, 2125–2140. [https://doi.org/10.1016/S0277-3791\(03\)00181-1](https://doi.org/10.1016/S0277-3791(03)00181-1)
- 656 Huisink, M., 1999. Lateglacial river sediment budgets in the Mass valley, The Netherlands. *Earth Surface Processes and Landforms* 24, 93–109.  
657 [https://doi.org/10.1002/\(SICI\)1096-9837\(199902\)24:2<93::AID-ESP940>3.0.CO;2-R](https://doi.org/10.1002/(SICI)1096-9837(199902)24:2<93::AID-ESP940>3.0.CO;2-R)
- 658 Huisink, M., 1997. Late-glacial sedimentological and morphological changes in a lowland river in response to climatic change: the Maas, southern  
659 Netherlands. *Journal of Quaternary Science* 12, 209–223. [https://doi.org/10.1002/\(SICI\)1099-1417\(199705/06\)12:3<209::AID-  
661 JQS306>3.0.CO;2-P](https://doi.org/10.1002/(SICI)1099-1417(199705/06)12:3<209::AID-<br/>
660 JQS306>3.0.CO;2-P)
- 661 Isarin, R.F.B., 1997. Permafrost Distribution and Temperatures in Europe During the Younger Dryas. *Permafrost and Periglacial Processes* 8, 313–  
662 333. [https://doi.org/10.1002/\(SICI\)1099-1530\(199709\)8:3<313::AID-PPP255>3.0.CO;2-E](https://doi.org/10.1002/(SICI)1099-1530(199709)8:3<313::AID-PPP255>3.0.CO;2-E)
- 663 Isarin, R.F.B., Renssen, H., Vandenberghe, J., 1998. The impact of the North Atlantic Ocean on the Younger Dryas climate in northwestern and  
664 central Europe. *Journal of Quaternary Science* 13, 447–453. [https://doi.org/10.1002/\(SICI\)1099-1417\(1998090\)13:5<447::AID-  
666 JQS402>3.0.CO;2-B](https://doi.org/10.1002/(SICI)1099-1417(1998090)13:5<447::AID-<br/>
665 JQS402>3.0.CO;2-B)
- 666 Iversen, J., 1954. The Late-Glacial flora of Denmark and its relation to climate and soil. *Danmarks Geologiske Undersøgelse II.række* 80, 87–119.
- 667 Jessen, K., 1935. Archeological dating in the history of North Jutland's vegetation. *Acta Archeologica* 5, 185–214.
- 668 Kasse, C., Bohncke, S.J.P., Vandenberghe, J., Gábris, G., 2010. Fluvial style changes during the last glacial-interglacial transition in the middle  
669 Tisza valley (Hungary). *Proceedings of the Geologists' Association* 121, 180–194. <https://doi.org/10.1016/j.pgeola.2010.02.005>
- 670 Kasse, C., Hoek, W.Z., Bohncke, S.J.P., Konert, M., Weijers, J.W.H., Cassee, M.L., Van Der Zee, R.M., 2005. Late Glacial fluvial response of the  
671 Niers-Rhine (western Germany) to climate and vegetation change. *Journal of Quaternary Science* 20, 377–394.  
672 <https://doi.org/10.1002/jqs.923>
- 673 Laigre, L., Reynard, E., Arnaud-Fassetta, G., Baron, L., Glenz, D., 2012. Characterisation of the Rhône River palaeodynamics in Central Valais  
674 (Switzerland) with the electrical resistivity tomography method. *Géomorphologie : relief, processus, environnement* 18, 405–426.  
675 <https://doi.org/10.4000/geomorphologie.10020>
- 676 Leroyer, C., 1997. Hommes, Climat, Végétation au Tardi-et-Postglaciaire dans le Bassin parisien : apports de l'étude palynologique des fonds de  
677 vallée. Paris 1.
- 678 Limondin-Lozouet, N., Antoine, P., 2001. Palaeoenvironmental changes inferred from malacofaunas in the Lateglacial and early Holocene fluvial  
679 sequence at Conty, northern France. *Boreas* 30, 148–164. <https://doi.org/10.1111/j.1502-3885.2001.tb01219.x>
- 680 Litt, T., Brauer, A., Goslar, T., Merkt, J., Bałaga, K., Müller, H., Ralska-Jasiewiczowa, M., Stebich, M., Negendank, J.F.W., 2001. Correlation and  
681 synchronisation of Lateglacial continental sequences in northern central Europe based on annually laminated lacustrine sediments.  
682 *Quaternary Science Reviews* 20, 1233–1249. [https://doi.org/10.1016/S0277-3791\(00\)00149-9](https://doi.org/10.1016/S0277-3791(00)00149-9)
- 683 Lowe, J.J., Rasmussen, S.O., Björck, S., Hoek, W.Z., Steffensen, J.P., Walker, M.J.C., Yu, Z.C., 2008. Synchronisation of palaeoenvironmental events  
684 in the North Atlantic region during the Last Termination: a revised protocol recommended by the INTIMATE group. *Quaternary Science  
685 Reviews* 27, 6–17. <https://doi.org/10.1016/j.quascirev.2007.09.016>
- 686 Macklin, M.G., Jones, A.F., Lewin, J., 2010. River response to rapid Holocene environmental change: evidence and explanation in British  
687 catchments. *Quaternary Science Reviews* 29, 1555–1576. <https://doi.org/10.1016/j.quascirev.2009.06.010>
- 688 Maher, E., Harvey, A.M., France, D., 2007. The impact of a major Quaternary river capture on the alluvial sediments of a beheaded river system,  
689 the Rio Alias SE Spain. *Geomorphology* 84, 344–356. <https://doi.org/10.1016/j.geomorph.2005.07.034>
- 690 Makaske, B., 2001. Anastomosing rivers: a review of their classification, origin and sedimentary products. *Earth-Science Reviews* 53, 149–196.  
691 [https://doi.org/10.1016/S0012-8252\(00\)00038-6](https://doi.org/10.1016/S0012-8252(00)00038-6)
- 692 Makaske, B., Lavooi, E., de Haas, T., Kleinhans, M.G., Smith, D.G., 2017. Upstream control of river anastomosis by sediment overloading, upper  
693 Columbia River, British Columbia, Canada. *Sedimentology* 64, 1488–1510. <https://doi.org/10.1111/sed.12361>
- 694 Mather, A.E., Stokes, M., Whitfield, E., 2017. River terraces and alluvial fans: the case for an integrated Quaternary fluvial archive. *Quaternary  
695 Science Reviews* 166, 74–90. <https://doi.org/10.1016/j.quascirev.2016.09.022>
- 696 Matys Grygar, T., Elznicová, J., Tůmová, Š., Faměra, M., Balogh, M., Kiss, T., 2016. Floodplain architecture of an actively meandering river (the  
697 Ploučnice River, the Czech Republic) as revealed by the distribution of pollution and electrical resistivity tomography. *Geomorphology*  
698 254, 41–56. <https://doi.org/10.1016/j.geomorph.2015.11.012>
- 699 Mol, J., Vandenberghe, J., Kasse, C., 2000. River response to variations of periglacial climate in mid-latitude Europe. *Geomorphology* 33, 131–  
700 148. [https://doi.org/10.1016/S0169-555X\(99\)00126-9](https://doi.org/10.1016/S0169-555X(99)00126-9)
- 701 Moore, P.D., Webb, J.A., Collinson, M.E., 1991. *Pollen analysis*, second. ed. Blackwell, Oxford.
- 702 Mullenders, W., Desair-Coremans, M., Gilot, E., 1972. Recherches palynologiques et datations 14C sur les dépôts tourbeux de Holsbeek, in:  
703 Vermeesch, P. (Ed.), *Twee mesolitische sites te Holsbeek*. *Archeologia Belgica* 138, pp. 133-141.
- 704 Munaut, A.V., Paulissen, E., 1973. Evolution et paléocologie de la vallée de la petite Nèthe au cours du Post-Würm (Belgique). *Annales de la  
705 Société Géologique de Belgique* 96, 301-346.
- 706 North Greenland Ice Core Project members, Andersen, K.K., Azuma, N., Barnola, J.-M., Bigler, M., Biscaye, P., Caillon, N., Chappellaz, J., Clausen,  
707 H.B., Dahl-Jensen, D., Fischer, H., Flückiger, J., Fritzsche, D., Fujii, Y., Goto-Azuma, K., Grønvdold, K., Gundestrup, N.S., Hansson, M.,  
708 Huber, C., Hvidberg, C.S., Johnsen, S.J., Jonsell, U., Jouzel, J., Kipfstuhl, S., Landais, A., Leuenberger, M., Lorrain, R., Masson-Delmotte,  
709 V., Miller, H., Motoyama, H., Narita, H., Popp, T., Rasmussen, S.O., Raynaud, D., Rothlisberger, R., Ruth, U., Samyn, D., Schwander, J.,  
710 Shoji, H., Siggard-Andersen, M.-L., Steffensen, J.P., Stocker, T., Sveinbjörnsdóttir, A.E., Svensson, A., Takata, M., Tison, J.-L.,  
711 Thorsteinsson, T., Watanabe, O., Wilhelms, F., White, J.W.C., 2004. High-resolution record of Northern Hemisphere climate extending  
712 into the last interglacial period. *Nature* 431, 147–151. <https://doi.org/10.1038/nature02805>
- 713 Pastre, J.-F., Leroyer, C., 1997. La capture du Grand-Morin par la Marne (Bassin parisien, France) : âge et mécanisme. *Géographie physique et*

714 Quaternaire 51, 347. <https://doi.org/10.7202/033133ar>

715 Pastre, J.F., Leroyer, C., Limondin-Lozouet, N., Chaussé, C., Fontugne, M., Gebhardt, A., Hatté, C., Krier, V., 2000. Le Tardiglaciaire des fonds de  
716 vallée du Bassin Parisien (France) [The Late-Glacial from the Paris basin floodplains (France)]. *Quaternaire* 11, 107–122.  
717 <https://doi.org/10.3406/quate.2000.1660>

718 Pastre, J.-F., Limondin-Lozouet, N., Leroyer, C., Ponel, P., Fontugne, M., 2003. River system evolution and environmental changes during the  
719 Lateglacial in the Paris Basin (France). *Quaternary Science Reviews* 22, 2177–2188. [https://doi.org/10.1016/S0277-3791\(03\)00147-1](https://doi.org/10.1016/S0277-3791(03)00147-1)

720 Ponel, P., Coope, R., Antoine, P., Limondin-Lozouet, N., Leroyer, C., Munaut, A.-V., Pastre, J.-F., Guiter, F., 2005. Lateglacial palaeoenvironments  
721 and palaeoclimates from Conty and Houdancourt, northern France, reconstructed from Beetle remains. *Quaternary Science Reviews*  
722 24, 2449–2465. <https://doi.org/10.1016/j.quascirev.2004.12.010>

723 Prosser, I.P., Chappell, J., Gillespie, R., 1994. Holocene valley aggradation and gully erosion in headwater catchments, south-eastern highlands  
724 of Australia. *Earth Surface Processes and Landforms* 19, 465–480. <https://doi.org/10.1002/esp.3290190507>

725 Punt, W., 1976. *Northwest European Pollen Flora*, vol. I. Elsevier, Amsterdam

726 Punt, W., Blackmore, S., Clarke, G.C.S., 1988. *Northwest European Pollen Flora*, vol. V. Elsevier, Amsterdam.

727 Punt, W., Clarke, G.C.S., 1980. *Northwest European Pollen Flora*, vol. II. Elsevier, Amsterdam.

728 Punt, W., Clarke, G.C.S., 1981. *Northwest European Pollen Flora*, vol. III. Elsevier, Amsterdam.

729 Punt, W., Clarke, G.C.S., 1984. *Northwest European Pollen Flora*, vol. IV. Elsevier, Amsterdam.

730 Rasmussen, S.O., Bigler, M., Blockley, S.P., Blunier, T., Buchardt, S.L., Clausen, H.B., Cvijanovic, I., Dahl-Jensen, D., Johnsen, S.J., Fischer, H., Gkinis,  
731 V., Guillevic, M., Hoek, W.Z., Lowe, J.J., Pedro, J.B., Popp, T., Seierstad, I.K., Steffensen, J.P., Svensson, A.M., Vallelonga, P., Vinther,  
732 B.M., Walker, M.J.C., Wheatley, J.J., Winstrup, M., 2014. A stratigraphic framework for abrupt climatic changes during the Last Glacial  
733 period based on three synchronized Greenland ice-core records: refining and extending the INTIMATE event stratigraphy. *Quaternary  
734 Science Reviews* 106, 14–28. <https://doi.org/10.1016/j.quascirev.2014.09.007>

735 Reille, M., 1992. Pollen et spores d'Europe et d'Afrique du Nord. *Laboratoire de botanique historique et palynologie*, URA CNRS 1152.

736 Reille, M., 1995. Pollen et spores d'Europe et d'Afrique du Nord: supplément 1. Marseille: Laboratoire de Botanique Historique et Palynologie.

737 Reimer, P.J., Bard, E., Bayliss, A., Beck, J.W., Blackwell, P.G., Ramsey, C.B., Buck, C.E., Cheng, H., Edwards, R.L., Friedrich, M., Grootes, P.M.,  
738 Guilderson, T.P., Hafliadason, H., Hajdas, I., Hatté, C., Heaton, T.J., Hoffmann, D.L., Hogg, A.G., Hughen, K.A., Kaiser, K.F., Kromer, B.,  
739 Manning, S.W., Niu, M., Reimer, R.W., Richards, D.A., Scott, E.M., Southon, J.R., Staff, R.A., Turney, C.S.M., van der Plicht, J., 2013.  
740 IntCal13 and Marine13 Radiocarbon Age Calibration Curves 0–50,000 Years cal BP. *Radiocarbon* 55, 1869–1887.  
741 [https://doi.org/10.2458/azu\\_js\\_rc.55.16947](https://doi.org/10.2458/azu_js_rc.55.16947)

742 Rey, J., Martínez, J., Hidalgo, M.C., 2013. Investigating fluvial features with electrical resistivity imaging and ground-penetrating radar: The  
743 Guadalquivir River terrace (Jaen, Southern Spain). *Sedimentary Geology* 295, 27–37. <https://doi.org/10.1016/j.sedgeo.2013.07.003>

744 Salaün, F., Marre, A., 2005. L'anthropisation des marais de Saint Gond : conséquences sur leur physionomie et sur leur fonctionnement actuel  
745 (Marne - France). *Travaux de l'Institut Géographique de Reims* 31, 79–95. <https://doi.org/10.3406/tigr.2005.1499>

746 Schumm, S.A., 1977. *The fluvial system*. Wiley, New York.

747 Smith, D.G., 1986. Anastomosing river deposits, sedimentation rates and basin subsidence, Magdalena River, northwestern Colombia, South  
748 America. *Sedimentary Geology* 46, 177–196. [https://doi.org/10.1016/0037-0738\(86\)90058-8](https://doi.org/10.1016/0037-0738(86)90058-8)

749 Smith, D.G., Smith, N.D., 1980. Sedimentation in anastomosed river systems; examples from alluvial valleys near Banff, Alberta. *Journal of  
750 Sedimentary Research* 50, 157–164. <https://doi.org/10.1306/212F7991-2B24-11D7-8648000102C1865D>

751 Stuiver, M., Reimer, P.J., 1993. Extended 14C Data Base and Revised CALIB 3.0 14C Age Calibration Program. *Radiocarbon* 35, 215–230.  
752 <https://doi.org/10.1017/S0033822200013904>

753 Teller, J.T., Leverington, D.W., Mann, J.D., 2002. Freshwater outbursts to the oceans from glacial Lake Agassiz and their role in climate change  
754 during the last deglaciation. *Quaternary Science Reviews* 21, 879–887. [https://doi.org/10.1016/S0277-3791\(01\)00145-7](https://doi.org/10.1016/S0277-3791(01)00145-7)

755 Tricart, J., 1949. *La partie orientale du Bassin de Paris, étude morphologique*, Thèses Lettres. S.E.D.E.S., Paris.

756 Turner, F., Tolksdorf, J.F., Viehberg, F., Schwalb, A., Kaiser, K., Bittmann, F., von Bramann, U., Pott, R., Staesche, U., Breest, K., Veil, S., 2013.  
757 Lateglacial/early Holocene fluvial reactions of the Jeetzel river (Elbe valley, northern Germany) to abrupt climatic and environmental  
758 changes. *Quaternary Science Reviews* 60, 91–109. <https://doi.org/10.1016/j.quascirev.2012.10.037>

759 Valladas, H., 1994. Chronologie des sites du Magdalénien final du Bassin parisien, in: Taborin, Y. (Ed.), *Environnements et habitats magdaléniens  
760 dans le centre du Bassin parisien*. *Documents d'Archéologie Française* 43, pp. 65–68.

761 Van Huissteden, J., Vandenbergh, J., Van Geel, B., 1986. Late Pleistocene stratigraphy and fluvial history of the Dinkel basin (Twente, eastern  
762 Netherlands). *Eiszeitalter und Gegenwart* 36, 43–59.

763 Vandenbergh, J., 1981. Geomorphological and paleohydrographical research based on geoelectrical prospecting (South Campine, Belgium).  
764 *Bull. Soc. Belge Géol.* 90, 341–356.

765 Vandenbergh, J., 1995. Postglacial river activity and climate: state of the art and future prospects, in: Frenzel, B., Vandenbergh, J., Kasse, K.,  
766 Bohncke, S., Gläser, B. (Eds.), *European River Activity and Climatic Change during the Lateglacial and Early Holocene*.  
767 *Paläoklimaforschung/Palaeoclimate Research* 14, pp. 1–9.

768 Vandenbergh, J., 2003. Climate forcing of fluvial system development: an evolution of ideas. *Quaternary Science Reviews* 22, 2053–2060.  
769 [https://doi.org/10.1016/S0277-3791\(03\)00213-0](https://doi.org/10.1016/S0277-3791(03)00213-0)

770 Vandenbergh, J., 2015. River terraces as a response to climatic forcing: Formation processes, sedimentary characteristics and sites for human  
771 occupation. *Quaternary International* 370, 3–11. <https://doi.org/10.1016/j.quaint.2014.05.046>

772 Vandenbergh, J., Bohncke, S., Lammers, W., Zilverberg, L., 1987. Geomorphology and palaeoecology of the Mark valley (southern Netherlands):  
773 geomorphological valley development during the Weichselian and Holocene. *Boreas* 16, 55–67. <https://doi.org/10.1111/j.1502-3885.1987.tb00754.x>

774 Vandenbergh, J., de Smedt, P., 1979. Palaeomorphology in the eastern Scheldt basin (Central Belgium) - The dijle-demergrote nete confluence  
775 area -. *Catena* 6, 73–105. [https://doi.org/10.1016/S0341-8162\(79\)80005-3](https://doi.org/10.1016/S0341-8162(79)80005-3)

776 Vandenbergh, J., Kasse, C., Bohncke, S., Kozarski, S., 1994. Climate-related river activity at the Weichselian-Holocene transition: a comparative  
777 study of the Warta and Maas rivers. *Terra Nova* 6, 476–485. <https://doi.org/10.1111/j.1365-3121.1994.tb00891.x>

778 Vannièr, B., Bossuet, G., Walter-Simonnet, A.-V., Ruffaldi, P., Adatte, T., Rossy, M., Magny, M., 2004. High-resolution record of environmental  
779 changes and tephrochronological markers of the Last Glacial-Holocene transition at Lake Lautrey (Jura, France): LAST GLACIAL-  
780 HOLOCENE TRANSITION AT LAKE LAUTREY, FRANCE. *Journal of Quaternary Science* 19, 797–808. <https://doi.org/10.1002/jqs.873>

781

- 782 Wagner-Cremer, F., Lotter, A.F., 2011. Spring-season changes during the Late Pleniglacial and Bølling/Allerød interstadial. *Quaternary Science*  
783 *Reviews* 30, 1825–1828. <https://doi.org/10.1016/j.quascirev.2011.05.003>
- 784 Walker, M., Lowe, J., Blockley, S.P.E., Bryant, C., Coombes, P., Davies, S., Hardiman, M., Turney, C.S.M., Watson, J., 2012. Lateglacial and early  
785 Holocene palaeoenvironmental 'events' in Sluggan Bog, Northern Ireland: comparisons with the Greenland NGRIP GICC05 event  
786 stratigraphy. *Quaternary Science Reviews* 36, 124–138. <https://doi.org/10.1016/j.quascirev.2011.09.008>
- 787 Walker, M.J.C., 1995. Climatic changes in Europe during the last glacial/interglacial transition. *Quaternary International* 28, 63–76.  
788 [https://doi.org/10.1016/1040-6182\(95\)00030-M](https://doi.org/10.1016/1040-6182(95)00030-M)
- 789