



HAL
open science

Historical and fossil evidence of an extinct endemic species of *Leiocephalus* (Squamata: Leiocephalidae) from the Guadeloupe Islands

Corentin Bochaton, Laurent Charles, Arnaud Lenoble

► **To cite this version:**

Corentin Bochaton, Laurent Charles, Arnaud Lenoble. Historical and fossil evidence of an extinct endemic species of *Leiocephalus* (Squamata: Leiocephalidae) from the Guadeloupe Islands. *Zootaxa*, 2021, 4927 (3), pp.383-409. 10.11646/zootaxa.4927.3.4 . halshs-03358473

HAL Id: halshs-03358473

<https://shs.hal.science/halshs-03358473>

Submitted on 29 Sep 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Historical and fossil evidence of an extinct endemic species of *Leiocephalus***
2 **(Squamata: Leiocephalidae) from the Guadeloupe Islands**

3 CORENTIN BOCHATON^{1,2}, LAURENT CHARLES³ & ARNAUD LENOBLE¹

4 1- PACEA – UMR CNRS 5199, Université de Bordeaux, 33 615 Pessac, France. [corentin.bochaton@u-](mailto:corentin.bochaton@u-bordeaux.fr)
5 [bordeaux.fr](mailto:corentin.bochaton@u-bordeaux.fr); arnaud.lenoble@u-bordeaux.fr

6
7 2 - Max Planck Institute for the Science of Human History, Department of Archaeology, Kahlaische Straße 10,
8 07745 Jena, Germany.

9 3- Muséum de Bordeaux - sciences et nature, 5 Place Bardineau, Bordeaux 33 000, France. [l.charles@mairie-](mailto:l.charles@mairie-bordeaux.fr)
10 [bordeaux.fr](mailto:l.charles@mairie-bordeaux.fr)

11 **Abstract**

12 Documenting recent extinction events against the backdrop of increasing human-induced
13 environmental pressure is complicated by the lack of historical and subfossil evidence for most parts
14 of the world. This paucity of data renders it particularly difficult to evaluate the human impact on
15 fragile environments, such as small islands, that may have been heavily altered by historical human
16 exploitation. Here we describe a new species of an extinct *Leiocephalus* lizard from Guadeloupe,
17 *Leiocephalus roquetus* sp. nov. based on recent discoveries of both a previously undocumented
18 historically taxidermy specimen and of a large assemblage of subfossil bone remains from La Désirade
19 Island. This new species presents a primitive morphology compared to all extant species of its genus
20 and provides evidence for the past existence of a Lesser Antillean clade of *Leiocephalus* lizards that
21 was completely wiped out in the centuries following the European colonization of these islands. Our
22 study demonstrates how paleontological approaches can help to better understand the history of human
23 environmental impact as well as the degree of degradation of modern ecosystems.

24 **Keywords:** Curlytail lizard, Extinction, Human Impact, Lesser Antilles, Osteology.

25 **Introduction**

26 The West Indies are known to have been severely impacted over the last several hundred to thousand
27 years by both climatic and anthropogenic events that reshaped most of their biological diversity. The
28 rich historical, archaeological, and paleontological records of these islands reveal hundreds of
29 extinction events touching all taxa and regions (Cooke *et al.* 2017; Orihuela *et al.* 2020; Steadman *et*
30 *al.* 2015). The causes underlying these extinctions have been the subject of considerable debate, as
31 early research focused almost uniquely on the Pleistocene–Holocene transition as the main extinction
32 factor (e. g. Pregill & Olson 1981; Underwood 1964). However, numerous recent work has
33 demonstrated the considerable impact of human colonization and anthropization that led to the
34 extinction of a relatively substantial portion of the local biodiversity (Bochaton *et al.* 2015b; Cooke *et*
35 *al.* 2017; Soto-Centeno & Steadman 2015; Steadman *et al.* 1984, 2015). These studies also
36 demonstrated the West Indies to be a perfect case study for exploring the long term impact of humans
37 on island ecosystems in the framework of the current mass extinction crisis (Barnosky *et al.* 2011;
38 Ceballos *et al.* 2015, 2017). Of the taxa impacted by these extinction events, several were almost
39 completely wiped out from the Lesser Antilles. This group of strongly affected taxa comprises
40 multiple squamates, including diploglossine lizards (Bochaton *et al.* 2015a, 2016), *Boa* snakes
41 (Bochaton 2020; Bochaton & Bailon 2018) and leiocephalid lizards (Etheridge 1964; Pregill *et al.*
42 1988) the latter, which despite being one of the few squamate genera endemic to the West Indies, still
43 remains poorly understood. The monophyly of the *Leiocephalus* (Gray 1827) species restricted to the
44 West Indies was documented in the 1960s (Etheridge 1966) and confirmed by molecular evidence
45 (Pyron *et al.* 2013), with the members of this genus first included in their own subfamily and then
46 family (Leiocephalidae Frost & Etheridge 1989). The genus *Leiocephalus* currently comprises 24
47 species exclusively distributed in the Greater Antilles (Hedges 2020), including seven extinct species,
48 five of which are known only from fossil remains (Pregill 1992). Among these extinct species, two
49 were endemic to the Lesser Antilles, an area where no *Leiocephalus* species are currently known. The
50 first of these species is *Leiocephalus herminieri* (Duméril & Bibron 1837), whose fluid-preserved type
51 specimens were collected in Martinique during the 19th century (for a discussion concerning the origin
52 of these specimens see Breuil -2002-). The second species is *Leiocephalus cuneus* Etheridge, 1964, a
53 taxa that was originally described on the basis of fossil remains collected in Barbuda (Etheridge 1964;

54 Watters *et al.* 1984) but whose distribution was later extended to Antigua (Pregill *et al.* 1988), and
55 possibly to Anguilla, La Désirade, and Marie-Galante (Bailon *et al.* 2015; Boudadi-Maligne *et al.*
56 2016; Pregill 1992; Pregill *et al.* 1994; Stouvenot *et al.* 2014). The distribution of this taxon outside
57 Barbuda and Antigua is however doubtful, as no clear fossil evidence is currently available. In
58 Guadeloupe, the past occurrence of *Leiocephalus* lizards was mentioned in several historical texts
59 (Breuil 2002; Du Tertre 1654; de Rochefort 1658) although their precise taxonomic identification is
60 unclear. Historical data and biogeographic hypotheses all suggest that *Leiocephalus* may have
61 previously been widespread in the Lesser Antilles. The current fossil record may therefore be
62 substantially unrepresentative of the past diversity of this lizard. The phylogenetic relationships
63 between both modern and fossil *Leiocephalus* species are also poorly understood. Until recently, the
64 most comprehensive study of this genus concerned morphological characters combining both modern
65 and known fossil taxa (Pregill 1992). However, none of the clades recognized in this pioneering work
66 were confirming by the few non-exhaustive molecular analysis conducted since (Köhler *et al.* 2016;
67 Pyron *et al.* 2013). In addition, both morphological and molecular analyses seem to indicate a complex
68 colonization scenario comprising multiple colonization events. Overall, the history and past diversity
69 of the genus *Leiocephalus* in the Lesser Antilles is poorly documented. For example, we currently do
70 not know how these species were diversified in this area, where they came from, and when and why
71 they went extinct. Here we provide important new descriptions of both historical and fossil specimens
72 of *Leiocephalus* collected in Guadeloupe that we attribute to a new endemic species: *Leiocephalus*
73 *roquetus* sp. nov.

74 **Materials and method**

75 Dry museum specimen

76 The holotype of *Leiocephalus roquetus* sp. nov. is a dry taxidermy specimen stored in the Natural
77 History Museum of Bordeaux (MHNbX) in France. The specimen was donated to the museum around
78 1835 by Théodore Roger (1784–1838), an alcohol manufacturer and naturalist better known for his
79 collection of Lepidoptera. Very few details are available concerning the origin of the holotype

80 specimen; it is stored with a label reading “Holotropide de l’Herminier *Holotropis herminieri* Gray
81 Guadeloupe donation Roger” with the species author’s name “*herminieri*” erroneously attributed to
82 Gray, and “Guadeloupe” being the geographic origin of the specimen. This label is, in fact, a mid 20th
83 century copy of the original 19th century label, and it is possible that some information originally
84 attached to the specimen has been lost. The first catalog record to unambiguously mention this
85 specimen under the name *Holotropis herminieri* Duméril & Bibron, 1837 comes from the beginning of
86 the 20th century, suggesting that this specimen remained unidentified for years. The fact that the
87 locality “Guadeloupe” contradicts the (erroneous) species identification (*Leiocephalus herminieri*
88 being endemic to Martinique) is a clue that the origin was originally already attached to the specimen
89 as it was not inferred from what is potentially a more recent taxonomic attribution. Although the
90 general interest of this specimen was previously recognized (Breuil 2009), it remained unstudied until
91 now.

92 Subfossil material

93 In addition to the specimen from the Natural History Museum of Bordeaux, more than 300
94 *Leiocephalus* remains have been recorded from 31 different archaeological and paleontological
95 deposits in Guadeloupe islands (Bochaton 2016). However, only a single deposit, Pointe Gros
96 Rempart 6 (PGR6) on the island of La Désirade, yielded enough well-preserved bones to enable a
97 formal evaluation, description, and identification of the *Leiocephalus* species they represent. While
98 this site is still largely unpublished it nevertheless produced a noteworthy assemblage of squamate
99 remains (Bochaton 2020; Bochaton *et al.* 2018), including 218 of the 320 *Leiocephalus* remains
100 currently known from the Guadeloupe Islands. For this study, we compare the material from PGR6 to
101 the dry specimen from Bordeaux, retaining only the most well-preserved fossil remains for
102 description. We also integrated remains collected from other Guadeloupe islands, although no large,
103 well-preserved fossil assemblages outside of La Désirade are currently available to identify them to
104 species. The fossil bones from PGR6 are archaeological specimens and are therefore the property of
105 the French state. Currently stored at the *Service Regional de l’Archéologie de Guadeloupe*, we plan to

106 petition for these bones described as paratypes to be united with the holotype dry-specimen in a single
107 collection housed at the Natural History Museum of Bordeaux.

108 The site of Pointe Gros Rempart 6 (16819041.5600N, 6180049.1800W WGS84) is a pit cave situated
109 on the coastal plain of La Désirade Island. This island was once connected to the other main islands of
110 the Guadeloupe archipelago (Grande-Terre, Basse-Terre, and Petite-Terre) during the Pleistocene
111 when sea levels were 100 meters lower (Fig. 1–A). PGR6 contains a stratified fossil-bearing deposit
112 that has been the object of two excavations, the first in 2011 led by M. Boudadi-Maligne (Boudadi-
113 Maligne *et al.* 2016), and the second in 2016 led by A. Lenoble. The still largely unpublished fossil
114 assemblage from the more recent excavations produced 191 of the 218 *Leiocephalus* bones found from
115 the site. The fossil material was collected during dry sieving in 2011 and subsequent wet sieving of the
116 sediment using 2mm² mesh sieves in 2016. The sieved sediment was recorded by successive, 5cm
117 thick horizontal spits (*décapages*) during excavations.

118 The 2016 excavations helped clarify the chronology of the four stratigraphic units identified during the
119 initial work at the site (Fig. 1–B). An initial radiocarbon date obtained on a sample recovered during
120 the 2011 excavations provided an age of 299-507 cal. AD for the base of the layer 3. Two additional
121 radiocarbon dates, one on charcoal (Lyon-14287 –SacA-50616-) recovered from the base of layer 4
122 (square B10d, *décapage* 22) provided a date of 170 BC – 4 cal. AD (2060±30 BP), the other on a
123 Yellow-crowned night-heron bone (Lyon-14478 -SacA-51014-) from the top of the layer 3 (square
124 B10d, *décapage* 11) produced a date of 1297-1373 cal. AD (600 ± 30 BP). These new dates are in
125 good agreement with the initial chronological interpretation of the stratigraphy based on the
126 archaeological material (pottery sherds and bones of introduced species). Layer 1 has been attributed
127 to modern activity over the last 200 years, layer 2 to the Colonial period, and layers 3 and 4 to the pre-
128 Columbian Ceramic period between around 200 BC and 1728 AD. The pre-Columbian Ceramic era is
129 divided into different sub-periods spanning from 80 AD to around 1600 AD in Guadeloupe
130 (Fitzpatrick 2015; Hofman *et al.* 1999; Keegan *et al.* 2013) and to 1728 AD in La Désirade (Boudadi-
131 Maligne *et al.* 2016). The few ceramic remains recovered from PGR6 do not provide any additional
132 chronological resolution (for details see Boudadi-Maligne *et al.* 2016).

133

134 CT-scanning

135 We produced high-resolution 3D models of the head of the dry specimen from the Bordeaux Natural
136 History Museum in order to describe its skull morphology without damaging the specimen. Micro-CT
137 imaging was performed using a GE v|tome|x s μ CT scanner at the UMS 3626 PLACAMAT laboratory
138 of the University of Bordeaux (France) with acquisition parameters set to 100 kV (tension), 200 μ A
139 (Intensity), and 333 ms (exposure time) for a total of 2550 projections. A 0.1 mm copper filter was
140 used to reduce beam hardening artifacts. The geometry was set to obtain a 13.7 μ m voxel size in the
141 reconstructed three-dimensional images. The reconstruction was generated using the phoenix datos|x
142 v. 2.2.1.716 scanning software. Each bone of the specimen was manually segmented and
143 individualized by creating a sub-set of selected voxels in AVIZO v. 9.3 (VSG, SAS, Merignac,
144 France, <http://www.vsg3d.com>). This final manipulation was impossible to perform automatically
145 given the presence of a metallic rod within the head of the specimen, which produced visual artifacts
146 that we had to remove manually.

147 Phylogenetic analysis

148 Phylogenetic analysis was conducted using the characters and character matrix of Pregill (1992),
149 completed with the data we collected from the dry specimen and fossil bones from Guadeloupe. The
150 theoretical ancestor of Pregill (1992) was included in the analysis but, as it included undetermined (?)
151 character states, we incorporated two additional “ancestors” with either 0 or 1 character states as out-
152 group taxa. The most parsimonious trees were determined from the obtained matrix (Appendix 1)
153 using a parsimony branch and bound search with the PAUP4 v.3.99.167.0 software
154 (<https://paup.phylosolutions.com>). We then computed a consensus of the trees obtained.

155 Anatomical descriptions

156 The description of the external anatomy of the Holotype follows the same terminology as Pregill
157 (1992). The osteological description mostly follow the nomenclature Fejérvary-Langh (1923) and

158 Evans (2008). This terminology was complemented by other references (Bailon 1991; Klembara *et al.*
159 2010; Oelrich 1956; Smith *et al.* 2018), some of which are mentioned in the text when needed.

160

161 **Results**

162 Comparison of both the dry specimen from the Natural History Museum of Bordeaux (MHNBx) and
163 the fossil remains from Guadeloupe with existing data concerning the morphology of known
164 *Leiocephalus* species indicates the Guadeloupe specimens to bear both unique morphological
165 characters and combinations of morphological characters that exist in no other known *Leiocephalus*
166 species. As such, we have attributed the Guadeloupe specimens to a new species, *Leiocephalus*
167 *roquetus* sp. nov., for which we provide a formal description and complete diagnosis.

168 *Leiocephalus roquetus* sp. nov.

169 Fig. 2, 3,4,5,6, 7

170 **Diagnosis.** *Leiocephalus roquetus* differs from all other congeners by the morphology of its parietal
171 bone that bears adductor crests fused into a single crest in the posterior area of the bone in large
172 specimens. In addition of this character, *L. roquetus* presents a unique combination of characters:
173 absence of enlarged snout scales, and occurrence of five scales in contact with the first pair of enlarged
174 frontal scales (pattern of type I of Pregill (1992)), the occurrence of three internasal scales, a
175 prominent dorsal body crest, absence of keeled ventral scales, frontal bone exposed dorsally between
176 the nasal and prefrontal bones, skull rugosities on the frontal bone, premaxillary spine constricted just
177 above the dentigerous process, a well-developed posterior process of the septomaxilla, a distinct dorsal
178 process of the squamosal, a well-developed angular process of the dentary, and transition from
179 bicuspid to tricuspid teeth occurs between the height and sixth most anterior dental position of the
180 dentary. This new taxon also presents several differences with other known Lesser Antillean
181 *Leiocephalus* specimens (see below).

182 **Derivatio nominis:** The species is named in reference to the first common name “roquet” attributed
183 by Du Tertre (1654) and de Rochefort (1658) to the leiocephalid lizards of Guadeloupe. As pointed
184 out by Breuil (2002), this name was latter erroneously attributed to *Anolis* lizards and, as the
185 Amerindian name of *Leiocephalus* is unknown, we choose to retain the original name given to these
186 squamates. The English name of this lizard would be “Curlytail roquet” and the French name
187 “Léiocéphale roquet”.

188 **Type locality:** Guadeloupe, ? La Désirade Island, Lesser Antilles.

189 **General description of the holotype.** The holotype is a complete, dry taxidermy specimen (labeled
190 MHNbx 2002.1871) preserved in the Natural History Museum of Bordeaux (France). This specimen
191 has a snout-vent length (SVL) of 10.8 cm and measures 26.5 cm in total length (Fig. 2).
192 Microtomography reveals the skull to be fully preserved (Fig. 3). The key morphological features of
193 the holotype are as follows: enlarged nasal bones, the narrow nasal process of the premaxilla is
194 triangular, the dorsal part of the premaxilla is fully enclosed by the nasal bones, the premaxilla lacks
195 lateral spines and bears 7 tooth positions, the nasal-maxillary suture is arched toward the mid-line of
196 the snout, the nasal process of the frontal bone is dorsally exposed between the nasal and prefrontal
197 bones, the pterygoid bears teeth but palatine teeth are absent, lacrimal and postfrontal bones are
198 present, the septomaxilla bears a quadrangular lateral wing with a thin posteriorly oriented process, the
199 frontal is narrow with dermal ornamentation, the parietal foramen is on the fronto-parietal suture, the
200 adductor crests of the parietal table converge but do not touch and are posteriorly separated by a
201 depressed area, the anterior part of the parietal table presents a dermal ornamentation, the
202 supratemporal bone is on the lateral side of the supratemporal process of the parietal, the
203 supratemporal process of the squamosal is distinct, splenial and angular bones are present, the angular
204 process of the dentary is well-developed, most of the teeth are tricuspid and distinctively flared, the
205 transition from monocuspid/bicuspid to tricuspid teeth occurs at the seventh dental position on the
206 dentary and at the fourth dental position on the maxilla. Head scales are large, the specimen presents
207 four rows of snout scales of sub-equal size between the internasal and the anterior pair of frontal
208 scales, there are four moderately-sized parietal scales and the most lateral scales are smaller than the

209 median scales, two rows of post parietal scales are present, there is no enlarged lateral post-parietal
210 scale, there are three internasal scales, there are four lorilabial scales anterior to the first lorilabial
211 contacting the elongated sub-ocular scale, the cephalic scales lack ridges, the temporal scales are of
212 sub-equal size and no elongated temporal scales are present, the lateral nuchal scales are smaller than
213 surrounding body scales, the lateral body scales are the same size as dorsal and ventral scales, a
214 prominent middorsal body crest formed by overlapping scales is present, the dorsal crest (occiput to
215 vent) is composed of 53 scales, the tricarnate scales at the base of the first and second toes grow into
216 comb-like fringes but are not prominent, the venter is dull and patternless, there is no scapular patch,
217 no suprascapular blotches, no facial band, and the base of tail is laterally compressed.

218 **Description of paratype fossil bones and corresponding bones of the holotype.** In addition to the
219 holotype dry specimen, we associate, as paratypes, several fossil bones collected from the Pointe Gros
220 Rempart 6 deposit to the type series of *L. roquetus*. These bones are described below in association
221 with corresponding bones of the holotype specimen in order to demonstrate that both modern and
222 fossil materials correspond to the same taxon.

223 **Premaxilla**

224 **Holotype** (MHNbx 2002.1871) (Fig. 4–A): The premaxilla is a single bone and measures 2.8 mm
225 across the rostrum. In posterior view, the supradental shelf tends to extend moderately both laterally
226 and posteriorly. In anterior view, the bone presents a single incisive process oriented ventrally,
227 reaching half the height of the maxillary process. The nasal process is high and triangular with the
228 thinner dorsal part fully covered by imprints of the nasal bones. The portion of the nasal process that is
229 not covered by the nasal bones on the articulated skull is triangular with a slight constriction at the
230 base. The nasal process lacks lateral spikes *sensu* Pregill (1992). The imprints of the maxilla are
231 weakly extended on the lateral side of the maxillary processes (lateral processes *sensu* Oelrich 1956).
232 The bone bears seven tooth positions occupied by conical, monocuspid pleurodont teeth. The
233 morphology of these teeth cannot be further described as the majority are broken, probably during the
234 preparation the specimen.

235 **Paratype** “PGR6. square C11c. *décapage* 14.XII” (Fig. 4–B): This element is a complete subfossil
236 premaxilla collected from layer 3 of PGR6 (square C11c; *décapage* 14). This element measures 4.2
237 mm across the rostrum. This bone is similar to the holotype and allows further detail to be described
238 for the morphology of the teeth, which are monocuspid, blunt, and slightly pointed on this specimen.
239 As in the holotype and every premaxilla from PGR6, this element does not exhibit any dermal
240 ornamentation.

241 Additional material: Two additional well-preserved premaxillas were recovered from the PGR6
242 deposit and are identical to the holotype and paratype described above. The teeth preserved on these
243 bones are all monocuspids.

244 **Maxilla**

245 **Holotype** (MHNbx 2002.1871) (Fig. 4–C): The bone bears a 12 mm long dental row bearing 21 tooth
246 positions. The most anterior tooth is conical, the second and third are bicuspid with a large posterior
247 cusp and a reduced anterior cusp. The following teeth are tricuspid (with a dominant central cusp),
248 linguolabially compressed, distinctively flared in mesio-distal direction. The shape of these teeth is
249 similar to that of tricuspid teeth of the dentary (see below). In medial view, the supradental shelf (
250 *sensu* Rage & Augé 2010) is slightly arched toward the mid-length of the bone. The facial process is
251 triangular, longer than it is high. The posterior process is short, and is in equivalent in length to the
252 portion of the supradental shelf bearing the two last dental positions, and its posterior tip is
253 quadrangular in dorsal view. In labial view, the antero-dorsal margin of the facial process is strongly
254 inflected medially. The surface of the facial process is punctuated by several small foramina, and six
255 larger labial foramina are present. The bone lacks dermal ornamentation. In dorsal view, the
256 supradental shelf is triangular. The internal premaxillary process is more developed than the external
257 one. This process is high and thin, and its antero-dorsal margin is linked to the medial margin of the
258 facial process by a crest visible in medial view. A moderately deep V-shaped premaxillary fenestra
259 separates the internal and external premaxillary processes.

260 **Paratype** “PGR6. square B10d. *décapage* 22.III” (Fig. 4–D): This element is a complete subfossil
261 maxilla recovered from layer 4 of PGR6 (square B10d; *décapage* 22). This bone, although similar to
262 the holotype, nevertheless presents several differences. This maxilla has a 15.4 mm long dental row
263 with 22 tooth positions. The most anterior tooth is bicuspid and the most anterior tricuspid tooth
264 occupies the fifth tooth position.

265 Additional material: Six additional, well-preserved maxillae were recovered from the PGR 6 deposit.
266 These bones are similar to those described above but present variable numbers of tooth positions,
267 ranging between 17 for the smallest specimens (dental row length = 9 mm) to 22 for the largest
268 individuals (dental row = 13.56–16.3 mm). The most anterior tricuspid tooth of these specimens
269 appears at the fourth or fifth tooth position.

270 **Frontal bone**

271 **Holotype** (MHNbx 2002.1871) (Fig. 5–A): The two frontal bones are fused (azygous). In dorsal view,
272 the bone is elongated, 2.5 times wider in its posterior than in its anterior part, with a nearly straight
273 parietal margin. The minimal width of this element at the level of its median constriction is 3.1 mm.
274 The bone exhibits a moderately marked tuberculated dermal ornamentation on the entire dorsal surface
275 of the bone. The parietal foramen is present at the posterior end of the bone and partially opens onto
276 the fronto-parietal suture. In dorsal view, the anterior end of the bone presents deep imprints of the
277 nasal bones. Nasal and prefrontal bones are independent in the anterior area of the frontal, as
278 demonstrated by the occurrence of two well-marked crests on the nasal process separating imprints of
279 the nasal and prefrontal bones. In ventral view, the cranial crests (= *crista cranii*) are well
280 individualized but are not prominent, and are independent of in the median area of the bone. The
281 postfrontal bone imprint is visible at the anteromedial base of the posterolateral processes. Nearly the
282 entire anterior half of the lateral margin of the bone is occupied by an imprint of the prefrontal bone,
283 which is fully visible only in lateral view.

284 **Paratypes** “PGR6. square B10d. *décapage* 13.IV” (Fig. 5–B), and “PGR6. square B10d. *décapage*
285 13.V” (Fig. 5–C): These bones were both found in layer 3 of PGR6 (square B10d, *décapage* 13) and

286 correspond to a large individual with a minimal width of 4.9 mm and a small specimen, probably a
287 juvenile, with a minimal width of 2.2 mm. These specimens are morphological similar to the holotype.

288 **Parietal bone**

289 **Holotype** (MHNbX 2002.1871) (Fig. 5–D): The parietal bone is a single element and has maximal
290 antero-posterior length of 9.5 mm. In dorsal view, the medial area of the bone is slightly laterally
291 constricted . A slight central depression of the fronto-parietal suture of the bone indicates the position
292 of the parietal foramen whose posterior margin is delimited by the parietal. The adductor crests are
293 low and tend to converge posteriorly but are parallel in the posterior part of the bone. In the anterior
294 part of the bone these crests delimit a U-shape, heavily ornamented area comprising elongated
295 tubercles. In the posterior part of the bone, the adductor crests are separated by a slight postero-medial
296 depression. The supratemporal processes are well individualized and extend posteriorly. Deep and
297 clearly distinct nuchal fossae are present on the medial side of both supratemporal processes. These
298 fossae are separated medially by a small longitudinal crest overlying the parietal fossa (recessus
299 processi ascendentis *sensu* Smith (2011)). In lateral view, an imprint of the supratemporal bone covers
300 the full length of the supratemporal processes.

301 **Paratypes**: “PGR6. square C11d. *décapage* 12.VI” (Fig. 5–E): This bone was found in layer 3 of
302 PGR6, (square C11d, *décapage* 12). This element is a small medial fragment of a parietal bone,
303 similar in size to the holotype. The morphology of this element is similar to the holotype, including the
304 occurrence of weakly defined adductor crests separated by a small depression in the posterior part of
305 the bone.

306 “PGR6. square B9b. *décapage* 22.VII” (Fig. 5–F): This bone was found in layer 4 of PGR6 (square
307 B9b, *décapage* 22). This element is a nearly complete parietal in a more advanced ontogenetic stage
308 than the holotype and paratype described above. It is larger than the other parietal bones we identified,
309 with a maximal antero-posterior length of 12.3 mm. While similar to the holotype, the morphology of
310 the parietal table and adductor crests is completely different. In addition to being high and well

311 defined, the crests delimit a V-shape ornamented area and meet in the posterior part of the bone to
312 form a single crest reaching the postero-median extremity of the parietal table.

313 Specific remarks regarding the attribution of the paratype “PGR6. square B9b. *décapage* 22.VII” to
314 *Leiocephalus roquetus*: The morphology of this element is substantially different from the holotype,
315 leading us to explore whether it formed part of the morphological variability of the same taxa or
316 reflects a different species. This is all the more important considering that the morphology of this
317 particular parietal bone differs from all known *Leiocephalus*. The attribution of the paratype “VII” to
318 *Leiocephalus* is however supported by its numerous similarities with the holotype, including the
319 overall shape of the bone, the morphology of the supratemporal processes and parietal fossae, the
320 absence of parietal foramen, and the occurrence of a well-delimited ornamented area in the antero-
321 medial area of the bone. In addition, similar ontogenetic parietal morphological variability exists in
322 several other pleurodont squamates (e. g. *Anolis* -Bochaton *et al.* 2017-, *Iguana* -Bochaton *et al.* 2019-
323) including *Leiocephalus* itself (Pregill 1992) although the adult morphology usually differs from our
324 paratype “VII”. However, the adductor crests in *Leiocephalus* usually converge in the posterior area of
325 the bone. We included this bone in the morphological variability of *L. roquetus* given the occurrence
326 of the smaller parietal paratype “VI”, similar to the holotype, in the same deposit, which reinforces our
327 interpretation of paratype “VII” representing an older specimen with a more ontogenetically advanced
328 morphology.

329 **Pterygoid bone**

330 **Holotype** (MHNbx 2002.1871) (Fig. 6–A): The bone presents a long, thin posterior process
331 accounting for half of the length of the complete bone. In dorsal view, the transverse process extends
332 moderately laterally and bears a well-marked, triangular articular surface with the ectopterygoid. The
333 base of the posterior process has a deep pit for the insertion of the epipterygoid bone. In dorsal view,
334 the palatine process forms a long and relatively thin blade whose pointed anterior tip bears a well-
335 defined contact area with the palatine bone. In medial view, the anterior extremity of the posterior
336 process presents a deep indentation occupied by a well-marked articular surface for the basipterygoid

337 process of the basisphenoid bone. Posteriorly to this structure, an oval pterygoid furrow occupies most
338 of the medial area of the posterior process. In ventral view, the palatal process present superficial
339 traces of teeth sockets (*sensu* Mahler & Kearney 2006), and two teeth were present on the left
340 pterygoid of the holotype specimen. However, these teeth were not attached to the bone by any
341 ossified structure and were absent on the right pterygoid. It is therefore difficult to be certain if the
342 absence of pterygoid teeth is a general trait of *L. roquetus* as this character could reflect intra-
343 individual and intra-specific variability. Such variability was previously observed by Mahler &
344 Kearney (2006) in the genus *Leiocephalus*.

345 **Paratypes:** “PGR6. square C11d. *décapage* 14.X” (Fig. 6–B): This bone, recovered from layer 4 of
346 PGR6 (square C11d, *décapage* 14), is similar to the holotype but it lacks any trace of teeth sockets.
347 Instead, the anterior part of the palatal process bears two short bulbous processes of unknown nature in
348 places that would normally be occupied by teeth. We are unable to determine whether these processes
349 are teeth or bony formations. The morphology of the palatal process displays some variability in the
350 PGR6 fossils. This process can be more or less depressed and usually lacks teeth-related structures,
351 although the smallest fossil does present a single small bulbous process similar to those previously
352 evocated. Considering that the holotype pterygoid is much smaller than any of the fossil pterygoid
353 from PGR6, we assume that the occurrence of teeth on the pterygoid in *L. roquetus* may reflect
354 ontogenetic or individual variability.

355 **Squamosal bone**

356 **Holotype** (MHNbX 2002.1871) (Fig. 6–C): In lateral view, the bone presents a long, narrow anterior
357 ramus and two shorter posterior processes. The dorsal process (supratemporal process) is three times
358 longer than the ventral process (quadrate process) and is clearly distinct. The tip of this process is
359 rounded and the whole process is slightly oriented anteriorly. The ventral process is reduced and
360 pointed.

361 **Paratypes:** “PGR6. square B9b. *décapage* 22.IX” (Fig. 6–D): This bone was found in layer 4 of PGR6
362 (square B9b, *décapage* 22). This bone, while similar to the holotype, is larger and presents a clearly
363 more developed ventral process. The ventral process is broken.

364 **Septomaxilla**

365 The septomaxillae of the holotype are very thin which make difficult to observe their morphology in
366 detail using the CT-scan data and to capture good images of it. We were however able to observe the
367 main components of its morphology that are describe hereinafter. This bone was not found in the fossil
368 material.

369 **Holotype** (MHNbx 2002.1871) (not figured): The septomaxilla presents a well-developed blade-like
370 posterior process and a flat well-extended triangular anterior process 1.5 times longer than wide. The
371 dorsal surface of this process is slightly concave. The septal process is poorly developed.

372

373 **Quadrate bone**

374 **Holotype** (MHNbx 2002.1871) (Fig. 6–E): The bone is 5 mm high and sub-rectangular in anterior and
375 posterior views. In posterior view, it is divided into two asymmetrical portions by a posterior crest.
376 The lateral region is the most developed and comprises a deep conch delimited by a tympanic crest.
377 This region is dorsally notched to receive the ventral process (quadrate process) of the squamosal. The
378 medial region has a weakly-developed pterygoid lamina whose width is similar to or slightly thinner
379 than the posterior crest. In anterior view, the dorsal part of the tympanic crest region is occupied by a
380 weakly marked antero-dorsal depression.

381 **Paratypes:** “PGR6. square C11c. *décapage* 15.XI” (Fig. 6–F): This bone comes from layer 3 of PGR6
382 (square C11c, *décapage* 15). This bone is similar to the holotype but is larger (7.7 mm high) and its
383 antero-dorsal depression is more strongly marked and delimited than in the holotype which could
384 reflect ontogenetic variability.

385 **Dentary**

386 **Holotype** (MHNbx 2002.1871) (Fig. 7–A): The bone is straight and elongated. Its dental row is 13
387 mm long and bears 25 teeth. The six most anterior teeth positions bear simple mono-pointed conical
388 teeth, remaining teeth are tricuspid, linguolabially compressed, distinctively flared in mesio-distal
389 direction, and present large and pointed secondary cusps. In lateral view, the symphysis is thin and
390 oblong. The surangular and angular processes are well-developed and extend posteriorly well beyond
391 the level of the dental row. The surangular process extends posteriorly well beyond the apex of
392 coronoid. The angular process is more pointed than the surangular process, which is blunter and
393 shorter. Its length is equivalent to two-thirds the surangular process. The dentary also presents a well-
394 individualized coronoid process extending inside the coronoid bone in the complete mandible. An
395 intra-mandibular septum is presents below the dental row, between the last and fourth-to-last dental
396 positions. An intramandibular lamella (*sensu* Smith 2009) is present and well-developed. In medial
397 view, a large imprint of the anteromedial process of the coronoid bone is visible on the posterior
398 extremity of the bone and extends below the three last dental positions of the dental row. In ventral
399 view, the Meckelian groove is nearly fully enclosed by the dorsal and ventral flanges of the dentary
400 that are fully fused on the three-quarters of the length of the dental row. The anterior extremity of the
401 Meckelian groove remains open by a slender opening below the first to seventh most anterior tooth
402 positions.

403 **Paratypes**: “PGR6. square C11d. *décapage* 16.I” (Fig. 7–B): This bone is a nearly complete subfossil
404 dentary missing only its posterior extremity. This bone was recovered from layer 3 of PGR6 (square
405 C11d; *décapage* 16). This bone is similar to the holotype however the most anterior tricuspid tooth
406 appears at the 8th tooth position. It also bears less dental positions (n=21) due to its shorter dental row
407 (14.6 mm). The anterior opening of the Meckelian groove is also less posteriorly extended than on the
408 holotype as it extends only below the five most anterior dental positions.

409 “PGR6. square C11c. *décapage* 11.II” (Fig. 7–C): This bone is a nearly complete subfossil dentary
410 missing its posterior extremity and whose anterior part is dissolved, probably due to it being digested
411 just as several other bone remains in the fossil assemblage presenting characteristic marks of digestion
412 (Fernández-Jalvo & Andrews 2016). This bone was also found in layer 3 of PGR6 layer (square C11c;

413 *décapage* 11) and exhibits a similar dental length (14.4 mm) and number of teeth (n=25) to the
414 holotype.

415 Additional material: Two additional but less well-preserved dentaries were also recovered from PGR6.
416 Dental rows measure 14.7 mm and 12 mm and bear, respectively, 21 and 20 tooth positions. The
417 smallest dentary presents a juvenile morphology and the most anterior tricuspid tooth occupies the
418 sixth most anterior tooth position.

419 Articular

420 **Holotype** (Fig. 7–D): In dorsal view, the elongated articular is four times longer than wide with well-
421 developed articular and retroarticular processes. The articular process is triangular with a medially
422 oriented apex. The retroarticular process is rectangular and separated from the articular process by a
423 well-marked ridge. The articular process extends 2 mm medially from the medial border of the
424 condyle.

425 **Paratype**: “PGR6. square B9b. *décapage* 17.VIII” (Fig. 7–E): This element was recovered from layer
426 3 of PGR6 (square B9b, *décapage* 17) and comprises complete and fused articular and surangular
427 bones. The retroarticular process is almost square in dorsal view. The articular process is similar to
428 that of the holotype but is more extended medially. The limit between the articular and the
429 retroarticular processes is not marked by any structure and is therefore difficult to delimit. The
430 articular process extends 4.1 mm medially from the medial border of the condyle.

431 Specific remarks regarding the morphological variability of the articular bone: The holotype and the
432 largest articular bone found from PGR6 (Paratype “PGR6. square B9b. *décapage* 17.VIII”) differ
433 somewhat in regards the morphology of the articular and retroarticular processes. These processes are
434 widened medially in the largest specimen, and the limit between the two processes becomes
435 indistinguishable. As other pleurodont squamate genera display a similar degree morphological
436 variability, we interpret this difference as reflecting ontogenetic variability in *Leiocephalus roquetus*.

437 Additional remarks regarding fossil post-cranial elements:

438 In addition of the cranial elements described above, the PGR6 deposit produced several postcranial
439 elements of *Leiocephalus* that we attribute to *L. roquetus*. However, as comparing these elements with
440 the holotype specimen was impossible, we describe their main morphological characters without
441 considering these bones as type material.

442 **Dorsal Vertebrae:** We attributed forty-seven dorsal vertebrae recovered from PGR6 to *Leiocephalus*
443 *roquetus*. The morphology of these vertebrae varies depending on their position on the vertebral
444 column, as indicated by the morphology of the synapophyses. One of the most significant differences
445 is the morphology of the hypapophysis, which is narrow and forms a longitudinal ridge in cervical and
446 anterior trunk vertebrae and becomes uniformly wide and flat in median and posterior trunk vertebrae.
447 However, all complete vertebrae, independent of their position on the column, present a high, vertical
448 neural spine (Fig. 7–F). This combination of characters is typical of most *Leiocephalus* species (Pregill
449 1992).

450 **Caudal vertebrae:** The ten caudal vertebrae recovered from PGR6 present a slender medial spine
451 projected vertically above the neural arch at the position of the fracture plane. This character is an
452 apomorphy in *Leiocephalus* (Etheridge 1966; Pregill 1992).

453 **General remarks:**

454 The diagnosis of *L. roquetus* is based on a combination of a single historical dry-specimen (holotype)
455 and fossil bone remains (paratypes) sharing common anatomical characteristics and likely
456 geographical origin. Both of these types of material would have independently led to the same
457 conclusion (the description of a new species) but the combined information they provide allow for a
458 complete assessment of the external morphology and osteological ontogenetic variability of *L.*
459 *roquetus* sp. nov. The combined consideration of the holotype juvenile dry-specimen and the
460 paratypes fossil remains from PGR 6 is enabled by the strict osteological similarity between these
461 materials as it is demonstrated by our cladistic analysis.

462 *L. roquetus* presents a combination of characters shared with *L. herminieri* from Martinique, which
463 would place it in a hypothetical Lesser Antillean clade (see below): the absence of enlarged snout

464 scales and the occurrence of five scales in contact the first pair of enlarged frontal scales (the type I
465 pattern of Pregill -1992-), frontal bone exposed dorsally between the nasal and prefrontal bones, skull
466 rugosities on the frontal bone, and a prominent dorsal body crest. The fossil species *L. cuneus* from
467 Antigua bears the two osteological characters shared between *L. roquetus* and *L. herminieri*, however,
468 as these characters also occur in Greater Antillean taxa, it is impossible to link *L. cuneus* to a putative
469 Lesser Antillean clade. Moreover, such a hypothesis is not supported by our phylogenetic analysis (see
470 below).

471 Although potentially forming part of the same clade, *L. roquetus* presents several morphological
472 differences with the two other *Leiocephalus* described in the Lesser Antilles in addition to the
473 morphology of its parietal bone.

474 - with *L. herminieri*: absence of keeled ventral scales, occurrence of three internasal scales,
475 premaxillary spine constricted just above the dentigerous process, occurrence of well-developed
476 posterior process of the septomaxilla, occurrence of a distinct dorsal process of the squamosal,
477 occurrence of a well-developed angular process of the dentary, the most anterior occurrence of
478 tricuspid teeth between the 6th and the 9th most anterior dental position of the dentary.

479 -with *L. cuneus*, the only other fossil *Leiocephalus* described in the Lesser Antilles and whose past
480 occurrence was suspected in Guadeloupe: absence of bicuspid teeth on the premaxilla, absence of
481 dermal ornamentation on the premaxilla, premaxillary spine constricted just above the dentigerous
482 process, occurrence of “teeth” on the pterygoid bones of some specimens, absence of tricuspid teeth on
483 the 4th and 5th dental positions of the dentary, absence of a V-shaped indentation separating the angular
484 from the retroarticular process of the articular bone.

485 In terms of the size of *L. roquetus*, although the holotype specimen is moderate in size (snout-vent
486 length of 10.8 cm), most of the fossil bones attributed to this species were larger, indicating that the
487 type specimen is likely to be a juvenile. Isolated bones do not, however, provide direct evidence for
488 the size of the complete specimen, and we can only provide a rough estimate of the putative size of
489 adult *L. roquetus*. The fossil bones from PGR6 are of similar size to the largest bones reported by R.

490 Etheridge (1964) to estimate the size of *L. cuneus* (200 mm SVL). Certain elements are both smaller
491 (premaxilla, quadrate, dentary, articular) or larger (maxilla, frontal, parietal), leading us to estimate *L.*
492 *roquetus* to be roughly of similar size as *L. cuneus*, which is currently the largest known *Leiocephalus*
493 species.

494 **Phylogenetic relationships of *L. roquetus* in regard to other *Leiocephalus* species**

495 To complement our morphological analysis of *L. roquetus*, we placed this newly described taxon in
496 the only available morphological phylogeny of *Leiocephalus* (Pregill 1992). The parsimony analysis
497 generated 177 optimal trees of 130 steps (Consistency Index -CI- =0.408 and Retention Index -RI-
498 =0.603). We then computed the consensus of these trees (Fig. 8). Our parsimony analysis produced
499 almost exactly the same results as those reported by Pregill (1992) with similar data (CI=0.441,
500 RI=0.590). The addition of additional ancestors did not modify the topology of the tree obtained by
501 Pregill (1992). The basal position of *L. herminieri* inferred by Pregill (1992) in regard to other
502 *Leiocephalus* is partly confirmed by our analysis but this part of the tree is more poorly resolved than
503 that of the previous study. Our newly described *L. roquetus* appears to be the closest relative of this
504 also extinct, taxon. Our results indicate the existence of a putative but now completely extinct Lesser
505 Antillean clade, which might reflect an individual colonization event of the Lesser Antilles that may
506 predates the colonization of the Greater Antilles. This hypothesis could be seen as in contradiction
507 with the position of *L. cuneus*, which is not part of this putative Lesser Antillean group despite having
508 been described in this area (primarily Antigua and Barbuda). This could be explained either by the low
509 number of characters it was possible to record on *L. cuneus* due to the fact that only part of its
510 osteology and none of its external morphology is currently known. However, we also performed a
511 second analysis treating the complete holotype specimen and the fossil paratypes of *L. roquetus*
512 independently (CI=0.405, RI=0.608). This analysis confirmed the holotype and paratype specimens to
513 be more closely related to each other than to any other taxon and that *L. cuneus* was not part of that
514 clade. There is thus also a possibility that *L. cuneus* has a Greater Antillean origin and is related to a
515 different colonization event. However, the paucity of available fossil data makes it impossible to
516 explore this question further.

517 **The past biodiversity of *Leiocephalus* in the Guadeloupe Islands**

518 Fossil evidence complemented by the historical specimen presented here allow us to describe an
519 endemic and recently extinct species of *Leiocephalus* from Guadeloupe, whose past distribution in
520 Guadeloupe remains unclear. *L. roquetus* was definitely present on La Désirade, which forms part of
521 an island group comprising Grande-Terre, Basse-Terre, and Petite-Terre. This could suggest that these
522 islands were occupied by a single species, as is generally the case for Lesser Antillean squamate taxa
523 (Henderson & Powell 2009). However, the exact geographic origin of the type specimen of *L.*
524 *roquetus* is unclear, and fossil occurrences of *Leiocephalus* are extremely rare in Guadeloupe. The
525 occurrence of *L. cuneus* in Guadeloupe (Grande-Terre and La Désirade) had been previously
526 suggested based uniquely on a small sample of fossil dentary bones on which the most anterior
527 tricuspid tooth appears in the fifth most anterior dental position (Pregill 1992; Pregill *et al.* 1994). In
528 addition to this material not be available for study, the lack of well-preserved *Leiocephalus* remains in
529 the archaeological records of Grande-Terre and Basse-Terre makes evaluating this possibility difficult
530 (Bochaton 2016). However, a premaxilla reported by Bochaton (2016) from the archaeological site of
531 Anse à la Gourde (Grande Terre) lacked both bicuspid teeth and the dermal ornamentation typical of
532 *L. cuneus*. The taxonomic identity of the *Leiocephalus* species which inhabited the main Guadeloupe
533 islands therefore remains an open question. Regarding the putative occurrence of *L. cuneus* on La
534 Désirade, we restudied the dentary described by Boudadi-Maligne *et al.* (2016). Our analysis found
535 that the most anterior tricuspid tooth of this bone appears on the sixth and not on the fifth most
536 anterior dental position, as initially reported, making it similar the dentary of *L. roquetus*. Regarding
537 Marie-Galante, despite its rich fossil record, *Leiocephalus* remains are extremely rare in the
538 archeological and paleontological deposits. The few available bones all have the most anterior
539 tricuspid tooth on the fifth most anterior dental position on the dentaries (Stouvenot *et al.* 2014;
540 Bochaton pers. obs.), a trait absent from *L. roquetus*. However, the unpublished premaxilla from Cadet
541 3 on Marie-Galante (Bochaton pers. obs.) lacks bicuspid teeth and the dermal ornamentation typical of
542 *L. cuneus* , making it impossible to definitively attribute it to this species. In the end, while we can be

543 certain that *L. roquetus* previously inhabited La Désirade, the fossil material is too scarce to identify
544 fossils from the other islands of Guadeloupe to specific taxa.

545 **Discussion**

546 Our works add a third species to the list of now-extinct Lesser Antillean *Leiocephalus*. This very low
547 species count compared to other Lesser Antillean lizards, especially anoles, is likely to be a heavy
548 under-representation of the real past diversity of these lizards.

549 Causes and timing of the extinction of *Leiocephalus* in the Lesser Antilles

550 While the extinction of *Leiocephalus* during the colonial period in Guadeloupe was previously
551 suggested by archaeological evidence (Bochaton 2016), the specimen housed at the Natural History
552 Museum of Bordeaux, collected in the first half of the 19th century, adds more precision to the last
553 occurrence date of this lizard. An early 19th century extinction of this lizard would be consistent with
554 the last mention of *Leiocephalus* in Martinique, demonstrating the species to have disappeared from
555 both islands only very recently. The potential causes underlying these extinctions could be the same as
556 those frequently advanced for squamates: the impact of introduced mammalian predators (Henderson
557 1992; Lorvelec *et al.* 2007), human-induced changes to landscapes and intensive agricultural practices
558 (Boudadi-Maligne *et al.* 2016; Corke 1992), or a combination of several of these factors (Iverson
559 1978). The general causes for the regional extinction of *Leiocephalus* are far from clear. This genus
560 was probably among the most heavily impacted squamate genera, as it is the only genus that was
561 completely wiped out from the Lesser Antilles sometime after the Late Pleistocene. The second most
562 impacted genera, *Boa* and *Diploglossus*, also present in Guadeloupe in the past, still have relic
563 populations on respectively Dominica and Montserrat islands (Corry *et al.* 2010; Henderson & Powell
564 2009). Extant *Leiocephalus* mainly inhabit littoral areas, which potentially contributed to the early
565 demise of the genus in the Lesser Antilles (Breuil 2009), especially as the dry forests typical of these
566 environments were heavily exploited by humans from the outset of the colonial period (Lugo *et al.*
567 1981). However, the key to fully understand the history of *Leiocephalus* requires the construction of a
568 precise regional chronology, unfortunately and although well-documented in Guadeloupe and

569 Martinique, the putative occurrence and extinction dates of *Leiocephalus* on the other Lesser Antillean
570 islands is far from clear.

571 In Anguilla, *Leiocephalus* is estimated to have gone extinct in the mid-Holocene
572 (Roughgarden & Pacala 1989) based on data collected from a single site. However, this appears
573 insufficient to support a mid-Holocene extinction, as the occurrence of a given taxa in a natural or
574 archaeological deposit can be influenced by several factors (e.g. accumulation agent, changes in local
575 environmental conditions). The additional mention of *Leiocephalus* on Anguilla (Pregill *et al.* 1994)
576 remains undated, exposing one of the main issues for the study of fossil *Leiocephalus* in the Lesser
577 Antilles; the extreme rarity of remains in faunal assemblages from archaeological deposits dating to
578 the two last millennia. In fact, apart from the Indian Town Trail site in Barbuda (Watters *et al.* 1984)
579 and several unpublished deposits in Guadeloupe investigated by one of us (Bochaton 2016),
580 *Leiocephalus* remains have never been reported from archaeological deposits in the Lesser Antilles.

581 The lack of dated fossils and their rarity raises two possibilities; 1) *Leiocephalus* species were
582 almost entirely extinct more than two thousand years ago and are thus logically absent from Lesser
583 Antillean archaeological deposits that predate the human colonization of the islands or, 2) their
584 absence reflects biases in the identification of fossil remains or the composition of faunal assemblages
585 from open-air archaeological deposits. These issues are not specific to *Leiocephalus*, as recently
586 demonstrated by the similar complete absence of *Boa* snakes in regional archaeological sites, pattern
587 which has potentially been linked to specific Amerindian cultural practices (Bochaton 2020). A similar
588 lack of past occurrence data was also used to support an early Holocene extinction of *Diploglossus* in
589 the Lesser Antilles (Underwood 1964; although see Bochaton *et al.* 2016 for rare archaeological data
590 contradicting this position). It is possible that the small size and relative morphological similarity with
591 other lizard bones renders identifying *Leiocephalus* specimens from archaeological deposits difficult.
592 In this respect, it is interesting to note that rare archaeological taxa, such as *Leiocephalus* (Bailon *et al.*
593 2015; Boudadi-Maligne *et al.* 2016; Kemp & Hadly 2016; Pregill *et al.* 1994; Roughgarden & Pacala
594 1989; Stouvenot *et al.* 2014; Watters *et al.* 1984), *Boa* (Bochaton 2020; Bochaton & Bailon 2018), and
595 *Diploglossus* (Bochaton *et al.* 2016), were mentioned only by trained paleo-herpetologists. This lack

596 of archaeological data concerning *Leiocephalus* is thus not a good argument for their absence during
597 the Amerindian and historical periods. These lizards might have disappeared during the second half of
598 the 19th century colonial period, which would explain why they were overlooked by the biologists who
599 provided the first descriptions of fauna in the West Indies.

600 Past diversity and colonization history of *Leiocephalus* in the Lesser Antilles

601 The geological history of the West-Indies is complex as the geography of this area, especially its
602 southern part, has undergone massive changes across the last 50 million years. The West-Indies have
603 been mainly colonized in Late Eocene to early Oligocene times from South America (Blackburn *et al.*
604 2020; Delsuc *et al.* 2019) thanks to a land bridge linking the Greater Antilles to the continent
605 (Iturralde-Vinent 2006; Marivaux *et al.* 2020). This landmass however disappeared in the Late
606 Oligocene while a much younger chain of islands emerged eastward (the modern Lesser Antilles)
607 (Philippon *et al.* 2020). As an effect of this rich geological history several events can be drawn to
608 explain the colonization of the Lesser Antillean islands with overwater dispersal events from South
609 America, the Greater Antilles, or another still visible or nowadays disappeared island. Regarding
610 *Leiocephalus*, hypotheses of its initial colonization of the Lesser Antilles vary according to the
611 phylogenetic position of this genus. These lizards were first considered as “Tropiduridae” (Frost &
612 Etheridge 1989) native to South America, prior to being found to be closer to North American genera
613 (Hedges 1996). Recent molecular data indicates Leiocephalidae to be either a basal pleurodont
614 squamate (Pyron *et al.* 2013) or a sister taxon to all pleurodont squamates (Zheng & Wiens 2016),
615 making it difficult to pinpoint where they first appeared in the West Indies. Leiocephalidae and other
616 pleurodont squamates diverged 91 My years ago (Zheng & Wiens 2016), which would considerably
617 predate the K/T crisis and the divergence time of all Antillean taxa (Hedges 2006). Given the current
618 absence of molecular data for Lesser Antillean *Leiocephalus*, it is difficult to evaluate the reliability of
619 our morphology-based cladistic analysis. In addition, considering the changing position of
620 Leiocephalidae since the initial work of Pregill (1992), it is also possible that the character states
621 identified as “primitive” or “derived” need to be reevaluated in light of the past phylogenetic position of
622 *Leiocephalus*. This renders interpreting our results concerning the existence of a basal Lesser Antillean

623 clade more complex. Despite these uncertainties, if we accept the phylogenetic proximity between *L.*
624 *herminieri* and *L. roquetus*, and a basal position in their genus clade, our results would suggest an
625 initial colonization of the Cenozoic Antilles from South America just as it is the case for most other
626 terrestrial taxa. However, in the absence of molecular data for each modern species and fossil data
627 predating the Late Pleistocene for most of the West Indies, it is currently impossible to integrate this
628 potential scenario within what is likely to be very old and complex colonization that included multiple
629 extinction events. The history of *Leiocephalus* in the West Indies is probably significantly more
630 complex than the majority of modern taxa, rodents for example (Marivaux *et al.* 2020) which
631 colonized the region following either the formation of the Aves Ridge during the Oligocene or who
632 dispersed after the emergence of the modern islands (Hedges 2006).

633 **Conclusion**

634 Our results highlight the interest of revisiting old museum and private collections that, at first glance,
635 appear to hold no genuine scientific interest. The discovery of a dry specimen of an extinct taxon only
636 previously documented by fossil remains allowed us to describe an endemic extinct species, provide
637 additional details about the timing of its extinction, and advance a hypothesis concerning the past
638 radiation of the genus *Leiocephalus* in the Lesser Antilles. This exceptional discovery is likely due to
639 the important circulation of “exotic” goods from the Caribbean between the 17th and 19th centuries and
640 suggests that other interesting natural history specimens have ended up in private collections and then
641 in regional museums the world over. Like the example described here, these specimens are often of
642 uncertain origin and history. However, as we have shown, comparisons with fossil data of well-
643 documented origin can overcome these issues. The combination of fossil and modern data provide a
644 robust comparative framework for future studies of Caribbean *Leiocephalus* and reveals a wide
645 diversity of extinct lizards that likely remains to be described in the Lesser Antilles.

646

647 **Acknowledgments**

648 This study was funded by the FEDER program 2016–2020 “Ecosystème insulaire tropical, réponse de
649 la faune vertébrée terrestre à 6 000 ans d’anthropisation de la Guadeloupe (ECSIT) / Terrestrial
650 vertebrate responses to 6000 years of human activity in a tropical island ecosystem” of the CNRS and
651 a European PO-FEDER 2014-2020 grant (grant n°2016-FED-503), as well as support from the
652 Guadeloupe Regional Council, and the DAC of Guadeloupe.

653 The authors are extremely grateful to Nathalie Mémoire and the Natural History Museum of Bordeaux
654 for loaning us the dry historical specimen of *Leiocephalus*. We also thank Ronan Ledevin for his help
655 in the CT-scanning of the specimen and the two anonymous reviewers who provided feedback and
656 comments on this paper.

657

658 **References**

659 Bailon, S. (1991) Amphibiens et Reptiles du Pliocène et du Quaternaire de France et d’Espagne: Mise
660 en place et évolution des faunes. Ph. D Dissertation. Paris VII.

661 Bailon, S., Bochaton, C. & Lenoble, A. (2015) New data on Pleistocene and Holocene herpetofauna of
662 Marie-Galante (Blanchard Cave, Guadeloupe Islands, French West Indies): Insular faunal
663 turnover and human impact. *Quaternary Science Reviews* 128, 127–137.
664 <https://doi.org/10.1016/j.quascirev.2015.09.023>

665 Barnosky, A.D., Matzke, N., Tomiya, S., Wogan, G.O.U., Swartz, B., Quental, T.B., Marshall, C.,
666 McGuire, J.L., Lindsey, E.L., Maguire, K.C., Mersey, B. & Ferrer, E.A. (2011) Has the Earth’s
667 sixth mass extinction already arrived? *Nature* 471, 51–57.

668 Blackburn, D.C., Keeffe, R.M., Vallejo-Pareja, M.C. & Vélez-Juarbe, J. (2020) The earliest record of
669 Caribbean frogs: a fossil coquí from Puerto Rico. *Biology Letters* 16, 20190947.
670 <https://doi.org/10.1098/rsbl.2019.0947>

671 Bochaton, C. (2016) Squamates du Pléistocène supérieur et de l’Holocène de l’archipel
672 guadeloupéen : Évolution de la biodiversité et interactions avec l’Homme. PhD thesis.
673 Muséum national d’Histoire naturelle.

674 Bochaton, C. (2020) First records of modified snake bones in the Pre-Columbian archaeological
675 record of the Lesser Antilles: Cultural and paleoecological implications. *Journal of Island and*
676 *Coastal Archaeology*. <https://doi.org/10.1080/15564894.2020.1749195>

677 Bochaton, C. & Bailon, S. (2018) A new fossil species of *Boa* Linnaeus, 1758 (Squamata, Boidae) from
678 the Pleistocene of Marie-Galante Island (French West Indies). *Journal of Vertebrate*
679 *Paleontology* 38, e1462829. <https://doi.org/10.1080/02724634.2018.1462829>


- 680 Bochaton, C., Bailon, S., Herrel, A., Grouard, S., Ineich, I., Tresset, A. & Cornette, R. (2017) Human
681 impacts reduce morphological diversity in an insular species of lizard. *Proceedings of the*
682 *Royal Society B: Biological Sciences* 284, 20170921. <https://doi.org/10.1098/rspb.2017.0921>
- 683 Bochaton, C., Boistel, R., Casagrande, F., Grouard, S. & Bailon, S. (2016) A fossil *Diploglossus*
684 (Squamata, Anguidae) lizard from Basse-Terre and Grande-Terre islands (Guadeloupe, French
685 West-Indies). *Scientific Report* 28475, 1–12.
- 686 Bochaton, C., Boistel, R. & Charles, L. (2015a) X-ray microtomography provides first data about the
687 feeding behavior of an endangered lizard, the Monsterrat galliwasp (*Diploglossus*
688 *montisserrati*). *Open Science* 2, 150461.
- 689 Bochaton, C., Daza, J.D. & Lenoble, A. (2018) Identifying gecko species from Lesser Antillean
690 paleontological assemblages: Intraspecific osteological variation within and interspecific
691 osteological differences between *Thecadactylus rapicauda* (Houttuyn, 1782)
692 (Phyllodactylidae) and *Hemidactylus mabouia* (Moreau de Jonnès, 1818) (Gekkonidae).
693 *Journal of Herpetology* 52, 313–320. <https://doi.org/10.1670/17-093>
- 694 Bochaton, C., Grouard, S., Cornette, R., Ineich, I., Tresset, A. & Bailon, S. (2015b) Fossil and subfossil
695 herpetofauna from Cadet 2 Cave (Marie-Galante, Guadeloupe Islands, F. W. I.): Evolution of
696 an insular herpetofauna since the Late Pleistocene. *Comptes Rendus Palévol* 14, 101–110.
- 697 Bochaton, C., Ivanov, M. & Claude, J. (2019) Osteological criteria for the specific identification of
698 Monitor lizards (*Varanus* Merrem, 1820) remains in subfossil deposits of Sundaland and
699 continental Southeast Asia. *Amphibia-Reptilia* 40, 219–232.
700 <https://doi.org/10.1163/15685381-20181101>
- 701 Boudadi-Maligne, M., Bailon, S., Bochaton, C., Casagrande, F., Grouard, S., Serrand, N. & Lenoble, A.
702 (2016) Evidence for historical human-induced extinctions of vertebrate species on La
703 Désirade (French West Indies). *Quaternary Research* 85, 54–65.
- 704 Breuil, M. (2002) *Histoire naturelle des amphibiens et reptiles terrestres de l'archipel guadeloupéen:*
705 *Guadeloupe, Saint-Martin, Saint-Barthélemy*. Publications Scientifiques du Muséum National
706 d'Histoire Naturelle, France, 339 pp.
- 707 Breuil, M. (2009) The terrestrial herpetofauna of Martinique: past, present, future. *Applied*
708 *Herpetology* 6, 123–149.
- 709 Ceballos, G., Ehrlich, P.R., Barnosky, A.D., García, A., Pringle, R.M. & Palmer, T.M. (2015) Accelerated
710 modern human-induced species losses: Entering the sixth mass extinction. *Science Advances*
711 1, e1400253. <https://doi.org/10.1126/sciadv.1400253>
- 712 Ceballos, G., Ehrlich, P.R. & Dirzo, R. (2017) Biological annihilation via the ongoing sixth mass
713 extinction signaled by vertebrate population losses and declines. *Proceedings of the National*
714 *Academy of Sciences* 114, E6089–E6096. <https://doi.org/10.1073/pnas.1704949114>
- 715 Cooke, S.B., Dávalos, L.M., Mychajliw, A.M., Turvey, S.T. & Upham, N.S. (2017) Anthropogenic
716 extinction dominates Holocene declines of West Indian mammals. *Annual Review of Ecology,*
717 *Evolution, and Systematics* 48, 301–327. [https://doi.org/10.1146/annurev-ecolsys-110316-](https://doi.org/10.1146/annurev-ecolsys-110316-022754)
718 022754

- 719 Corke, D. (1992) The status and conservation needs of the terrestrial herpetofauna of the windward
720 islands (West Indies). *Biological Conservation* 62, 47–58. [https://doi.org/10.1016/0006-](https://doi.org/10.1016/0006-3207(92)91151-H)
721 3207(92)91151-H
- 722 Corry, E., Martin, L., Morton, M.N., Hilton, G.M., Young, R.P. & Garcia, G. (2010) *A Species Action Plan*
723 *for the Montserrat galliwasp* *Diploglossus montisserrati*. Department of Environment,
724 Montserrat.
- 725 Delsuc, F., Kuch, M., Gibb, G.C., Karpinski, E., Hackenberger, D., Szpak, P., Martínez, J.G., Mead, J.I.,
726 McDonald, H.G., MacPhee, R.D.E., Billet, G., Hautier, L. & Poinar, H.N. (2019) Ancient
727 Mitogenomes Reveal the Evolutionary History and Biogeography of Sloths. *Current Biology*
728 29, 2031–2042.e6. <https://doi.org/10.1016/j.cub.2019.05.043>
- 729 Du Tertre, J.-B. (R. P.) (1654) *Histoire générale des isles de S. Christophe, de la Guadeloupe, de la*
730 *Martinique, et autres dans l'Amérique*. A Paris : Chez Jacques Langlois et Emmanuel Langlois,
731 Paris, 542 pp. Available from: http://archive.org/details/cihm_34860 (May 6, 2014)
- 732 Duméril, A.M.C. & Bibron, G. (1837) *4 Erpétologie générale ou histoire naturelle complète des*
733 *reptiles*. Librairie encyclopédique de Roret, Paris, France, 573 pp.
- 734 Etheridge, R. (1964) Late Pleistocene lizards from Barbuda, British West Indies. *Bulletin of the Florida*
735 *State Museum* 9, 46–75.
- 736 Etheridge, R. (1966) The systematic relationship of West Indian and South American lizards referred
737 to the iguanid genus *Leiocephalus*. *Copeia* 1966, 79–91.
- 738 Evans, S.E. (2008) The skull of lizards and Tuatara. In: C. Gans, A. S. Gaunt, and K. Adler (Eds), *The*
739 *Skull of Lepidosauria*. Biology of the Reptilia. Society for the Study of Amphibians and
740 Reptiles, New-York, pp. 1–347.
- 741 Fejérvary-Langh, A.M. (1923) Beiträge zu einer Monographie der fossilen Ophisaurier. *Palaeontologia*
742 *Hungarica* 1, 123–220.
- 743 Fernández-Jalvo, Y. & Andrews, P. (2016) *Atlas of Taphonomic Identifications: 1001+ Images of Fossil*
744 *and Recent Mammal Bone Modification*. Springer, London, UK, 359 pp.
- 745 Fitzpatrick, S.M. (2015) The pre-columbian Caribbean: colonization, population dispersal, and island
746 adaptations. *PaleoAmerica* 1, 305–331. <https://doi.org/10.1179/2055557115Y.0000000010>
- 747 Frost, D.R. & Etheridge, R. (1989) A Phylogenetic Analysis and Taxonomy of Iguanian Lizards (Reptilia:
748 Squamata). *University of Kansas Museum of Natural History Miscellaneous Publication* 81, 1–
749 65.
- 750 Gray, J.E. (1827) A Description of a new Genus and some new Species of Saurian Reptiles, with a
751 Revision of the Species of Chameleons. *The Philosophical Magazine or Annals of Chemistry,*
752 *Mathematics, Astronomy, Natural History and General Science* 2, 207–214.
- 753 Hedges, S.B. (1996) The Origin of West Indian Amphibians and Reptiles. In: R. Powell and R. W.
754 Henderson (Eds), *Contributions to West Indian herpetology: a tribute to Albert Schwartz*.
755 Contributions to Herpetology. Society for the Study of Amphibians and Reptiles, New York
756 (USA), pp. 95–128.


- 757 Hedges, S.B. (2006) Paleogeography of the Antilles and origin of West Indian terrestrial vertebrates.
758 *Annals of the Missouri Botanical Garden* 93, 231–244.
- 759 Hedges, S.B. (2020) Caribherp: amphibian and reptiles of Caribbean islands. Available from:
760 <http://www.caribherp.org> (March 30, 2020)
- 761 Henderson, R. (1992) Consequences of predator introductions and habitat destruction on amphibians
762 and reptiles in the post-columbus West Indies. *Caribbean Journal of Science* 28, 1–10.
- 763 Henderson, R. & Powell, R. (2009) *Natural History of West Indian Reptiles and Amphibians*. University
764 Press of Florida, Gainesville, Florida, 496 pp.
- 765 Hofman, C.L., Hoogland, M.L.P. & Delpuech, A. (1999) New perspectives on a Huecan Saladoid
766 assemblage on Guadeloupe: the case of Morel I. In: *Archaeological investigations on St.*
767 *Martin (Lesser Antilles). The sites of Norman Estate, Anse des Pères and Hope Estate, with a*
768 *contribution to the “la Hueca problem.”* Archaeological Studies Leiden University. Leiden,
769 Netherlands, pp. 303–310.
- 770 Iturralde-Vinent, M.A. (2006) Meso-Cenozoic Caribbean paleogeography: implications for the
771 historical biogeography of the region. *International Geology Review* 48, 791–827.
772 <https://doi.org/10.2747/0020-6814.48.9.791>
- 773 Iverson, J.B. (1978) The impact of feral cats and dogs on populations of the West Indian rock iguana,
774 *Cyclura carinata*. *Biological Conservation* 14, 63–73.
- 775 Keegan, W.F., Hofman, C.L. & Ramos, R.R. (2013) *The Oxford Handbook of Caribbean Archaeology*.
776 Oxford University Press, Oxford, 617 pp.
- 777 Kemp, M.E. & Hadly, E.A. (2016) Early Holocene turnover, followed by stability, in a Caribbean lizard
778 assemblage. *Quaternary Research* 85, 255–261. <https://doi.org/10.1016/j.yqres.2015.12.007>
- 779 Klembara, J., Böhme, M. & Rummel, M. (2010) Revision of the anguine lizard *Pseudopus laurillardii*
780 (Squamata, Anguillidae) from the Miocene of Europe, with comments on paleoecology. *Journal*
781 *of Paleontology* 84, 159–196.
- 782 Köhler, G., Bobadilla, M.J.R. & Hedges, S.B. (2016) A new dune-dwelling lizard of the genus
783 *Leiocephalus* (Iguania, Leiocephalidae) from the Dominican Republic. *Zootaxa* 4121, 517.
784 <https://doi.org/10.11646/zootaxa.4121.5.2>
- 785 Lorvelec, O., Pascal, M., Pavis, C. & Feldmann, P. (2007) Amphibians and reptiles of the French West
786 Indies: Inventory, threats and conservation. *Applied Herpetology* 4, 131–161.
787 <https://doi.org/10.1163/157075407780681356>
- 788 Lugo, E., Schmidt, R. & Brown, R. (1981) Tropical forests in the Caribbean. *Ambio* 10, 318–324.
- 789 Mahler, D.L. & Kearney, M. (2006) The Palatal Dentition in Squamate Reptiles: Morphology,
790 Development, Attachment, and Replacement. *Fieldiana Zoology* 2006, 1–61.
791 [https://doi.org/10.3158/0015-0754\(2006\)108\[1:TPDISR\]2.0.CO;2](https://doi.org/10.3158/0015-0754(2006)108[1:TPDISR]2.0.CO;2)
- 792 Marivaux, L., Vélez-Juarbe, J., Merzeraud, G., Pujos, F., Viñola López, L.W., Boivin, M., Santos-
793 Mercado, H., Cruz, E.J., Grajales, A., Padilla, J., Vélez-Rosado, K.I., Philippon, M., Léticée, J.-L.,
794 Münch, P. & Antoine, P.-O. (2020) Early Oligocene chinchilloid caviomorphs from Puerto Rico

- 795 and the initial rodent colonization of the West Indies. *Proceedings of the Royal Society B:*
796 *Biological Sciences* 287, 20192806. <https://doi.org/10.1098/rspb.2019.2806>
- 797 Oelrich, T.M. (1956) The Anatomy of the Head of *Ctenosaura pectinata* (Iguanidae). *Miscellaneous*
798 *publications Museum of Zoology, University of Michigan* 94, 1–172.
- 799 Orihuela, J., Viñola, L.W., Vázquez, O.J., Mychajliw, A.M., de Lara, O.H., Lorenzo, L. & Soto-Centeno,
800 J.A. (2020) *Assessing the role of humans in Greater Antillean land vertebrate extinctions:*
801 *new insights from Cuba*. Paleontology. preprint.
- 802 Philippon, M., Cornée, J.-J., Münch, P., Hinsbergen, D.J.J. van, BouDagher-Fadel, M., Gailler, L.,
803 Boschman, L.M., Quillevère, F., Montheil, L., Gay, A., Lebrun, J.F., Lallemand, S., Marivaux, L.,
804 Antoine, P.-O. & Team, with the G. (2020) Eocene intra-plate shortening responsible for the
805 rise of a faunal pathway in the northeastern Caribbean realm. *PLOS ONE* 15, e0241000.
806 <https://doi.org/10.1371/journal.pone.0241000>
- 807 Pregill, G.K. (1992) Systematics of the west indian lizard genus *Leiocephalus* (Squamata: Iguania:
808 Tropicuridae). *Miscellaneous publication (University of Kansas. Museum of Natural History)*
809 84, 1–69.
- 810 Pregill, G.K. & Olson, S.L. (1981) Zoogeography of West Indian Vertebrates in Relation to Pleistocene
811 Climatic Cycles. *Annual Review of Ecology and Systematics* 12, 75–98.
- 812 Pregill, G.K., Steadman, D.W., Olson, S.L. & Grady, F.V. (1988) Late Holocene fossil vertebrates from
813 Burma Quarry, Antigua, Lesser Antilles. *Smithsonian contribution to zoology* 463, 1–27.
- 814 Pregill, G.K., Steadman, D.W. & Watters, D.R. (1994) Late Quaternary vertebrate faunas of the Lesser
815 Antilles: historical components of Caribbean biogeography. *Bulletin of Carnegie Museum of*
816 *Natural History* 30, 1–51.
- 817 Pyron, R.A., Burbrink, F.T. & Wiens, J.J. (2013) A phylogeny and revised classification of Squamata,
818 including 4161 species of lizards and snakes. *BMC Evolutionary Biology* 13, 1–53.
- 819 Rage, J.C. & Augé, M. (2010) Squamate reptiles from the middle Eocene of Lissieu (France). A
820 landmark in the middle Eocene of Europe. *Geobios* 43, 253–268.
- 821 de Rochefort, C.D. (1658) *Histoire naturelle et morale des Antilles de l'Amérique. Histoire générale*
822 *des Antilles habitées par les Français*. Arnould Liers. Rotterdam, 558 pp.
- 823 Roughgarden, J. & Pacala, S. (1989) Taxon cycle among *Anolis* lizard populations: review of evidence.
824 In: D. Otte and J. A. Jendler (Eds), *Speciation and it's consequences*. Sinauer Associates, Inc,
825 United States of America, pp. 403–432.
- 826 Smith, K.T. (2009) Eocene Lizards of the Clade Geiseltaliellus from Messel and Geiseltal, Germany,
827 and the Early Radiation of Iguanidae (Reptilia: Squamata). *Bulletin of the Peabody Museum of*
828 *Natural History* 50, 219–306. <https://doi.org/10.3374/014.050.0201>
- 829 Smith, K.T. (2011) The evolution of mid-latitude faunas during the Eocene: Late Eocene lizards of the
830 Medicine Pole Hills reconsidered. *Bulletin of the Peabody Museum of Natural History* 52, 3–
831 104.

- 832 Smith, K.T., Bhullar, B.-A.S., Köhler, G. & Habersetzer, J. (2018) The Only Known Jawed Vertebrate
 833 with Four Eyes and the Bauplan of the Pineal Complex. *Current Biology* 28, 1101-1107.e2.
 834 <https://doi.org/10.1016/j.cub.2018.02.021>
- 835 Soto-Centeno, J.A. & Steadman, D.W. (2015) Fossils reject climate change as the cause of extinction of
 836 Caribbean bats. *Scientific Reports* 5, 7971. <https://doi.org/10.1038/srep07971>
- 837 Steadman, D.W., Albury, N.A., Kakuk, B., Mead, J.I., Soto-Centeno, J.A., Singleton, H.M. & Franklin, J.
 838 (2015) Vertebrate community on an ice-age Caribbean island. *Proceedings of the National
 839 Academy of Sciences* 112, E5963–E5971. <https://doi.org/10.1073/pnas.1516490112>
- 840 Steadman, D.W., Pregill, G.K. & Olson, S.L. (1984) Fossil vertebrates from Antigua, Lesser Antilles:
 841 evidence for late Holocene human-caused extinctions in the West Indies. *Proceedings of the
 842 National Academy of Sciences of the United States of America* 81, 4448–4451.
- 843 Stouvenot, C., Grouard, S., Bailon, S., Bonnissent, D., Lenoble, A., Serrand, N. & Sierpe, V. (2014)
 844 L’abri sous roche Cadet 3 (Marie-Galante): un gisement à accumulations de faune et à
 845 vestiges archéologiques. In: B. Bérard and C. Losier (Eds), *Archéologie Caraïbe*. Taboui.
 846 Sidestone Press, Leiden, Netherlands, pp. 77–102.
- 847 Underwood, G. (1964) An anguid lizard from the Leeward islands. *Breviora* 200, 1–10.
- 848 Watters, D.R., Reitz, E.J., Steadman, D.W. & Pregill, G.K. (1984) Vertebrates from archaeological sites
 849 on Barbuda, West Indies. *Annals of Carnegie Museum* 53, 383–412.
- 850 Zheng, Y. & Wiens, J.J. (2016) Combining phylogenomic and supermatrix approaches, and a time-
 851 calibrated phylogeny for squamate reptiles (lizards and snakes) based on 52 genes and 4162
 852 species. *Molecular Phylogenetics and Evolution* 94, Part B, 537–547.
 853 <https://doi.org/10.1016/j.ympev.2015.10.009>




855 Figure 1: A) Map of the Lesser Antilles with the location of the Guadeloupe islands; B) Map of the
856 Guadeloupe islands with the location of the PGR6 deposit and indication of the -100 meters marine
857 isobath; C) Stratigraphic correspondence between the squares and *décapages* of the 2016 PGR6
858 excavation and those of the 2011 excavation with locations of the three 14c dates performed.



859

860 Figure 2: Dorsal (A) and ventral (B) views of the complete holotype specimen of *Leiocephalus*
861 *roquetus* sp. nov. (MHNBx 2002.1871). Scale bar= 5 mm.




862

863 Figure 3: Head and skull of the Holotype of *Leiocephalus roquetus* sp. nov. (MHNbX 2002.1871). A,

864 B: Picture and drawing of the dorsal scales of the head of the Holotype; C, D: Picture and drawing of

865 the lateral scales of the head of the Holotype; E, F: Dorsal and lateral views of the 3D model of the


866 skull of the holotype; G: Labial and lingual views of the 3D model of the right mandible of the
 867 holotype. Scale bars= 3 mm.



868


869 Figure 4: Premaxilla and maxilla of *Leiocephalus roquetus* sp. nov. A: 3D Model of the premaxilla of
 870 the holotype (MHNbx 2002.1871); B: Paratype “PGR6. square C11c. *décapage* 14.XII” collected in
 871 PGR6 deposit. C: 3D Model of the right maxilla of the holotype (MHNbx 2002.1871); D: Paratype
 872 “PGR6. square B10d. *décapage* 22.III” (right) collected in PGR6 deposit. Abbreviations: **c. i. p. p.**:
 873 crest of the internal premaxillary process, **e. p. p.**: external premaxillary process, **f. p.**: facial process, **f.**

874 **t. t.:** first most anterior tricuspid tooth, **i. m.:** imprint of the maxilla, **i. n.:** imprint of the nasal bone, **i.**
 875 **p.:** incisive process, **i. p. p.:** internal premaxillary process, **m. p.:** maxillary process, **n. p.:** nasal
 876 process, **p. f.:** premaxillary fenestra, **s. s.:** supradental shelf, **z. p.:** zygomatic process. Scale bars=
 877 2mm.



878
 879 Figure 5: Frontal and parietal bones of *Leiocephalus roquetus* sp. nov. A: 3D Model of the frontal
 880 bone of the holotype (MHNbx 2002.1871); B: Paratype frontal bone “PGR6. square B10d. *décapage*
 881 13.IV” collected in PGR6 deposit; C: Paratype frontal bone “PGR6. square B10d. *décapage* 13.V”


882 collected in PGR6 deposit; D: 3D Model of the parietal bone of the holotype (MHNbX 2002.1871); E:
 883 Paratype parietal bone “PGR6. square C11d. *décapage* 12.VI” collected in PGR6 deposit; F: Paratype
 884 parietal bone “PGR6. square B9b. *décapage* 22.VII” collected in PGR6 deposit. Abbreviations: **a. c.:**
 885 adductor crest, **c. c.:** cranial crest, **d. o.:** dermal ornamentation, **f-p. s.:** fronto-parietal suture, **i. n.:**
 886 imprint of the nasal bone, **i. pr.:** imprint of the prefrontal bone, **i. ps.:** imprint of the postfrontal bone,
 887 **i. s.:** imprint of the supratemporal bone, **n. c.:** nasal process crest, **n. f.:** nuchal fossa, **pa. f.:** parietal
 888 fossa, **p. d.:** postero-medial depression, **p. f.:** parietal foramen, **s. p.:** supratemporal process. Scale
 889 bars= 2mm.



890

891 Figure 6: Pterygoid, squamosal and quadrate bones of *Leiocephalus roquetus* sp. nov. A: 3D Model of
 892 the right pterygoid of the holotype (MHNbX 2002.1871); B: Paratype left pterygoid bone “PGR6.
 893 square C11d. *décapage* 14.X” collected PGR6 deposit; C: 3D Model of the right squamosal of the
 894 holotype (MHNbX 2002.1871); D: Paratype left squamosal bone “PGR6. square B9b. *décapage*


895 22.IX” collected in PGR6 deposit; E: 3D Model of the left quadrate of the holotype (MHNbx
896 2002.1871); F: Paratype left quadrate bone “PGR6. square C11c. *décapage* 15.XI” collected in PGR6
897 deposit. Abbreviations: **a.-d. d.**: antero dorsal depression, **a. e.**: articular surface with the ectopterygoid
898 bone, **a. p.**: articular surface with the palatine bone, **a. r.**: anterior ramus, **c.**: conch, **d. p.**: dorsal
899 process, **p. c.**: posterior crest, **p. e.**: pit for the insertion of the epipterygoid, **p. l.**: pterygoid lamina, **p.**
900 **t.**: pterygoid teeth?, **pa. p.**: palatine process, **po. p.**: posterior process, **t. c.**: tympanic crest, **t. p.**:
901 transverse process, **t. s.**: tooth socket, **v. p.**: ventral process. Scale bars= 2mm.



902

903 Figure 7: Dentary, articular bones and vertebrae of *Leiocephalus roquetus* sp. nov. A: 3D Model of the
 904 right dentary of the holotype (MHNbX 2002.1871); B: Paratype dentary “PGR6. square C11d.
 905 *décapage* 16.I” (left) collected in PGR6 deposit; C: Paratype dentary “PGR6. square C11c. *décapage*
 906 11.II” (left) collected in PGR6 deposit; D: 3D Model of the right articular of the holotype (MHNbX
 907 2002.1871); E: Paratype articular “PGR6. square B9b. *décapage* 17.VIII” (right) collected in PGR6

908 deposit; F: Anterior trunk vertebra collected in the layer 4 of PGR6 (square B10d, *décapage* 20); G:
909 Caudal vertebra collected in the layer 2/3 of PGR6 (square B10d, *décapage* 10). Abbreviations: **a. p.**:
910 angular process, **c. p.**: coronoid process, **f. t. t.**: first most anterior tricuspid tooth, **hy.**: hypapophysis, **i.**
911 **c.**: lateral imprint of the coronoid bone, **i. s.**: intra-mandibular septum, **m. s.**: medial spine, **n. s.**: neural
912 spine, **o. M. g.**: anterior opening of the Meckelian groove, **r. p.**: retroarticular process, **s.**: symphysis,
913 **sy.**: synapophysis, **s. p.**: surangular process. Scale bars=2 mm.



915 Figure 8: Consensus phylogenetic tree of modern and fossil Lesser Antillean species of *Leiocephalus*
916 showing the strong proximity between *L. roquetus* sp. nov. and *L. herminieri*. This tree was obtained
917 from the 177 most parsimonious trees (130 steps). In this tree, “ancestor ?” is the theoretical ancestor
918 used by Pregill (1992). The two other theoretical ancestors “ancestor A” and “ancestor B” are similar
919 to “ancestor ?” but for these “taxa” the characters recorded as “?” in the “ancestor” of Pregill (1992)
920 have been respectively replaced by “0” and “1” character states.

921

922 Appendix 1. Character matrix used in the phylogenetic analysis

923

924