



HAL
open science

Lost Manuscripts and Extinct Texts: A Dynamic Model of Cultural Transmission

Jean-Baptiste Camps, Julien Randon-Furling

► **To cite this version:**

Jean-Baptiste Camps, Julien Randon-Furling. Lost Manuscripts and Extinct Texts: A Dynamic Model of Cultural Transmission. CHR 2022: Computational Humanities Research Conference, Computational Humanities Research, Dec 2022, Antwerp, Belgium. halshs-03827975

HAL Id: halshs-03827975

<https://shs.hal.science/halshs-03827975>

Submitted on 26 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

Lost Manuscripts and Extinct Texts: A Dynamic Model of Cultural Transmission

Jean-Baptiste Camps^{1,2,*,†}, Julien Randon-Furling^{3,4,*,†}

¹Venice Center for Digital and Public Humanities, Univ. Ca' Foscari, Fondamenta Malcanton 5449, Venezia, 30123, Italy

²Centre Jean-Mabillon, École nationale des chartes, Paris Sciences & Lettres, 65 rue de Richelieu, Paris, 75002, France

³Centre Borelli, Univ. Paris-Saclay, ENS Paris-Saclay, CNRS, SSA, INSERM, 91190, Gif-sur-Yvette, France

⁴SAMM, FP2M (FR2036), Université Paris-1 Panthéon-Sorbonne, CNRS, Paris, 75013, France

Abstract

How did written works evolve, disappear or survive down through the ages? In this paper, we propose a unified, formal framework for two fundamental questions in the study of the transmission of texts: how much was lost or preserved from all works of the past, and why do their genealogies (their “phylogenetic trees”) present the very peculiar shapes that we observe or, more precisely, reconstruct? We argue here that these questions share similarities to those encountered in evolutionary biology, and can be described in terms of “genetic” drift and “natural” selection. Through agent-based models, we show that such properties as have been observed by philologists since the 1800s can be simulated, and confronted to data gathered for ancient and medieval texts across Europe, in order to obtain plausible estimations of the number of works and manuscripts that existed and were lost.

Keywords

Agent-based models, Stochastic models, Loss of cultural artefacts, Text transmission, Stemmatology

1. Introduction

How much do we preserve of the written knowledge, science [1] and culture of the past? And how representative is what we know compared to what existed? Such fundamental questions depend on the process through which texts were distributed materially.

Before the advent of the printing press, written texts were circulated in manuscript form. In order to make the text available, the author would dictate it to a secretary, or write a draft on wax tablets, papyrus, parchment or, eventually, paper, and this original, authorial, manuscript would then have to be copied manually by a scribe in the form of a new manuscript, and then circulated. Copies could then be used to create more manuscripts, again by manual copying, perhaps by other scribes in other regions at a later date. During this process, successive modifications were introduced in the text, either by error or intentionally, to make the text more suited to its intended audience. These alterations in the written sequence that forms the text could then be transmitted to its “descendants” by a given manuscript. Wear and tear, accidents, fashions caused

CHR 2022: Computational Humanities Research Conference, December 12 – 14, 2022, Antwerp, Belgium

*Corresponding author.

†These authors contributed equally.

✉ jean-baptiste.camps@chartes.psl.eu (J. Camps); julien.randon-furling@cantab.net (J. Randon-Furling)

🌐 <https://github.com/Jean-Baptiste-Camps/> (J. Camps)



© 2022 Copyright for this paper by its authors. Use permitted under Creative Commons License Attribution 4.0 International (CC BY 4.0).



CEUR Workshop Proceedings (CEUR-WS.org)

the destruction of some manuscripts, while others enjoyed the long life of library preservation. In the end, knowledge was lost and some texts went extinct, while other texts gained traction or were eventually preserved for future generations.

In textual studies and philology, since the development of the “common errors” methods from the 19th century onward [2, 3], the analysis of the alterations to the text allows the philologists to reconstruct the relations between the surviving copies (witnesses) of a given work, and to represent them as a tree-like graph, called a *stemma codicum* (fig. 1). This reconstructive process is not entirely different from the methods used by biologists to reconstruct the links between existing or extinct species, based on their shared characteristics that are supposed to derive from a common ancestry, and to represent it as a phylogenetic tree. Methods from this biological subfield, known as cladistics, have even sometimes been directly applied to texts, with controversial results [4, 5, 6, 7, 8]. These trees represent the result of an enquiry into the relationships between the surviving witnesses, together with the hypothetical lost nodes that can be deduced from them. To arrive at it, researchers have to study the variants – more precisely, common innovations or errors (i.e., mutations) – observed in the text of the surviving witnesses (fig. 1, A). The reconstructed tree will show only what can be deduced from surviving witnesses: the witnesses themselves, and as many hypothetical nodes as are needed to explain their relationships (fig. 1, B). While it may be the case with more recent works that all nodes are known and the graph represents the full transmission of the text – for instance with the genealogy of printed editions (fig. 1, C) –, most of the time the tree represents only a (potentially very small) subset of what existed (fig. 1, D).

A long standing observation, not yet fully understood, about the structure of the reconstructed trees was made by the French philologist Joseph Bédier, almost a century ago, in 1928 [13]. He observed that most trees reconstructed by philologists show a root bifurcation (a root with outdegree 2): in most reconstructions, the original (or archetype) had two, and only two, direct descendants, preventing an accurate reconstruction of the original text by majority principle (e.g., two witnesses vs one). Instead of searching an explanation for such data in the dynamics of text transmission, he interpreted this “forest of bifid trees” as the result of a methodological flaw or an unconscious bias, spurring a century long debate and causing a long lasting methodological schism inside textual scholarship that remains to be fully resolved [14, 15].

Bédier’s initial observation has been replicated, with estimates of the proportion of root bifurcation varying, from Bédier’s 95.5%, to somewhat lower estimates ranging from 70% to 83% [16, 17, 18]. Yet, some have argued that the prevalence of root bifurcation could be an explainable feature of manuscript transmission of texts as it reached us. Tentative explanations include combinatorial estimations of the proportion of root bifurcation for a given number of witnesses, under the assumption that all configurations are equally likely [19, 17, 20], or consider the effects of decimation (i.e., manuscript loss) [21], for instance by applying a uniform loss probability to static preexisting trees [10] or by calculating a node specific loss probability to simulated trees [22]. Even if it generated little follow-ups, there have also been rare attempts of using birth and death process for exploring the dynamics of manuscript transmission [23, 24].

The abundance of roots – and more generally nodes [18] – with out-degree 2, is not the only property that can be observed in many stemmas. The asymmetry between branches (cf. fig. 1, C and D), the presence or not of lateral transmission (generally called “contamination”; fig. 1, E) are other properties worthy of investigations, as well as those that can indicate that




Figure 1: From the real tree to the reconstruction. **A** an artificial example of the transmission of a very short text and its plausible reconstruction, where circles depict extant manuscripts, rectangles lost ones), and gray, nodes that are recoverable when reconstructing the tree; **B** another artificial example, where the distribution of lost manuscripts causes the loss of a whole branch, and where, consequently, the last common ancestor is not the original (root), but a lower witness (archetype); **C** the observed phylogeny of the successive editions of Fortunio, *Regole grammaticali della volgar lingua* (1516) [9, 10]; **D** the reconstructed phylogeny (*stemma*) of the Old French *Song of Roland* [11]; **E** the reconstructed phylogeny of the Anglo-French *Guy de Warewic* (Ewert, 1932), showing many cases of lateral transmission (“contamination”) in dashed lines, and even one instance of lateral transmission from outside the tree; **F** the same tree, but including (box shape) the outline of a lost branch (cf. [12]).

the tree made from extant witnesses represents only a small portion of the original tradition (i.e., lateral transmission from outside the tree; root identifiable not with the original but with a later manuscript; fig. 1, F). It is reasonable to assume that some of these properties reflect the dynamics of manuscript transmission, while others keep trace of important destruction, decimating manuscripts and removing even full branches. For the texts, this can be seen as an evolutionary process, where two antagonist tendencies are at work: the apparition of textual variants in individuals, causing the increase of diversity in the tradition, and the extinction of full branches, causing some variants to prevail upon others and so reducing diversity.

Here again, these observations can be put in perspective with problems occurring in evolutionary biology, where, too, two antagonists tendencies are at work, mutation and fixation, either by drift or natural selection, in a context where processes of speciation and extinction are strongly linked and where extant species represent only a small subset of the species that have existed [25]. In both cases, survival might be the exception and extinction the rule, be it by “bad genes or bad luck”, a process that can be seen in terms of gambler’s paradox [26, 27].

Basic processes of reproduction and destruction create complex shapes in the trees, from which one might want to deduce whether they can be fully explained by random process akin to genetic drift, or if differences in selective values are to be suspected. In other terms, going back to textual traditions, if cultural context, through literary taste, canon or fashion for instance, creates a form of evolutionary pressure on textual traditions. Any insights gained on this question would have an applicability beyond the question of the transmission of antique and medieval texts, because it seems that similar dynamics are at work in the diffusion of content in other medium, including print [10] or even the web [28], and have been observed in areas such as the cognitive evolution of scientific fields and the dynamics of scientific memes [29, 30].

Data on cases as different as the songs of the Medieval Occitan troubadours from southern France or the incunabula editions printed in Renaissance Italy outline the same Pareto-like world, where a large number of texts are kept only in a single or handful of documents, while a limited number of “successful” texts are kept in a large number of copies (fig. 2, **A** and **B**), where most authors are known only for one or two texts, while a very limited number of writers can have dozens of texts preserved (fig. 2, **C** and **D**)... Such a process is also apparent in the constitution of a literary canon of a limited number of authors and texts. This ‘canonisation’ can be seen has a progressive loss of diversity, where an ever shrinking number of authors and texts take on an ever growing share of the circulated documents (fig. 2, **E** and **F**). But is this due to chance or to a selective process ?

In fact, some properties as were just mentioned for textual traditions have some pendants in evolutionary models, concerning for instance the very unequal distribution of descendants [31], the varying patterns of biodiversity varying in time and space, studied in macroecology and biogeography [32, 33] and the dynamics of speciation and extinction that manifest themselves in the shape of phylogenies and the loss of branches from the tree of life [34, 25]. For this reason, there are inspiration and resources to be found in the study of mathematical properties of evolutionary trees, regarding the establishment of a null-model [35].

Can we gain some insight on what existed, what was lost, and the driving forces between extinction or survival of texts — by drift or selection? In order to do so, we need a better understanding of the dynamical processes of manuscript transmission.




Figure 2: A Pareto-like world with diminishing diversity. **A** distribution of witnesses per troubadour text; **B** distribution of the extant copies per work for incunabula printed in Italy; **C** distribution of texts per author for troubadours and **D** incunabula; **E** Shannon diversity, Generations as sites, texts as species, witnesses as individuals; **F** number of texts and witnesses per author.

2. An agent-based model of a stochastic process

In this paper we present a selection of the first results obtained with a stochastic model for the transmission of manuscripts in the Middle Ages. Following Weitzman [23, 24], we use so-called *birth-and-death* processes. These are random processes introduced in probability theory to describe, among other things, simple population dynamics and genealogies. For the simplest versions of these processes, it is possible to derive analytically (i.e. with mathematical formulae) certain quantities of interest: the expected number of individuals (here for us, manuscripts) still present at a time t , the extinction probability, the survival probability,... But also, quantities of particular interest in the context of manuscript genealogies: the probability that the latest

common ancestor (lca) be an archetype rather than the original, or the probability of root bifidity for the reconstructed stemma. Here we favour a numerical approach, that allows us to explore more complicated variants of a birth-and-death process through an agent-based computer simulation. The agents correspond to manuscripts, and during each time step of the simulation each agent has a probability λ of being copied (once) and a probability μ of disappearing. Following Cisne, we limit the increase rate of the population by letting the birth rate λ depend on the size of the population at the previous step k_{t-1} :

$$\lambda_t = \lambda \frac{K}{k_{t-1}}, \quad (1)$$

where K is a theoretical upper bound on the maximum possible size of the manuscript population extant at any given time during the period under consideration.

3. Phase diagrams obtained through computer simulations

We ran agent-based simulations of a Cisne-type tradition with a total time frame of 500 pseudo-years — that is, a time-step in the simulation corresponds roughly to 3 months. This was derived using an estimate for the time taken to produce an average 200 page manuscript (though the speed of scribes is known to vary a lot, from 1 to 10 leaves a day) [36]. We chose $K = 100\,000 = 10^5$, as an order of magnitude (rather than 10^4 or 10^6) for the total number of manuscripts in a given tradition that could be extant simultaneously at any one time¹.

As for the remaining free parameters, namely the base “birth” or copy rate, λ , and the “death” rate μ , we explored all possible pairs of values of these parameters within their range (from 0 to 1 in theory, but reduced here to 10^{-4} to 10^{-3} by field expertise, i.e. rough estimates from philological knowledge)². The space of all possible values is usually called the phase space in physics and other mathematical sciences, and a representation of the value taken by a given

¹This order of magnitude is a very rough estimate based on human population and our own assumptions: the medieval population of countries such as France or Italy was in the 10^7 range; we estimate that the maximum saturation of this market for a given book would be reached if around 1% of the population were to own a copy.

²Historical and philological knowledge of loss rates is very scarce and elusive, but can still be approached from various angles, such as the collection of data from ancient library catalogues, inventories, wills, as well as allusions and intertextuality [37, 38, 39]. Buringh [39] provides estimates for the Latin West, with a geometric mean of loss around -25% per century, with variations from -11% in the 9th to -32% in the 14th and 15th centuries (with local variations between medieval institutions from -3% to -71% per century). The global loss rate for non-illustrated manuscripts of several well known collections have been estimated around 93-97% [40, 41, 42, 43]. But there is a potential bias in accounting only for well known institutional collections, from which some manuscripts are known to have survived: trying to account for fully lost libraries, Buringh [39] is compelled to revise his estimates higher, to -25% by century until the 12th, up to -43% in the 15th. For incunabula, using editions whose original number of print made is known, it is possible to gather loss estimates by counting known surviving exemplars in public or private collections: doing so for Venetian incunabula, Trovato [44] finds very variable loss rates according to textual and material typology, from 73% for the *Decretales* printed on parchment to 99.3% for more popular chivalrous literature (*Orlando furioso* for instance). This shows the importance both of variation in time and space, and of textual contents and material typology. In some extreme cases, loss can be very close to 100%, for reasons that may combine the fragility of the document form, lack of consideration for the documents or large scale historical events such as political instability, invasions or major cultural changes; examples are provided by cases as different as the Merovingian royal diplomas on papyrus or the Lombard royal charters [45], the Mayan (pre-colombian) manuscripts or medieval notarial acts [46]. Production estimates have also been attempted on the basis of the

observable quantity (eg the extinction probability) when parameters are varied across the phase space is called the phase diagram of this quantity. We thus produced phase diagrams for a number of relevant observables (Fig. 3). Note that, since there is a stochastic component in the model, each phase diagram is computed by averaging over the results of a relatively large number of simulations, here 100 – this means that we produced 100 artificial manuscript traditions for each pair $(\lambda, \mu) \in [10^{-4}, 10^{-3}]$, varying values by increments of 10^{-4} ; hence each phase diagram required 10 000 simulations.

The approach then consists in identifying, within the phase diagrams (also called heat maps), regions in which the values for the observables are consistent either with measured quantities (as is done in the natural sciences) or with estimates for these quantities coming from other, independent and altogether different, models. We have circled in red such regions on the phase diagrams in Fig. 3.

These heat maps show that the results obtained through these simulations are internally consistent in terms of not only population size and survival rates, but also in terms of structural properties of the resulting trees. In particular, the results obtained for a ratio $\frac{\mu}{\lambda}$ between $\frac{5}{8}$ and $\frac{6}{7}$ are surprisingly consistent with the observed properties of some medieval traditions, in particular those from chivalric narratives in Old French. In particular, values of 0.55 for the survival of works and 0.05 for the survival of manuscripts (fig. 3), **A** and **B**, red squared area, bottom-right tile) are identical to those provided by Kestemont et al. for Old French chivalric romances, using unrelated methods from ecodiversity [49]. Yet, for what regards specifically Old French epics, known as *chanson de geste* – a genre predating the later form of the *roman*, and whose circulation and reception considerably differs for a long time –, the median final population of 2 and the third quartile of LCA outdegree of 2 (though, median LCA Shannon index is 0.69)³. This would lead us to revise Kestemont et al. estimates to 0.22 (instead of 0.55) for the specific survival of Old French epics (*chansons de geste*) and 0.01 (instead of 0.5) for the survival of epic manuscripts (a figure closer to that observed by Trovato [44] for their later Italian successors). On the other hand, specific values, this time, for (Arthurian and Antique matters) *romans* yield a third quartile of LCA outdegree of 3 more coherent with Kestemont

quantity of sealing wax acquired by a given producer (a chancellery for instance [47]). More founded loss estimates have also been gathered by counting how many of the acts mentioned in imperial or royal registers are kept in original or consigned in the archives of the recipients: this gives a loss rate of originals varying from 80% (acts from the emperor Charles IV in 1360-1361) to 90% for the acts from Louis X of France, increasing to 99% for the judgements rendered by his Parliament, suggesting here as well a massive effect of typological variation [46, 48], resulting in very strong biases in the body of documents available to us. For our simulation needs, if we start from Trovato's estimates (potentially more reliable, because based on editions whose original number of copies is known), we get a survival rate whose order of magnitude is between 0.1 and 0.01 (between 10^{-2} and 10^{-3}) in 500 years (2000 steps in our model), in similar ranges as Buringh's 0.75 in 100 years and those reported by Holtz and Canteaut; from this we can deduce a step loss rate for a given total survival rate. For instance, for 1% survival rate, $(1 - \mu)^{2000} = 0.01$, which simplifies to $\mu = 0.002$. So we retain values of μ between 10^{-4} to 10^{-3} . Given that books could not have been produced order of magnitudes higher or slower than they were destroyed (or we would be either drown in medieval manuscripts or keep none), we explore the same range for λ . Of course, fixed rates are a limitation, and do not yet account for substantial variations in time (such as massive extinction events, like, e.g., the fall of the Roman empire, the fire in Alexandria library, the shift from *volumen* to *codex* or from caroline to gothic script, etc.).

³Data about the traditions of the *chansons de geste* follow Vitale-Brovarone's [50] and Camps' [51]. Information on the shape of stemmata have been computed based on a restriction to *chansons de geste* and deduplicating of the collection provided by OpenStemmata [52, 53].

et al. general estimate (median tradition size, according to Martina [54], is 2 – and mean 4.8 – for the sole *romans en vers*, similarly to *chansons de geste*, but is expected to be higher for later *romans en prose*).




Figure 3: Heat maps (phase diagrams) for the simulation of Cisne-like models; first row contains population size and survival properties, namely **A** Survival rate of traditions (i.e., trees); **B** survival rate of manuscripts (i.e., nodes); **C** the median final population of surviving witnesses for traditions with at least one; second row contains data about the age of manuscripts in the simulation, for **D** all manuscripts ever created; **E** extant ones; as well as **F** the date of the oldest extant manuscript; the third row concerns structural properties of the trees themselves, **F** the median distance between the lowest common ancestor of the surviving manuscripts and the actual root of the original tree; **G** the third quartile of the LCA (archetype) out-degree and **H** and the median Shannon index for the families (the main branches stemming from the LCA). For each pair of parameter values between 0.0001 and 0.001, 100 simulations were run for 1000 active and 1000 inactive steps. Grey areas correspond to irrelevant values of the parameters, or unstable regions. The red square shows parameter regions where the observables computed on the simulated manuscript populations are consistent with observations made for Medieval French epics or with plausible estimates made using different methods.

The situation is, for the moment (and until further data is acquired) consistent yet deserving of further inquiry concerning the distribution of age of surviving manuscripts: in the simulation’s

red-squared area, the median date of surviving manuscripts would be in the 800's step (around 200 years after the original) and the median date of the oldest for each tradition in the 150-200 years range. For *chansons de geste*, the median date of surviving manuscripts would be between 1250 and 1300 [50, 51] – 150 to 200 years after the first documents of the genre itself (the end of the 11th century for the composition of the oldest version of the *Roland*). For *romans en vers*, it is, like the genre itself, slightly offset in time, with a peak between 1275 and 1325 [54].

4. Discussion

Combining previous inquiries by Weitzman and Cisne with the power of computer simulations and the methodology of statistical physics, we are able to reproduce the evolutionary process that underlies the observable data for, at least, some textual traditions such as those from medieval French epics and romances. The results obtained can even corroborate or refine results obtained by unrelated methodologies, such as those recently published by Kestemont et al. [49], indicating that these relatively simple birth-and-death process have relevancy in philology as well as they have in Evolutionary Biology for instance. This method then provides us a way to account both for population dynamics in time, loss or production estimates, as well as the shape of the stemmata (the phylogenies) of manuscripts, answering the century long Bédier observation [13], whose lack of solution until now has been at the core of a lasting schism in philological studies.

The range of further investigations opened by this research is considerably large. Models using individual variable rates of λ and μ could be used to account for phenomenon such as efforts of preservation of old and venerable artefacts, or higher selective values of some copies, or accelerated destruction due to small scale (e.g., burnt libraries), larger scale (e.g., the Dissolution of English monasteries, French Wars of Religion,...) or global events (e.g., shift in book types such as from *volumen* to *codex* or caroline to *gothic* scripts, major cultural changes like the Renaissance, ...). Modelling should also include the actual variation of the texts, to reflect the introduction of variants (mutations) in some families, and processes of transmission of inherited variants, as well as lateral transmission.

More generally, once having established this 'null model', deviations due to different factors should be explored, such as higher selective values for some mutations (*variants*) or individuals, fluctuations in time and space and the existence of different 'ecological niches' (e.g., the Anglo-Norman public versus the readers of Franco-Italian epics), typological variation in books or texts, chocks and bottlenecks, etc. The question of the age of surviving manuscripts should also be explored and accounted for, especially in the light of potential variations of λ and μ in time. For instance, the demand and rate of copy for a given text could be expected to be highest shortly after its initial release, when it is most fitted to the taste and fashion of the time, perhaps reinforce itself if the text gets a quick breakthrough, and then decrease with the passing of years. Similarly, the rate of destruction could vary at a global or local level, as some shocks lead to peaks of destruction or canonicalisation and conservation efforts lead to lower rates. Last but not least, the model should account for non standard transmission, in particular lateral transmission (contamination), a process not uncommon in textual transmission but that is also encountered in the natural world (e.g., lateral gene transfer).

The generality of the models considered here makes them applicable not only to medieval texts, but to any type of cultural transmission, at least in written form, from manuscript circulation to the elaboration of a canon of works. Further investigations should try to verify it on the broadest possible range of cases, starting with Western Medieval and Antique texts, but preferably encompassing cultural productions from very different time periods and continents.

References

- [1] J. L. Cisne, How science survived: Medieval manuscripts' "demography" and classic texts' extinction, *Science* 307 (2005-02-25) 1305–1307. URL: <http://science.sciencemag.org/content/307/5713/1305>. doi:10.1126/science.1104718.
- [2] S. Timpanaro, *La genesi del metodo del Lachmann*, 4th ed., UTET Libreria, Torino, 2003.
- [3] O. E. Haugen, The genealogical method, in: R. Philipp (Ed.), *Handbook of Stemmatology*, De Gruyter, 2020, pp. 57–138. URL: <https://doi.org/10.1515/9783110684384-003>.
- [4] A. C. Barbrook, C. J. Howe, N. Blake, P. Robinson, The phylogeny of the Canterbury Tales, *Nature* 394 (1998) 839–839.
- [5] C. J. Howe, A. C. Barbrook, M. Spencer, P. Robinson, B. Bordalejo, L. R. Mooney, Manuscript evolution, *Trends in Genetics* 17 (2001) 147–152. URL: <https://www.sciencedirect.com/science/article/pii/S0168952500022101>. doi:10.1016/S0168-9525(00)02210-1.
- [6] M. Spencer, E. A. Davidson, A. C. Barbrook, C. J. Howe, Phylogenetics of artificial manuscripts, *Journal of Theoretical Biology* 227 (2004) 503–511. URL: <https://www.sciencedirect.com/science/article/pii/S0022519303004442>. doi:10.1016/j.jtbi.2003.11.022.
- [7] C. Macé (Ed.), *The Evolution of Texts: confronting stemmatological and genetical methods. Proceedings of the international workshop held in Louvain-la Neuve on September 1-2, 2004*, Istituti editoriali e poligrafici internazionali, Pisa, 2006.
- [8] A. Hoenen, History of computer-assisted stemmatology, in: R. Philipp (Ed.), *Handbook of Stemmatology*, De Gruyter, 2020, pp. 294–303. URL: <https://www.degruyter.com/document/doi/10.1515/9783110684384/html>.
- [9] B. Richardson (Ed.), *Giovan Francesco Fortunio: Regole grammaticali della volgar lingua*, Antenore, Roma, 2001.
- [10] V. Guidi, P. Trovato, Sugli stemmi bipartiti. decimazione, asimmetria e calcolo delle probabilità, *Filologia italiana* 1 (2004) 9–48.
- [11] C. Segre (Ed.), *La chanson de Roland*, number 16 in *Documenti di filologia*, R. Ricciardi, Milan et Naples, 1971.
- [12] A. Hoenen, The stemma as a computational model, in: R. Philipp (Ed.), *Handbook of Stemmatology*, De Gruyter, 2020, pp. 226–241. URL: <https://www.degruyter.com/document/doi/10.1515/9783110684384/html>.
- [13] J. Bédier, La tradition manuscrite du lai de l'ombre. réflexions sur l'art d'éditer les anciens textes (premier article), *Romania* 54 (1928) 161–196.
- [14] C. A. Baker, M. Barbato, M. Cavagna, Y. Greub (Eds.), *L'ombre de Joseph Bédier: théorie et pratiques éditoriales au XXe siècle*, ÉLiPhi, Strasbourg, 2018.

- [15] F. Duval, *La 'Tradition manuscrite du Lai de l'Ombre' de Joseph Bédier ou la critique textuelle en question*, Honoré Champion, Paris, 2021.
- [16] W. P. Shepard, Recent theories of textual criticism, *Modern Philology* 28 (1930) 129–141. Publisher: University of Chicago Press.
- [17] A. Castellani, Bédier avait-il raison?: La méthode de Lachmann dans les éditions de textes du Moyen Age. Leçon inaugurale donnée à l'université de Fribourg le 2 juin 1954, number 20 in *Discours universitaires, Nouvelle série = Freiburger Universitätsreden*, Neue Folge, Éditions Universitaires, Fribourg, 1957.
- [18] O. E. Haugen, The silva portentosa of stemmatology: Bifurcation in the recension of Old Norse manuscripts, *Digital Scholarship in the Humanities* 31 (2015) 594–610. URL: <https://academic.oup.com/dsh/article/31/3/594/2340338>. doi:10.1093/dsh/fqv002.
- [19] P. Maas, Leitfehler und stemmatische Typen, *Byzantinische Zeitschrift* 37 (1937) 289–294. doi:10.1515/byzs.1937.37.2.289.
- [20] A. Hoenen, S. Eger, R. Gehrke, How many stemmata with root degree k ?, in: *Proceedings of the 15th Meeting on the Mathematics of Language*, 2017, pp. 11–21.
- [21] W. W. Greg, Recent theories of textual criticism, *Modern Philology* 28 (1931) 401–404. Publisher: University of Chicago Press.
- [22] A. Hoenen, Silva Portentissima – Computer-Assisted Reflections on Bifurcativity in Stemmas, in: *Digital Humanities 2016: Conference Abstracts*. Jagiellonian University & Pedagogical University, DH 2016, 2016, pp. 557–560. URL: <http://dh2016.adho.org/abstracts/311>.
- [23] M. P. Weitzman, Computer simulation of the development of manuscript traditions., *ALLC BULL.* 10 (1982) 55–59.
- [24] M. P. Weitzman, The Evolution of Manuscript Traditions, *Journal of the Royal Statistical Society. Series A (General)* 150 (1987) 287–308. URL: <http://www.jstor.org/stable/2982040>. doi:10.2307/2982040.
- [25] K. Yessoufou, T. J. Davies, Reconsidering the Loss of Evolutionary History: How Does Non-random Extinction Prune the Tree-of-Life?, in: R. Pellens, P. Grandcolas (Eds.), *Biodiversity Conservation and Phylogenetic Systematics: Preserving our evolutionary heritage in an extinction crisis*, *Topics in Biodiversity and Conservation*, Springer International Publishing, Cham, 2016, pp. 57–80. URL: https://doi.org/10.1007/978-3-319-22461-9_4. doi:10.1007/978-3-319-22461-9_4.
- [26] D. M. Raup, *Extinction: bad genes or bad luck?*, WW Norton & Company, 1992.
- [27] P. Canettieri, V. Loreto, M. Rovetta, G. Santini, *Philology and Information Theory*, *Cognitive Philology* 1 (2008). URL: <http://ojs.uniroma1.it/index.php/cogphil/article/view/8816>.
- [28] A.-L. Barabási, R. Albert, Emergence of Scaling in Random Networks, *Science* 286 (1999) 509–512. URL: <https://science.sciencemag.org/content/286/5439/509>. doi:10.1126/science.286.5439.509.
- [29] R. A. Bentley, Random Drift versus Selection in Academic Vocabulary: An Evolutionary Analysis of Published Keywords, *PLOS ONE* 3 (2008) e3057. URL: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0003057>. doi:10.1371/journal.pone.0003057.
- [30] D. Chavalarias, J.-P. Cointet, Phylomemetic Patterns in Science Evolution—The Rise and Fall of Scientific Fields, *PLOS ONE* 8 (2013) e54847. URL: <https://journals.plos.org/plosone/>

- article?id=10.1371/journal.pone.0054847. doi:10.1371/journal.pone.0054847.
- [31] J. Cosette, A. Moussy, F. Onodi, A. Auffret-Cariou, T. M. A. Neildez-Nguyen, A. Paldi, D. Stockholm, Single cell dynamics causes Pareto-like effect in stimulated T cell populations, *Scientific reports* 5 (2015) 1–10.
- [32] J. S. Cabral, L. Valente, F. Hartig, Mechanistic simulation models in macroecology and biogeography: state-of-art and prospects, *Ecography* 40 (2017) 267–280. URL: <https://onlinelibrary.wiley.com/doi/abs/10.1111/ecog.02480>. doi:<https://doi.org/10.1111/ecog.02480>.
- [33] T. F. Rangel, N. R. Edwards, P. B. Holden, J. A. F. Diniz-Filho, W. D. Gosling, M. T. P. Coelho, F. A. S. Cassemiro, C. Rahbek, R. K. Colwell, Modeling the ecology and evolution of biodiversity: Biogeographical cradles, museums, and graves, *Science* 361 (2018). URL: <https://science.sciencemag.org/content/361/6399/eaar5452>. doi:10.1126/science.aar5452.
- [34] G. M. Mace, J. L. Gittleman, A. Purvis, Preserving the Tree of Life, *Science* 300 (2003) 1707–1709. URL: <https://science.sciencemag.org/content/300/5626/1707>. doi:10.1126/science.1085510.
- [35] F. Bienvenu, F. Débarre, A. Lambert, The split-and-drift random graph, a null model for speciation, *Stochastic Processes and their Applications* 129 (2019) 2010–2048. URL: <https://www.sciencedirect.com/science/article/pii/S0304414918303004>. doi:10.1016/j.spa.2018.06.009.
- [36] E. Overgaauw, Fast or slow, professional or monastic. The writing speed of some late-medieval scribes, *Scriptorium* 49 (1995) 211–227. URL: https://www.persee.fr/doc/scrip_0036-9772_1995_num_49_2_1726. doi:10.3406/scrip.1995.1726, publisher: Persée - Portail des revues scientifiques en SHS.
- [37] Wilson R M, *The Lost Literature Of Medieval England*, Methuen, 1952. URL: <http://archive.org/details/in.ernet.dli.2015.86593>.
- [38] H. Bardon, *La littérature latine inconnue*, C. Klincksieck, Paris, 1952.
- [39] E. Buringh, *Medieval Manuscript Production in the Latin West*, Brill, 2010-11-19. URL: <http://booksandjournals.brillonline.com/content/books/9789047428640>. doi:10.1163/9789047428640.
- [40] M. Kestemont, F. Karsdorp, Estimating the Loss of Medieval Literature with an Unseen Species Model from Ecodiversity, in: *Proceedings of the Workshop on Computational Humanities Research*, volume 2723 of *CEUR*, 2020, pp. 44–55. URL: <http://ceur-ws.org/Vol-2723/short10.pdf>.
- [41] U. Neddermeyer, *Von der Handschrift zum gedruckten Buch: Schriftlichkeit und Leseinteresse im Mittelalter und in der frühen Neuzeit quantitative und qualitative Aspekte*, phdthesis, Philosophische Fakultät Köln, 1998.
- [42] H. Wijsman, *Luxury Bound: Illustrated Manuscript Production and Noble and Princely Book Ownership in the Burgundian Netherlands (1400-1550)*, volume 16 of *Burgundica*, Brepols Publishers, Turnhout, 2010. URL: <https://www.brepolonline.net/doi/book/10.1484/M.BURG-EB.5.105851>. doi:10.1484/M.BURG-EB.5.105851.
- [43] F. v. Oostrom, *Stemmen op schrift: geschiedenis van de Nederlandse literatuur vanaf het begin tot 1300*, Bert Bakker, Amsterdam, 2013.
- [44] P. Trovato, *Everything you always wanted to know about Lachmann's method: a non-standard handbook of genealogical textual criticism in the age of post-structuralism*,

- cladistics, and copy-text, Libreriauniversitaria.it edizioni, Limena, 2014. Préf. de Michael D. Reeve.
- [45] D. Ganz, W. Goffart, Charters Earlier than 800 from French Collections, *Speculum* 65 (1990) 906–932. URL: <https://www.journals.uchicago.edu/doi/abs/10.2307/2863567>. doi:10.2307/2863567.
- [46] E. Holtz, Überlieferungs- und Verlustquoten spätmittelalterlicher Herrscherurkunden, in: M. Lawo, O. B. Rader (Eds.), *Turbata per aequora mundi: Dankesgabe an Eckhard Müller-Mertens*, Harrassowitz, 2001, pp. 67–80.
- [47] R.-H. Bautier, Introduction, in: A. Lapeyre, R. Scheurer (Eds.), *Les notaires et secrétaires du roi sous les règnes de Louis XI, Charles VIII et Louis XII, 1461-1515*, volume 1, Bibliothèque nationale, Paris, 1978, pp. IX–XXXIX.
- [48] O. Canteaut, Quantifier l'activité des chancelleries à l'aune de la tradition des actes : l'exemple de la chancellerie des derniers Capétiens (1314-1328), in: O. Canteaut, O. Guyot-jeannin, O. Poncet (Eds.), *Actes royaux et princiers à l'ère du numérique (Moyen Âge-Temps modernes)*, Pau, 2020, pp. 103–114.
- [49] M. Kestemont, F. Karsdorp, E. de Bruijn, M. Driscoll, K. A. Kapitan, P. Ó Macháin, D. Sawyer, R. Sleiderink, A. Chao, Forgotten books: The application of unseen species models to the survival of culture, *Science* 375 (2022) 765–769. URL: <https://www.science.org/doi/abs/10.1126/science.abl7655>. doi:10.1126/science.abl7655, publisher: American Association for the Advancement of Science.
- [50] A. Vitale-Brovarone, La diffusion manuscrite des chansons de geste: une vue d'ensemble, in: E. Garavelli, M. Helkkula, O. Välikangas (Eds.), *Tra Italia e Francia. Entre France et Italie. In honorem Elina Suomela-Härmä*, number 69 in *Mémoire de la Société Néophilologique de Helsinki*, Helsinki, 2006, pp. 473–488.
- [51] J.-B. Camps, *La 'Chanson d'Otinél': édition complète du corpus manuscrit et prolégomènes à l'édition critique*, thèse de doctorat, dir. Dominique Boutet, Paris-Sorbonne, Paris, 2016. URL: <https://halshs.archives-ouvertes.fr/tel-01664932>. doi:10.5281/zenodo.1116735.
- [52] J.-B. Camps, S. Gabay, G. F. Riva, *Open Stemmata: A Digital Collection of Textual Genealogies*, in: *EADH2021: Interdisciplinary Perspectives on Data*, 2nd International Conference of the European Association for Digital Humanities, Krasnoyarsk, 2021. URL: <https://halshs.archives-ouvertes.fr/halshs-03260086>.
- [53] J.-B. Camps, G. Fernandez Riva, S. Gabay, *Open Stemmata: Database*, 2021. URL: <https://github.com/OpenStemmata/database/>, original-date: 2020-11-20T15:56:06Z.
- [54] P. A. Martina, *La produzione manoscritta del romanzo francese in versi : modelli materiali e modelli di cultura*, These de doctorat, Sorbonne université, 2018. URL: <https://www.theses.fr/2018SORUL051>.