HAL open science

Impact of uncertainty intolerance on clinical reasoning: a systematic review of the 21st century literature

Clémence Brun, Oulmann Zerhouni, Alexis Akinyemi, Laurène Houtin, Richard Monvoisin, Nicolas Pinsault

To cite this version:

Clémence Brun, Oulmann Zerhouni, Alexis Akinyemi, Laurène Houtin, Richard Monvoisin, et al.. Impact of uncertainty intolerance on clinical reasoning: a systematic review of the 21st century literature. 36th Annual European Health Psychology Society Conference 2022, Aug 2022, Bratislava, Slovakia. . ⟨halshs-03843560⟩

HAL Id: halshs-03843560

https://shs.hal.science/halshs-03843560v1

Submitted on 8 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Autorisation HAL

HAL Authorization

Impact of uncertainty intolerance on clinical reasoning: a scoping review of the 21st century literature

Clémence Brun[1,3], Oulmann Zerhouni[2], Alexis Akinyemi[2,3], Laurène Houtin[2,3], Richard Monvoisin[1], Nicolas Pinsault[1]
[1]Univ. Grenoble Alpes, TIMC-IMAG UMR CNRS 5525, ThEMAS team, [2]Univ. Paris Nanterre - Laboratoire Parisien de Psychologie Sociale, [3]AD-HOC Lab

INTRODUCTION

Choquet (2019, p. 18) defines **clinical reasoning** as "*the set of thought and decision-making processes that allow the clinician to determine the most appropriate actions in a specific context of health problem solving*". Among factors influencing clinical reasoning and thus medical decisions, uncertainty is currently extensively studied.

Intolerance of uncertainty (IU), *i.e.*, "*the tendency of an individual to consider as unacceptable the possibility that a negative event will occur, regardless of the probability of its occurrence*" (Carleton et al., 2007, p. 105), can have a significant impact on clinical reasoning.

Various studies show that a high level of **IU has deleterious effects on problem-solving abilities** and information processing, which can lead to inaction and avoidance of ambiguous situations (Begin et al., 2022).

However, studies on healthcare professionals' reactions to uncertainty and their impact on clinical reasoning are still rare.

This study aimed to **assess the current state of the literature regarding the influence of IU on the quality of clinical reasoning**. To answer this question, we conducted a scoping review.

METHOD

A **digital scoping review** was conducted by screening Google Scholar, Medline, PsycINFO and PBSC using English keywords identified using classifications (*e.g.*, MeSH, TermSciences terminology databases).

We used the following search equation: **(« clinical reasoning » OR diagnostic OR « intolerance of uncertainty ») AND (review OR definition OR introduction OR process* OR « decision making » OR uncertainty) OR (« clinical reasoning » AND diagnostic)**.

We included articles i) containing terms specified in our search equation, ii) talking about healthcare professionals as defined by the Public Health Code, and iii) reporting on the impact of uncertainty intolerance or uncertainty management on clinical reasoning. We excluded: i) articles with titles containing keywords but whose topic was incorrect, ii) articles published before 2000, iii) articles not written in English and iv) articles that were not freely available in PDF or readable format.

We selected articles by: a) removing duplicates, b) reading the titles and abstracts of the remaining articles saving only those relevant, c) reading full-text articles and excluding those which did not match our criteria and including the remaining articles to our review.

RESULTS

1. Practitioners' experience seems to influence their uncertainty management : increased **experience is related to increased comfort with uncertainty**.
2. The way **uncertainty is communicated to peers and patients impacts practitioners' well-being.**



3. Our analysis also revealed 2 categories of behaviours influenced by IU: investigative and prescriptive behaviours.

Regarding **investigative behaviours**, we found that IU could be associated with information withholding and reasoning bias, potentially resulting in diagnostic errors. IU was also associated with more referrals to peers and less use of new types of medical interventions.
Regarding **prescribing behaviours**, IU was associated with overtesting, which may be unnecessary or even dangerous. IUI was also associated with more prescriptions of antibiotics in situations where they should be prescribed with caution.
Finally, IU could influence abortion recommendations in cases of cystic fibrosis diagnosis in prenatal genetic tests.

DISCUSSION

Selected articles reported practical consequences, without mentioning cognitive impacts in detail. Future should focus more on the real-time psychological impact of IU on the different phases of clinical reasoning.

Also, a significant proportion of the articles that dealt with uncertainty in the health field **investigated the influence of uncertainty on patients' decision making rather than health professionals' decision making or how health professionals communicate their medical decisions and uncertainties** (Rising et al., 2018). Research seems currently focused on the impact of a hypothesized uncertainty, which should be more experimentally proven, on decision making.
Future studies should also examine these various concepts while recontextualizing them in decision support approaches, namely clinical decision support systems (CDSS, Harada et al., 2021). These systems are bound to grow in the future, especially since scientific studies have shown their interest for various conditions such as common chronic diseases. It is essential to work jointly on these aspects in order to ensure the development of programs that promote optimized diagnosis that are compatible with these new systems.

References :
- Begin, A. S., Hidrue, M., Lehrhoff, S., Del Carmen, M. G., Armstrong, K., & Wasfy, J. H. (2022). Factors associated with physician tolerance of uncertainty: An observational study. Journal of General Internal Medicine, 37(6), 1415–1421. https://doi.org/10.1007/s11606-021-06776-8
- Carleton, R. N., Norton, M. A. P. J., & Asmundson, G. J. G. (2007). Fearing the unknown: a short version of the Intolerance of Uncertainty Scale. Journal of Anxiety Disorders, 21(1), 105–117. https://doi.org/10.1016/j.janxdis.2006.03.014
- Choquet, G. (2019). Évaluation d'une formation sur l'incertitude par l'échelle de stress lié à l'incertitude (PRU) chez des internes en stage de médecine générale. Médecine humaine et pathologie. ffdumas-02498345f
- Harada, T., Miyagami, T., Kunitomo, K., & Shimizu, T. (2021). Clinical decision support systems for diagnosis in primary care: A scoping review. International Journal of Environmental Research and Public Health, 18(16), 8435. https://doi.org/10.3390/ijerph18168435
- Rising, K. L., Papanagnou, D., McCarthy, D., Gentsch, A., & Powell, R. (2018). Emergency medicine resident perceptions about the need for increased training in communicating diagnostic uncertainty. Cureus, 10(1), e2088. https://doi.org/10.7759/cureus.2088

adhoclab TIMC ehps 2022

Contact: clemence.brun@adhoclab.fr